



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 7, 2020 – 02:08 AM BST

PDB ID : 5U8Q
Title : Structure of the ectodomain of the human Type 1 insulin-like growth factor receptor in complex with IGF-I
Authors : Lawrence, M.; Xu, Y.
Deposited on : 2016-12-14
Resolution : 3.27 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1177 (3.32-3.24)
Clashscore	141614	1044 (3.30-3.26)
Ramachandran outliers	138981	1026 (3.30-3.26)
Sidechain outliers	138945	1025 (3.30-3.26)
RSRZ outliers	127900	1141 (3.32-3.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	885	<div> <div>7%</div> <div> <div></div> <div>74%</div> <div>16%</div> <div>9%</div> </div> </div>
2	B	70	<div> <div>4%</div> <div> <div></div> <div>61%</div> <div>7%</div> <div>31%</div> </div> </div>
3	H	126	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>12%</div> <div>6%</div> </div> </div>
4	L	108	<div> <div>4%</div> <div> <div></div> <div>80%</div> <div>18%</div> <div>..</div> </div> </div>
5	C	2	<div> <div></div> <div> <div></div> <div>50%</div> <div>50%</div> </div> </div>
5	E	2	<div> <div></div> <div> <div></div> <div>100%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	D	4	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NAG	C	2	-	-	-	X

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 8704 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Insulin-like growth factor 1 receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	801	Total	C	N	O	S	0	0	0
			6427	4066	1112	1202	47			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	THR	deletion	UNP P08069
A	?	-	THR	deletion	UNP P08069
A	?	-	MET	deletion	UNP P08069
A	?	-	SER	deletion	UNP P08069
A	?	-	SER	deletion	UNP P08069
A	?	-	ARG	deletion	UNP P08069
A	?	-	SER	deletion	UNP P08069
A	?	-	ARG	deletion	UNP P08069
A	?	-	ASN	deletion	UNP P08069
A	?	-	THR	deletion	UNP P08069
A	?	-	THR	deletion	UNP P08069
A	?	-	ALA	deletion	UNP P08069
A	?	-	ALA	deletion	UNP P08069
A	?	-	ASP	deletion	UNP P08069
A	?	-	THR	deletion	UNP P08069
A	?	-	TYR	deletion	UNP P08069
A	?	-	ASN	deletion	UNP P08069
A	?	-	ILE	deletion	UNP P08069
A	?	-	THR	deletion	UNP P08069
A	?	-	ASP	deletion	UNP P08069
A	738	ALA	PRO	conflict	UNP P08069
A	739	GLY	GLU	conflict	UNP P08069
A	740	ASN	GLU	conflict	UNP P08069
A	741	ASN	LEU	conflict	UNP P08069

- Molecule 2 is a protein called Insulin-like growth factor I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	48	Total	C	N	O	S	0	0	0
			375	235	63	70	7			

- Molecule 3 is a protein called Fv 24-60 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	H	118	Total	C	N	O	S	0	0	0
			915	581	150	181	3			

- Molecule 4 is a protein called Fv 24-60 light chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	L	107	Total	C	N	O	S	0	0	0
			820	511	140	166	3			

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	C	2	Total	C	N	O		0	0	0
			28	16	2	10				
5	E	2	Total	C	N	O		0	0	0
			28	16	2	10				

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	D	4	Total	C	N	O		0	0	0
			50	28	2	20				

- Molecule 7 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	S	0	0
			5	4	1		
7	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 8 is D-MALATE (three-letter code: MLT) (formula: $C_4H_6O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			9	4	5		

- Molecule 9 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:

C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	A	1	Total	C	N	O	0	0
			14	8	1	5		
9	A	1	Total	C	N	O	0	0
			14	8	1	5		
9	A	1	Total	C	N	O	0	0
			14	8	1	5		

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Chain A:
-
- 74% 7% 16% 9%
- ALA
GLY
ASN
ASN
GLU
THR
Y745
R751
V752
D753
V759
L763
L768
Y769
R770
I773
H774
H778
E779
S788
A798
E799
V808
L820
E824
P825
E826
I832
Y835
E836
I837
V850
Y855
K862
L863
N864
A873
A877
T878
V882
- ILE
GLU
GLU
VAL
THR
GLU
ASN
PRG
LYS
THR
GLU
VAL
CYS
GLY
GLY
GLU
LYS
GLY
PRG
CYS
CYS
ALA
CYS
PRG
LYS
THR
GLU
ALA
CYS
PRG
LYS
THR
GLU
ALA
GLN
ALA
GLU
LYS
GLU
ALA
GLU
LYS
GLU
ALA
GLU
GLU
TYR
R689
N694
H697
I700
P705
GLU
ARG
LYS
ARG
ARG
ARG
ASP
VAL
ASP
MET
GLN
VAL
ASN
ASP
- V353
K503
N504
V505
Q511
D512
A513
G514
G515
S518
L526
P527
G535
V551
I564
S569
Y573
S582
I583
P584
A590
S595
I598
P604
V616
R617
H618
Q621
D624
G625
Y626
R637
P638
R641
V642
T643
A644
T647
G648
G649
T650
K651
N652
K653
L654
I655
L656
L657
L658
L659
L660
L661
L662
L663
L664
L665
L666
L667
L668
L669
L670
L671
L672
L673
L674
L675
L676
L677
L678
L679
L680
L681
L682
L683
L684
L685
L686
L687
L688
L689
L690
L691
L692
L693
L694
L695
L696
L697
L698
L699
L700
L701
L702
L703
L704
L705
L706
L707
L708
L709
L710
L711
L712
L713
L714
L715
L716
L717
L718
L719
L720
L721
L722
L723
L724
L725
L726
L727
L728
L729
L730
L731
L732
L733
L734
L735
L736
L737
L738
L739
L740
L741
L742
L743
L744
L745
L746
L747
L748
L749
L750
L751
L752
L753
L754

- Chain B:
-
- | Amino Acid | Frequency (bits) |
|------------|------------------|
| GLY | 0.00 |
| PRO | 0.00 |
| GLU | 0.00 |
| T4 | 0.00 |
| L14 | 0.00 |
| D20 | 0.00 |
| R21 | 0.00 |
| G22 | 0.00 |
| P23 | 0.00 |
| N26 | 0.00 |
| LYS | 0.00 |
| PRO | 0.00 |
| THR | 0.00 |
| GLY | 0.00 |
| GLY | 0.00 |
| SER | 0.00 |
| SER | 0.00 |
| ARG | 0.00 |
| ARG | 0.00 |
| ALA | 0.00 |
| P39 | 0.00 |
| V44 | 0.00 |
| E48 | 0.00 |
| P63 | 0.00 |
| LEU | 0.00 |
| LYS | 0.00 |
| PRO | 0.00 |
| ALA | 0.00 |
| LYS | 0.00 |
| SER | 0.00 |
| ALA | 0.00 |

- Molecule 3: Fv 24-60 heavy chain




- Molecule 4: Fv 24-60 light chain



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 6: alpha-D-mannopyranose-(1-4)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	88.69 Å 197.66 Å 117.65 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	22.17 – 3.27 22.17 – 3.27	Depositor EDS
% Data completeness (in resolution range)	98.6 (22.17-3.27) 99.1 (22.17-3.27)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.20 (at 3.30 Å)	Xtriage
Refinement program	phenix.refine 1.11.1-2575_1692, PHENIX 1.11.1-2575_1692	Depositor
R, R_{free}	0.260 , 0.303 0.260 , 0.303	Depositor DCC
R_{free} test set	1524 reflections (4.72%)	wwPDB-V
Wilson B-factor (Å ²)	111.0	Xtriage
Anisotropy	0.718	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 97.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8704	wwPDB-V
Average B, all atoms (Å ²)	177.0	wwPDB-V

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, NAG, MLT, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.24	0/6582	0.44	0/8931
2	B	0.24	0/381	0.39	0/511
3	H	0.24	0/936	0.44	0/1270
4	L	0.25	0/838	0.47	0/1136
All	All	0.24	0/8737	0.44	0/11848

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6427	0	6220	81	0
2	B	375	0	347	5	0
3	H	915	0	888	10	0
4	L	820	0	792	13	0
5	C	28	0	25	1	0
5	E	28	0	25	0	0
6	D	50	0	43	0	0
7	A	10	0	0	0	0
8	A	9	0	4	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	A	42	0	39	0	0
All	All	8704	0	8383	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 106 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:511:GLN:H	1:A:515:GLY:HA3	1.58	0.69
1:A:114:GLU:HG2	1:A:115:LYS:HG3	1.75	0.68
1:A:433:GLU:HA	1:A:436:THR:HB	1.75	0.67
1:A:337:GLY:HA3	1:A:340:ILE:HD11	1.78	0.66
1:A:144:PRO:HG2	1:A:147:GLU:HB2	1.78	0.64

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	789/885 (89%)	723 (92%)	61 (8%)	5 (1%)	25	58
2	B	44/70 (63%)	40 (91%)	4 (9%)	0	100	100
3	H	116/126 (92%)	111 (96%)	5 (4%)	0	100	100
4	L	105/108 (97%)	99 (94%)	6 (6%)	0	100	100
All	All	1054/1189 (89%)	973 (92%)	76 (7%)	5 (0%)	29	62

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	286	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	11	ASN
1	A	379	LEU
1	A	380	GLY
1	A	504	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	715/787 (91%)	706 (99%)	9 (1%)	69	82
2	B	41/57 (72%)	41 (100%)	0	100	100
3	H	100/107 (94%)	100 (100%)	0	100	100
4	L	95/95 (100%)	93 (98%)	2 (2%)	53	75
All	All	951/1046 (91%)	940 (99%)	11 (1%)	71	83

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	310	THR
1	A	315	THR
1	A	626	TYR
1	A	294	CYS
1	A	624	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
4	L	90	GLN
4	L	92	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	NAG	C	1	1,5	14,14,15	0.28	0	17,19,21	0.45	0
5	NAG	C	2	5	14,14,15	0.27	0	17,19,21	0.52	0
6	NAG	D	1	1,6	14,14,15	0.46	0	17,19,21	0.38	0
6	NAG	D	2	6	14,14,15	0.29	0	17,19,21	0.45	0
6	MAN	D	3	6	11,11,12	0.95	0	15,15,17	1.17	2 (13%)
6	MAN	D	4	6	11,11,12	0.72	1 (9%)	15,15,17	1.21	2 (13%)
5	NAG	E	1	1,5	14,14,15	0.28	0	17,19,21	0.44	0
5	NAG	E	2	5	14,14,15	0.23	0	17,19,21	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	C	1	1,5	-	1/6/23/26	0/1/1/1
5	NAG	C	2	5	-	0/6/23/26	0/1/1/1
6	NAG	D	1	1,6	-	0/6/23/26	0/1/1/1
6	NAG	D	2	6	-	2/6/23/26	0/1/1/1
6	MAN	D	3	6	-	0/2/19/22	0/1/1/1
6	MAN	D	4	6	-	2/2/19/22	0/1/1/1
5	NAG	E	1	1,5	-	2/6/23/26	0/1/1/1
5	NAG	E	2	5	-	1/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	D	4	MAN	C1-C2	2.07	1.56	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	D	4	MAN	C1-O5-C5	3.57	117.03	112.19
6	D	3	MAN	C1-O5-C5	2.69	115.84	112.19
6	D	3	MAN	O2-C2-C3	-2.25	105.62	110.14
6	D	4	MAN	O2-C2-C3	-2.17	105.78	110.14

There are no chirality outliers.

5 of 8 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	E	1	NAG	O5-C5-C6-O6
5	E	1	NAG	C4-C5-C6-O6
6	D	4	MAN	O5-C5-C6-O6
6	D	4	MAN	C4-C5-C6-O6
6	D	2	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
9	NAG	A	1004	1	14,14,15	0.25	0	17,19,21	0.54	0
7	SO4	A	1001	-	4,4,4	0.14	0	6,6,6	0.05	0
7	SO4	A	1002	-	4,4,4	0.14	0	6,6,6	0.05	0
9	NAG	A	1007	1	14,14,15	0.40	0	17,19,21	0.57	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	MLT	A	1003	-	2,8,8	0.50	0	3,10,10	0.86	0
9	NAG	A	1012	1	14,14,15	0.31	0	17,19,21	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	NAG	A	1007	1	-	0/6/23/26	0/1/1/1
9	NAG	A	1004	1	-	1/6/23/26	0/1/1/1
9	NAG	A	1012	1	-	1/6/23/26	0/1/1/1
8	MLT	A	1003	-	-	2/2/8/8	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	A	1003	MLT	C1-C2-C3-C4
8	A	1003	MLT	O3-C2-C3-C4
9	A	1004	NAG	C3-C2-N2-C7
9	A	1012	NAG	C3-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	801/885 (90%)	0.42	65 (8%) 12 11	119, 174, 259, 316	0
2	B	48/70 (68%)	0.27	3 (6%) 20 20	140, 171, 205, 214	0
3	H	118/126 (93%)	0.16	4 (3%) 45 43	109, 147, 196, 222	0
4	L	107/108 (99%)	0.23	4 (3%) 41 39	102, 157, 202, 242	0
All	All	1074/1189 (90%)	0.37	76 (7%) 16 16	102, 170, 251, 316	0

The worst 5 of 76 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	171	TYR	7.1
1	A	233	ALA	6.4
1	A	146	LYS	5.8
1	A	799	GLU	5.6
1	A	512	ASP	5.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	C	2	14/15	0.77	0.47	161,231,239,242	0
6	MAN	D	4	11/12	0.81	0.59	189,225,236,236	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	MAN	D	3	11/12	0.83	0.45	219,242,253,261	0
6	NAG	D	1	14/15	0.84	0.18	136,164,184,214	0
5	NAG	C	1	14/15	0.87	0.20	177,211,224,225	0
6	NAG	D	2	14/15	0.87	0.34	168,202,218,239	0
5	NAG	E	1	14/15	0.90	0.27	119,153,169,179	0
5	NAG	E	2	14/15	0.91	0.37	150,177,196,200	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
9	NAG	A	1007	14/15	0.64	0.33	188,239,290,327	0
7	SO4	A	1002	5/5	0.74	0.36	172,185,208,221	0
9	NAG	A	1012	14/15	0.74	0.39	159,188,214,215	0
9	NAG	A	1004	14/15	0.81	0.61	172,207,225,227	0
7	SO4	A	1001	5/5	0.82	0.27	163,205,216,216	0
8	MLT	A	1003	9/9	0.83	0.31	137,153,160,161	0

6.5 Other polymers [i](#)

There are no such residues in this entry.