



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 04:55 pm BST

PDB ID : 4U9I
Title : High Resolution Structure Of The Ni-R State Of [Nife]Hydrogenase From Desulfovibrio Vulgaris Miyazaki F
Authors : Ogata, H.; Nishikawa, K.; Lubitz, W.
Deposited on : 2014-08-06
Resolution : 1.06 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1202 (1.10-1.02)
Clashscore	141614	1252 (1.10-1.02)
Ramachandran outliers	138981	1204 (1.10-1.02)
Sidechain outliers	138945	1202 (1.10-1.02)
RSRZ outliers	127900	1178 (1.10-1.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	S	265	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0;">%</div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 100%; background: linear-gradient(to right, red, orange, yellow, green);"></div> <div style="position: absolute; bottom: -10px; left: 0;">94%</div> <div style="position: absolute; bottom: -10px; right: 0;">6% •</div> </div> </div>
2	L	533	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 100%; background: linear-gradient(to right, red, orange, yellow, green);"></div> <div style="position: absolute; bottom: -10px; left: 0;">94%</div> <div style="position: absolute; bottom: -10px; right: 0;">5%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MPD	L	604	-	-	X	-
5	MPD	S	2007	-	-	-	X
8	TRS	L	605	-	X	-	-

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 13619 atoms, of which 6299 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Periplasmic [NiFe] hydrogenase small subunit.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	S	265	4139	1336	2026	355	404	18	0	13	0

- Molecule 2 is a protein called Periplasmic [NiFe] hydrogenase large subunit.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	L	533	8595	2765	4272	746	797	15	0	18	0

- Molecule 3 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Fe	S		
3	S	1	8	4	4	0	0
3	S	1	8	4	4	0	0

- Molecule 4 is FE3-S4 CLUSTER (three-letter code: F3S) (formula: Fe_3S_4).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	S	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 5 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $\text{C}_6\text{H}_{14}\text{O}_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	S	1	Total	C	O	0	0
			8	6	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	S	1	Total	C	O	0	0
			8	6	2		
5	S	1	Total	C	O	0	0
			8	6	2		
5	S	1	Total	C	O	0	0
			8	6	2		
5	S	1	Total	C	O	0	0
			8	6	2		
5	L	1	Total	C	O	0	0
			8	6	2		
5	L	1	Total	C	O	0	0
			8	6	2		

- Molecule 6 is hydrido[hydridonickel(2+)]bis(hydrocyanato-1kappaC)(hydroxymethyl)iron (three-letter code: NWN) (formula: $\text{C}_3\text{H}_4\text{FeN}_2\text{NiO}$).




Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
6	L	1	Total	C	Fe	H	N	Ni	O	0	0
			9	3	1	1	2	1	1		

- Molecule 7 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	L	1	Total	Mg	0	0
			1	1		

- Molecule 8 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	L	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	S	275	Total	O	0	0
			275	275		
9	L	505	Total	O	0	0
			505	505		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Periplasmic [NiFe] hydrogenase small subunit

Chain S: 



- Molecule 2: Periplasmic [NiFe] hydrogenase large subunit

Chain L: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.85Å 99.69Å 122.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.20 – 1.06 49.21 – 1.06	Depositor EDS
% Data completeness (in resolution range)	94.8 (49.20-1.06) 86.5 (49.21-1.06)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.10 (at 1.06Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1388)	Depositor
R, R_{free}	0.110 , 0.132 0.111 , 0.132	Depositor DCC
R_{free} test set	16569 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	8.2	Xtriage
Anisotropy	0.312	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 66.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	13619	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, SF4, MG, F3S, TRS, NWN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	S	0.70	5/2169 (0.2%)	0.71	1/2956 (0.0%)
2	L	0.74	15/4436 (0.3%)	0.74	3/6028 (0.0%)
All	All	0.73	20/6605 (0.3%)	0.73	4/8984 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	S	0	1

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	S	61	GLU	CD-OE1	12.77	1.39	1.25
2	L	470	GLU	CD-OE1	12.21	1.39	1.25
2	L	439[A]	GLU	CD-OE2	-11.70	1.12	1.25
2	L	439[B]	GLU	CD-OE2	-11.70	1.12	1.25
2	L	336	GLU	CD-OE1	-8.15	1.16	1.25
2	L	381	GLU	CD-OE2	8.12	1.34	1.25
2	L	433	GLU	CD-OE2	-7.07	1.17	1.25
2	L	271	GLU	CD-OE1	6.90	1.33	1.25
2	L	470	GLU	CD-OE2	6.61	1.32	1.25
2	L	418	GLU	CD-OE2	-6.40	1.18	1.25
2	L	34	GLU	CD-OE1	6.23	1.32	1.25
2	L	418	GLU	CD-OE1	6.03	1.32	1.25
1	S	209[A]	GLU	CD-OE2	-5.99	1.19	1.25
1	S	209[B]	GLU	CD-OE2	-5.99	1.19	1.25
1	S	57[A]	GLU	CD-OE2	-5.66	1.19	1.25
1	S	57[B]	GLU	CD-OE2	-5.66	1.19	1.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	L	464[A]	GLU	CD-OE1	-5.61	1.19	1.25
2	L	464[B]	GLU	CD-OE1	-5.61	1.19	1.25
2	L	345[A]	GLU	CD-OE1	-5.32	1.19	1.25
2	L	345[B]	GLU	CD-OE1	-5.32	1.19	1.25

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	L	59	ARG	NE-CZ-NH1	5.75	123.18	120.30
1	S	33	ASP	CB-CG-OD1	5.66	123.39	118.30
2	L	544	ASP	CB-CG-OD2	-5.13	113.68	118.30
2	L	32	ARG	NE-CZ-NH1	5.05	122.83	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	S	26	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	S	2113	2026	2011	13	0
2	L	4323	4272	4247	16	0
3	S	16	0	0	0	0
4	S	7	0	0	0	0
5	L	16	0	28	8	0
5	S	48	0	84	7	0
6	L	8	1	0	0	0
7	L	1	0	0	0	0
8	L	8	0	12	0	0
9	L	505	0	0	3	0
9	S	275	0	0	8	0
All	All	7320	6299	6382	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 3.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:S:2009:MPD:HM1	5:S:2009:MPD:H53	1.21	1.19
5:S:2009:MPD:C5	5:S:2009:MPD:HM1	2.03	0.87
2:L:360:LYS:HZ1	5:L:603:MPD:H11	1.45	0.82
5:S:2009:MPD:H53	5:S:2009:MPD:CM	2.10	0.81
5:S:2008:MPD:H51	9:L:825:HOH:O	1.81	0.79
1:S:96[B]:ASP:OD1	9:S:2101:HOH:O	2.03	0.76
2:L:166[A]:ASP:OD2	9:L:701:HOH:O	2.07	0.72
2:L:360:LYS:NZ	5:L:603:MPD:H11	2.10	0.66
2:L:417:PRO:HB3	5:L:604:MPD:H4	1.79	0.64
5:S:2009:MPD:C5	5:S:2009:MPD:CM	2.74	0.64
1:S:68:HIS:HA	5:S:2007:MPD:H4	1.80	0.63
2:L:381:GLU:OE2	2:L:514[B]:LYS:NZ	2.34	0.61
2:L:133:ALA:HA	2:L:136[B]:PHE:CZ	2.36	0.60
1:S:209[A]:GLU:OE2	9:S:2102:HOH:O	2.17	0.58
1:S:91:ASN:OD1	9:S:2103:HOH:O	2.18	0.56
1:S:135:ASP:OD1	1:S:138:LYS:NZ	2.40	0.55
1:S:61:GLU:OE1	1:S:65:ASN:ND2	2.34	0.53
2:L:417:PRO:CB	5:L:604:MPD:H4	2.38	0.53
1:S:134:ASN:HB3	5:S:2004:MPD:H4	1.90	0.52
5:L:604:MPD:H12	5:L:604:MPD:O4	2.12	0.49
2:L:163:THR:OG1	2:L:166[A]:ASP:OD2	2.26	0.49
1:S:96[B]:ASP:OD2	9:S:2104:HOH:O	2.19	0.48
5:L:604:MPD:O4	5:L:604:MPD:C1	2.61	0.48
1:S:266:GLN:NE2	9:S:2107:HOH:O	2.38	0.47
1:S:68:HIS:HD2	9:S:2349:HOH:O	1.97	0.47
2:L:395:TYR:CZ	5:L:604:MPD:H53	2.49	0.47
2:L:484:HIS:CD2	2:L:498[B]:LEU:HG	2.51	0.46
2:L:395:TYR:CE2	5:L:604:MPD:H53	2.51	0.45
2:L:439[B]:GLU:HG3	9:L:993:HOH:O	2.16	0.44
2:L:436:VAL:O	2:L:439[B]:GLU:HG3	2.17	0.44
1:S:96[A]:ASP:OD2	9:S:2105:HOH:O	2.21	0.44
1:S:241:TRP:CH2	1:S:243:VAL:HB	2.53	0.43
2:L:157:ILE:HD13	2:L:279:VAL:HG21	2.02	0.42
2:L:500:VAL:CG1	2:L:501:PRO:HD2	2.50	0.41
1:S:256[B]:ASP:OD1	9:S:2106:HOH:O	2.22	0.41
2:L:376:PRO:HD3	2:L:503:THR:HG22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	S	276/265 (104%)	270 (98%)	6 (2%)	0	100	100
2	L	549/533 (103%)	534 (97%)	15 (3%)	0	100	100
All	All	825/798 (103%)	804 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	S	224/211 (106%)	223 (100%)	1 (0%)	91	73
2	L	455/437 (104%)	454 (100%)	1 (0%)	93	76
All	All	679/648 (105%)	677 (100%)	2 (0%)	92	75

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	S	177	ASN
2	L	161	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	S	68	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 14 ligands modelled in this entry, 1 is monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	MPD	S	2009	-	7,7,7	0.51	0	9,10,10	0.99	0
6	NWN	L	601	2	1,6,6	7.19	1 (100%)	-		
5	MPD	L	603	-	7,7,7	0.43	0	9,10,10	0.52	0
5	MPD	S	2008	-	7,7,7	0.37	0	9,10,10	0.83	0
5	MPD	S	2006	-	7,7,7	0.49	0	9,10,10	0.51	0
5	MPD	L	604	-	7,7,7	0.30	0	9,10,10	0.78	0
8	TRS	L	605	-	7,7,7	0.29	0	9,9,9	1.20	2 (22%)
3	SF4	S	2001	1	0,12,12	0.00	-	-		
5	MPD	S	2005	-	7,7,7	0.27	0	9,10,10	0.53	0
3	SF4	S	2002	1	0,12,12	0.00	-	-		
5	MPD	S	2004	-	7,7,7	0.28	0	9,10,10	0.56	0
4	F3S	S	2003	1	0,9,9	0.00	-	-		
5	MPD	S	2007	-	7,7,7	0.47	0	9,10,10	1.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MPD	S	2009	-	-	1/5/5/5	-
5	MPD	L	603	-	-	3/5/5/5	-
5	MPD	S	2008	-	-	0/5/5/5	-
5	MPD	S	2006	-	-	0/5/5/5	-
5	MPD	L	604	-	-	4/5/5/5	-
8	TRS	L	605	-	-	8/9/9/9	-
3	SF4	S	2001	1	-	-	0/6/5/5
5	MPD	S	2005	-	-	0/5/5/5	-
3	SF4	S	2002	1	-	-	0/6/5/5
5	MPD	S	2004	-	-	0/5/5/5	-
4	F3S	S	2003	1	-	-	0/3/3/3
5	MPD	S	2007	-	-	0/5/5/5	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	L	601	NWN	O3-C3	-7.19	1.18	1.40

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	L	605	TRS	O2-C2-C	2.29	118.25	111.00
8	L	605	TRS	C2-C-N	-2.21	101.38	107.98

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	L	603	MPD	C1-C2-C3-C4
5	L	603	MPD	O2-C2-C3-C4
5	L	604	MPD	C1-C2-C3-C4
5	L	604	MPD	O2-C2-C3-C4
8	L	605	TRS	C1-C-C2-O2
8	L	605	TRS	C3-C-C2-O2
8	L	605	TRS	N-C-C2-O2
8	L	605	TRS	N-C-C3-O3
8	L	605	TRS	C2-C-C1-O1
8	L	605	TRS	C2-C-C3-O3
8	L	605	TRS	N-C-C1-O1
5	S	2009	MPD	C2-C3-C4-C5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	L	603	MPD	CM-C2-C3-C4
5	L	604	MPD	CM-C2-C3-C4
8	L	605	TRS	C3-C-C1-O1
5	L	604	MPD	C2-C3-C4-O4

There are no ring outliers.

6 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	S	2009	MPD	4	0
5	L	603	MPD	2	0
5	S	2008	MPD	1	0
5	L	604	MPD	6	0
5	S	2004	MPD	1	0
5	S	2007	MPD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	S	265/265 (100%)	-0.61	3 (1%) 80 75	6, 10, 21, 32	0
2	L	533/533 (100%)	-0.62	2 (0%) 92 89	6, 10, 19, 31	0
All	All	798/798 (100%)	-0.62	5 (0%) 89 86	6, 10, 20, 32	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	S	3	GLY	4.1
2	L	161	LYS	3.6
2	L	325	LYS	2.6
1	S	4	PRO	2.3
1	S	215	TRP	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	MPD	S	2007	8/8	0.46	0.44	23,27,30,35	0
5	MPD	L	603	8/8	0.68	0.22	23,30,33,33	0
8	TRS	L	605	8/8	0.70	0.31	23,25,30,33	0
5	MPD	S	2009	8/8	0.75	0.20	20,20,20,20	0
5	MPD	S	2008	8/8	0.79	0.28	25,27,30,32	0
5	MPD	S	2004	8/8	0.83	0.23	17,22,28,32	0
5	MPD	S	2005	8/8	0.83	0.22	23,27,30,30	0
5	MPD	L	604	8/8	0.86	0.16	24,26,28,35	0
5	MPD	S	2006	8/8	0.92	0.16	17,22,25,29	0
6	NWN	L	601	8/8	1.00	0.06	3,5,6,6	0
3	SF4	S	2002	8/8	1.00	0.02	8,8,9,9	0
7	MG	L	602	1/1	1.00	0.06	6,6,6,6	0
4	F3S	S	2003	7/7	1.00	0.03	6,6,6,7	0
3	SF4	S	2001	8/8	1.00	0.04	5,5,6,6	0

6.5 Other polymers [i](#)

There are no such residues in this entry.