



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 25, 2020 – 04:28 PM BST

PDB ID : 4UAA
Title : CTX-M-14 Class A Beta-Lactamase in Complex with a Non-Covalent Inhibitor
at Sub-Angstrom Resolution
Authors : Nichols, D.A.; Chen, Y.
Deposited on : 2014-08-08
Resolution : 0.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 0.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1071 (1.04-0.70)
Clashscore	141614	1143 (1.04-0.68)
Ramachandran outliers	138981	1065 (1.04-0.68)
Sidechain outliers	138945	1066 (1.04-0.68)
RSRZ outliers	127900	1038 (1.04-0.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	263	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0;">%</div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 100%; background: linear-gradient(to right, red 0%, red 1%, orange 1%, orange 87%, yellow 87%, yellow 98%, green 98%, green 100%);"></div> <div style="position: absolute; bottom: -10px; left: 0;">87%11%.</div> </div> </div>
1	B	263	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 100%; background: linear-gradient(to right, red 0%, red 1%, orange 1%, orange 86%, yellow 86%, yellow 97%, green 97%, green 100%);"></div> <div style="position: absolute; bottom: -10px; left: 0;">86%11%...</div> </div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9763 atoms, of which 4356 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-lactamase CTX-M-14.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	263	Total	C	H	N	O	S	0	34	0
			4451	1387	2211	410	435	8			
1	B	260	Total	C	H	N	O	S	0	31	0
			4334	1350	2145	398	433	8			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	25	PCA	-	insertion	UNP H6UQI0
B	25	PCA	-	insertion	UNP H6UQI0

- Molecule 2 is N-[3-(2H-tetrazol-5-yl)phenyl]-6-(trifluoromethyl)-1H-benzimidazole-4-carboxamide (three-letter code: 3GK) (formula: C₁₆H₁₀F₃N₇O).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			27	16	3	7	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			27	16	3	7	1		
2	B	1	Total	C	F	N	O	0	0
			27	16	3	7	1		
2	B	1	Total	C	F	N	O	0	0
			27	16	3	7	1		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	1
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	440	Total	O	0	36
			460	460		
4	B	370	Total	O	0	33
			390	390		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Beta-lactamase CTX-M-14



• Molecule 1: Beta-lactamase CTX-M-14



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	44.97Å 106.69Å 47.66Å 90.00° 102.08° 90.00°	Depositor
Resolution (Å)	10.00 – 0.86 46.61 – 0.86	Depositor EDS
% Data completeness (in resolution range)	98.2 (10.00-0.86) 91.9 (46.61-0.86)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.95 (at 0.86Å)	Xtriage
Refinement program	SHELXL	Depositor
R, R_{free}	0.121 , 0.134 0.125 , 0.136	Depositor DCC
R_{free} test set	3632 reflections (1.00%)	wwPDB-VP
Wilson B-factor (Å ²)	8.3	Xtriage
Anisotropy	0.086	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 50.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.99	EDS
Total number of atoms	9763	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, PCA, 3GK

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.68	0/2279	1.14	14/3094 (0.5%)
1	B	0.67	0/2221	1.28	30/3015 (1.0%)
All	All	0.68	0/4500	1.21	44/6109 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	3

There are no bond length outliers.

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	256	ARG	NE-CZ-NH2	11.48	126.04	120.30
1	B	222[A]	ARG	NE-CZ-NH1	10.07	125.34	120.30
1	B	222[B]	ARG	NE-CZ-NH1	10.07	125.34	120.30
1	A	184	ARG	NE-CZ-NH1	9.99	125.30	120.30
1	A	256[A]	ARG	NE-CZ-NH1	9.87	125.24	120.30
1	A	256[B]	ARG	NE-CZ-NH1	9.87	125.24	120.30
1	B	222[A]	ARG	CG-CD-NE	9.48	131.72	111.80
1	B	222[B]	ARG	CG-CD-NE	9.48	131.72	111.80
1	B	222[A]	ARG	CB-CG-CD	-9.25	87.55	111.60
1	B	222[B]	ARG	CB-CG-CD	-9.25	87.55	111.60
1	B	161	ARG	NE-CZ-NH2	-9.09	115.76	120.30
1	A	284	ARG	CD-NE-CZ	8.36	135.30	123.60
1	B	98[A]	LYS	CG-CD-CE	8.14	136.33	111.90
1	B	98[B]	LYS	CG-CD-CE	8.14	136.33	111.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	222[A]	ARG	NH1-CZ-NH2	-7.79	110.83	119.40
1	B	222[B]	ARG	NH1-CZ-NH2	-7.79	110.83	119.40
1	A	288[A]	GLU	CB-CG-CD	7.49	134.43	114.20
1	A	288[B]	GLU	CB-CG-CD	7.49	134.43	114.20
1	B	284	ARG	CD-NE-CZ	7.40	133.96	123.60
1	A	284	ARG	NE-CZ-NH1	-7.37	116.61	120.30
1	B	191	ARG	NE-CZ-NH2	-7.17	116.71	120.30
1	B	222[A]	ARG	NE-CZ-NH2	7.01	123.80	120.30
1	B	222[B]	ARG	NE-CZ-NH2	7.01	123.80	120.30
1	A	184	ARG	NE-CZ-NH2	-6.88	116.86	120.30
1	A	164	ARG	NE-CZ-NH2	-6.82	116.89	120.30
1	B	275	ARG	NE-CZ-NH1	6.56	123.58	120.30
1	B	153	ARG	NE-CZ-NH1	6.26	123.43	120.30
1	B	184	ARG	NE-CZ-NH1	6.09	123.35	120.30
1	B	288[A]	GLU	CA-CB-CG	6.04	126.69	113.40
1	B	288[B]	GLU	CA-CB-CG	6.04	126.69	113.40
1	A	158	GLU	OE1-CD-OE2	-5.97	116.13	123.30
1	B	240	ASP	CB-CG-OD2	-5.94	112.95	118.30
1	B	54[A]	ASN	CB-CA-C	5.90	122.20	110.40
1	B	54[B]	ASN	CB-CA-C	5.90	122.20	110.40
1	B	65	ARG	NE-CZ-NH1	5.78	123.19	120.30
1	B	55[A]	THR	OG1-CB-CG2	-5.43	97.52	110.00
1	B	55[B]	THR	OG1-CB-CG2	-5.43	97.52	110.00
1	A	189	THR	OG1-CB-CG2	5.25	122.07	110.00
1	A	59[A]	LEU	CA-CB-CG	5.21	127.28	115.30
1	A	59[B]	LEU	CA-CB-CG	5.21	127.28	115.30
1	B	284	ARG	NE-CZ-NH1	-5.13	117.74	120.30
1	B	201	GLU	OE1-CD-OE2	-5.04	117.26	123.30
1	B	264	TYR	CB-CG-CD1	5.04	124.02	121.00
1	A	52	ALA	CB-CA-C	-5.03	102.56	110.10

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	54[A]	ASN	Mainchain
1	B	54[B]	ASN	Mainchain
1	B	55[B]	THR	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2240	2211	2271	26	0
1	B	2189	2145	2191	18	0
2	A	54	0	20	0	0
2	B	54	0	20	0	0
3	A	5	0	0	0	0
3	B	15	0	0	0	0
4	A	460	0	0	16	0
4	B	390	0	0	7	0
All	All	5407	4356	4502	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:288[A]:GLU:HG3	4:A:578:HOH:O	1.74	0.88
1:B:38[B]:LYS:HD2	4:B:610:HOH:O	1.88	0.74
1:A:75[B]:MET:HG2	4:A:459:HOH:O	1.90	0.72
1:A:93[A]:GLN:HG2	4:A:401:HOH:O	1.89	0.72
1:B:255[B]:GLY:O	1:B:256:ARG:HD3	1.89	0.71
1:A:27:SER:H	1:A:30:GLN:HE21	1.40	0.67
1:B:186[A]:MET:HE1	4:B:478:HOH:O	1.92	0.67
1:B:88[A]:LYS:NZ	4:B:403:HOH:O	2.27	0.66
1:B:75[B]:MET:HG2	4:B:478:HOH:O	1.98	0.62
1:B:54[A]:ASN:O	1:B:55[A]:THR:C	2.39	0.61
1:B:38[B]:LYS:HG2	4:B:638:HOH:O	2.03	0.58
1:A:256[B]:ARG:HD3	4:A:554:HOH:O	2.08	0.54
1:A:98[A]:LYS:HE3	4:A:634:HOH:O	2.08	0.53
1:A:256[B]:ARG:HD3	4:A:598[B]:HOH:O	2.09	0.53
1:B:53[A]:ASP:CG	1:B:55[A]:THR:HG1	2.12	0.53
1:B:201:GLU:HG3	4:B:519[B]:HOH:O	2.10	0.51
1:A:93[A]:GLN:CG	4:A:404:HOH:O	2.59	0.51
1:A:93[A]:GLN:NE2	4:A:404:HOH:O	2.45	0.49
1:A:75[B]:MET:HG3	1:A:160:PHE:CE2	2.48	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:53[A]:ASP:O	1:B:54[A]:ASN:HB2	2.12	0.49
1:A:188[B]:GLN:HG2	4:A:506:HOH:O	2.14	0.48
1:B:92[B]:ASN:ND2	1:B:93[B]:GLN:OE1	2.47	0.48
1:A:222[B]:ARG:NH2	4:A:407:HOH:O	2.48	0.47
1:A:59[B]:LEU:HD21	4:A:626:HOH:O	2.15	0.47
1:A:96[A]:GLU:OE1	1:A:98[A]:LYS:NZ	2.48	0.47
1:A:106[B]:ASN:ND2	1:A:110[B]:GLU:OE2	2.48	0.47
1:A:270[B]:GLN:NE2	4:A:405:HOH:O	2.47	0.46
1:A:26:THR:HG22	1:A:31[B]:GLN:HG3	1.98	0.46
1:A:256[B]:ARG:HG2	4:A:554:HOH:O	2.15	0.45
1:A:93[A]:GLN:HG2	4:A:404:HOH:O	2.17	0.45
1:B:102:LEU:HD11	1:B:113[A]:VAL:HG21	1.98	0.45
1:B:50:ASP:O	1:B:54[A]:ASN:N	2.49	0.44
1:A:75[B]:MET:SD	1:A:189:THR:HG21	2.59	0.43
1:A:56:GLN:NE2	1:A:188[B]:GLN:OE1	2.52	0.42
1:B:247:ILE:HG22	1:B:262[B]:VAL:HG12	2.02	0.42
1:B:288[A]:GLU:HB2	4:B:411:HOH:O	2.18	0.42
1:B:53[A]:ASP:OD1	1:B:55[A]:THR:OG1	2.37	0.42
1:A:67:PRO:HD3	4:A:682[A]:HOH:O	2.20	0.42
1:A:95[A]:VAL:HG11	1:A:137:LYS:HD2	2.02	0.41
1:B:50:ASP:HB3	1:B:55[A]:THR:HB	2.03	0.41
1:B:96[A]:GLU:HG2	1:B:98[A]:LYS:HG3	2.03	0.41
1:A:93[A]:GLN:HB3	4:A:404:HOH:O	2.20	0.40
1:A:273:GLU:O	1:A:273:GLU:HG2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	295/263 (112%)	289 (98%)	5 (2%)	1 (0%)	41 13

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	289/263 (110%)	280 (97%)	6 (2%)	3 (1%)	15	2
All	All	584/526 (111%)	569 (97%)	11 (2%)	4 (1%)	25	3

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	55[A]	THR
1	B	55[B]	THR
1	A	103	VAL
1	B	103	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	238/205 (116%)	236 (99%)	2 (1%)	81	50
1	B	231/205 (113%)	227 (98%)	4 (2%)	60	25
All	All	469/410 (114%)	463 (99%)	6 (1%)	71	32

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	130	SER
1	A	192	GLN
1	B	130	SER
1	B	254[A]	GLN
1	B	254[B]	GLN
1	B	271	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	56	GLN
1	A	141	GLN
1	A	192	GLN
1	B	271	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PCA	A	25	1	7,8,9	1.28	1 (14%)	9,10,12	1.71	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	A	25	1	-	0/0/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	25	PCA	OE-CD	2.24	1.27	1.23

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	25	PCA	OE-CD-CG	-3.77	120.18	126.76
1	A	25	PCA	OE-CD-N	2.42	130.50	124.86
1	A	25	PCA	CB-CA-C	-2.05	109.88	112.70

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	3GK	A	301	-	28,30,30	0.88	2 (7%)	36,44,44	0.96	3 (8%)
3	PO4	B	305[B]	-	4,4,4	1.31	0	6,6,6	0.94	0
2	3GK	B	301	-	28,30,30	0.81	2 (7%)	36,44,44	1.05	2 (5%)
3	PO4	A	303	-	4,4,4	0.64	0	6,6,6	0.67	0
3	PO4	B	303	-	4,4,4	0.99	0	6,6,6	0.69	0
2	3GK	A	302	-	28,30,30	1.19	3 (10%)	36,44,44	1.51	5 (13%)
3	PO4	B	304	-	4,4,4	1.40	1 (25%)	6,6,6	1.36	0
2	3GK	B	302	-	28,30,30	1.25	4 (14%)	36,44,44	1.76	8 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	3GK	A	301	-	-	0/18/18/18	0/4/4/4
2	3GK	B	302	-	-	4/18/18/18	0/4/4/4
2	3GK	B	301	-	-	0/18/18/18	0/4/4/4
2	3GK	A	302	-	-	4/18/18/18	0/4/4/4

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	302	3GK	C15-C16	-3.39	1.38	1.43
2	B	302	3GK	N2-N1	2.92	1.39	1.34
2	A	301	3GK	C20-C15	2.91	1.43	1.37
2	A	302	3GK	C18-C17	-2.71	1.37	1.41
2	B	302	3GK	C3-N4	2.65	1.37	1.33
3	B	304	PO4	P-O1	2.51	1.56	1.50
2	A	302	3GK	N5-N1	2.42	1.36	1.32
2	B	302	3GK	C20-C19	2.36	1.43	1.39
2	A	301	3GK	C15-C16	-2.28	1.40	1.43
2	B	301	3GK	C15-C16	-2.18	1.40	1.43
2	B	302	3GK	C15-C16	-2.10	1.40	1.43
2	B	301	3GK	C3-N4	2.06	1.36	1.33

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	302	3GK	C10-C11-C6	4.60	126.34	120.56
2	B	302	3GK	O14-C13-C15	-4.10	116.06	121.72
2	A	302	3GK	C19-C18-C17	-3.86	115.16	120.67
2	A	302	3GK	C20-C15-C16	-3.73	114.74	118.21
2	B	302	3GK	C19-C18-C17	-3.43	115.77	120.67
2	B	301	3GK	C19-C18-C17	-3.23	116.05	120.67
2	B	302	3GK	C11-C6-C7	-3.17	113.68	118.16
2	A	301	3GK	C19-C18-C17	-3.09	116.26	120.67
2	B	302	3GK	C20-C15-C16	-3.07	115.36	118.21
2	A	302	3GK	N4-N5-N1	-2.91	107.63	109.53
2	A	301	3GK	C20-C15-C16	-2.62	115.77	118.21
2	B	301	3GK	N2-N1-N5	-2.49	107.91	109.53
2	A	302	3GK	C3-N2-N1	2.28	106.90	104.87
2	B	302	3GK	N2-N1-N5	-2.19	108.10	109.53
2	A	302	3GK	C18-C19-C20	2.15	122.72	118.69
2	B	302	3GK	C6-C3-N4	2.15	127.76	124.12
2	A	301	3GK	C18-C19-C20	2.10	122.61	118.69
2	B	302	3GK	C7-C6-C3	2.07	123.09	120.05

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	302	3GK	O14-C13-C15-C16
2	A	302	3GK	O14-C13-C15-C20
2	A	302	3GK	N12-C13-C15-C16
2	B	302	3GK	O14-C13-C15-C16
2	A	302	3GK	N12-C13-C15-C20
2	B	302	3GK	N12-C13-C15-C20
2	B	302	3GK	N12-C13-C15-C16
2	B	302	3GK	O14-C13-C15-C20

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand 3GK A 301




Ligand 3GK B 301



Ligand 3GK A 302



Ligand 3GK B 302



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	262/263 (99%)	-0.63	2 (0%) 86 72	6, 9, 17, 25	0
1	B	260/263 (98%)	-0.63	1 (0%) 92 80	6, 9, 18, 27	0
All	All	522/526 (99%)	-0.63	3 (0%) 89 75	6, 9, 18, 27	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	54[A]	ASN	3.0
1	A	88[A]	LYS	2.2
1	A	254[A]	GLN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PCA	A	25	8/9	0.96	0.08	14,18,21,32	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	PO4	B	304	5/5	0.68	0.28	17,17,18,19	5
3	PO4	B	305[B]	5/5	0.86	0.22	14,23,28,29	5
2	3GK	B	302	27/27	0.92	0.08	11,13,14,17	27
3	PO4	A	303	5/5	0.93	0.14	17,17,21,21	5
3	PO4	B	303	5/5	0.94	0.12	14,15,17,22	5
2	3GK	A	302	27/27	0.95	0.08	8,9,13,15	27
2	3GK	B	301	27/27	0.99	0.04	6,8,11,14	0
2	3GK	A	301	27/27	0.99	0.04	6,7,10,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around 3GK A 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 3GK B 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.