



wwPDB X-ray Structure Validation Summary Report ⓘ

May 26, 2020 – 04:49 pm BST

PDB ID : 1UBL
Title : Three-dimensional Structure of The Carbon Monoxide Complex of [NiFe]hydrogenase From Desulfovibrio vulgaris Miyazaki F
Authors : Ogata, H.; Mizoguchi, Y.; Mizuno, N.; Miki, K.; Adachi, S.; Yasuoka, N.; Yagi, T.; Yamauchi, O.; Hirota, S.; Higuchi, Y.
Deposited on : 2003-04-04
Resolution : 1.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1286 (1.22-1.18)
Ramachandran outliers	138981	1240 (1.22-1.18)
Sidechain outliers	138945	1239 (1.22-1.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	S	267	
2	L	534	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MPD	S	2004	-	-	X	-

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 7116 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Periplasmic [NiFe] hydrogenase Small subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	S	267	Total	C	N	O	S	0	0	0
			2019	1282	342	378	17			


- Molecule 2 is a protein called Periplasmic [NiFe] hydrogenase Large subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	534	Total	C	N	O	S	0	0	0
			4177	2674	725	763	15			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
L	514	LYS	ASN	SEE REMARK 999	UNP P21852
L	515	LEU	VAL	SEE REMARK 999	UNP P21852

- Molecule 3 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	S	1	Total	Fe	S	0	0
			8	4	4		
3	S	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 4 is FE3-S4 CLUSTER (three-letter code: F3S) (formula: Fe_3S_4).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	S	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 5 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $\text{C}_6\text{H}_{14}\text{O}_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	S	1	Total 8	C 6	O 2	0	0
5	S	1	Total 8	C 6	O 2	0	0
5	S	1	Total 8	C 6	O 2	0	0
5	L	1	Total 8	C 6	O 2	0	0
5	L	1	Total 8	C 6	O 2	0	0

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	L	1	Total Mg 1 1	0	0

- Molecule 7 is (MU-SULPHIDO)-BIS(MU-CYS,S)-[TRICARBONYLIRON-DI-(CYS,S)NICKEL(II)](FE-NI) (three-letter code: FNE) (formula: $\text{C}_3\text{FeNiO}_3\text{S}$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	L	1	Total	C	Fe	Ni	O	0	0
			8	3	1	1	3		

- Molecule 8 is CARBON MONOXIDE (three-letter code: CMO) (formula: CO).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	L	1	Total	C O	0	0
			2	1 1		

- Molecule 9 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	S	293	Total 293	O 293	0	0
9	L	553	Total 553	O 553	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Periplasmic [NiFe] hydrogenase Small subunit



- Molecule 2: Periplasmic [NiFe] hydrogenase Large subunit



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	97.81 Å 125.50 Å 66.59 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.20	Depositor
% Data completeness (in resolution range)	89.1 (20.00-1.20)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS, SHELXL-97	Depositor
R, R_{free}	0.117 , 0.161	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	7116	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CMO, MPD, SF4, MG, FNE, F3S

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	S	0.76	0/2075	1.33	23/2830 (0.8%)
2	L	0.74	0/4288	1.28	36/5831 (0.6%)
All	All	0.75	0/6363	1.30	59/8661 (0.7%)

There are no bond length outliers.

The worst 5 of 59 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	S	5	ARG	NE-CZ-NH1	-13.97	113.32	120.30
2	L	506	LEU	CA-CB-CG	11.79	142.41	115.30
2	L	71	ARG	NE-CZ-NH2	-9.74	115.43	120.30
2	L	369	ARG	NE-CZ-NH1	9.34	124.97	120.30
1	S	5	ARG	CD-NE-CZ	-9.24	110.66	123.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	S	2019	0	1949	58	0
2	L	4177	0	4129	59	0
3	S	16	0	0	0	0
4	S	7	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	L	16	0	28	5	0
5	S	24	0	42	9	0
6	L	1	0	0	0	0
7	L	8	0	0	0	0
8	L	2	0	0	0	0
9	L	553	0	0	12	0
9	S	293	0	0	9	0
All	All	7116	0	6148	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 108 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:S:146:ASN:HD21	5:S:2004:MPD:H13	1.26	0.99
1:S:2:MET:HA	2:L:182:GLN:HG2	1.46	0.98
1:S:126:ASN:HD21	1:S:130:ALA:H	1.17	0.91
1:S:238:GLN:HE21	2:L:224:ARG:HH21	1.17	0.91
2:L:470:GLU:HG2	2:L:487:ARG:HD3	1.57	0.86

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	S	265/267 (99%)	258 (97%)	5 (2%)	2 (1%)	19	3
2	L	532/534 (100%)	518 (97%)	14 (3%)	0	100	100
All	All	797/801 (100%)	776 (97%)	19 (2%)	2 (0%)	41	16

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	S	4	PRO
1	S	5	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	S	213/213 (100%)	198 (93%)	15 (7%)	15	1
2	L	438/438 (100%)	415 (95%)	23 (5%)	22	2
All	All	651/651 (100%)	613 (94%)	38 (6%)	20	1

5 of 38 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	L	49	LYS
2	L	144	LYS
2	L	445	LEU
2	L	132	HIS
2	L	152	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
2	L	78	GLN
2	L	188	ASN
2	L	457	ASN
2	L	132	HIS
2	L	205	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	FNE	L	1004	8,2	3,7,9	2.37	2 (66%)	-		
3	SF4	S	1001	1	0,12,12	0.00	-	-		
3	SF4	S	1002	1	0,12,12	0.00	-	-		
5	MPD	S	2001	-	7,7,7	0.56	0	9,10,10	1.29	1 (11%)
8	CMO	L	1006	7	0,1,1	0.00	-	-		
5	MPD	S	2004	-	7,7,7	0.74	0	9,10,10	0.96	1 (11%)
5	MPD	L	2010	-	7,7,7	0.45	0	9,10,10	0.79	0
5	MPD	L	2006	-	7,7,7	0.52	0	9,10,10	0.56	0
4	F3S	S	1003	1	0,9,9	0.00	-	-		
5	MPD	S	2007	-	7,7,7	0.51	0	9,10,10	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MPD	S	2004	-	-	0/5/5/5	-
3	SF4	S	1002	1	-	-	0/6/5/5
5	MPD	S	2001	-	-	0/5/5/5	-
3	SF4	S	1001	1	-	-	0/6/5/5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MPD	L	2010	-	-	2/5/5/5	-
5	MPD	L	2006	-	-	4/5/5/5	-
4	F3S	S	1003	1	-	-	0/3/3/3
5	MPD	S	2007	-	-	0/5/5/5	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	L	1004	FNE	O3-C3	-3.21	1.11	1.16
7	L	1004	FNE	O1-C1	2.25	1.20	1.16

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	S	2001	MPD	CM-C2-C1	3.20	117.24	110.57
5	S	2004	MPD	O2-C2-CM	2.37	115.68	108.08

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	L	2006	MPD	O2-C2-C3-C4
5	L	2006	MPD	C1-C2-C3-C4
5	L	2006	MPD	CM-C2-C3-C4
5	L	2010	MPD	C2-C3-C4-C5
5	L	2010	MPD	C2-C3-C4-O4

There are no ring outliers.

4 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	S	2004	MPD	8	0
5	L	2010	MPD	1	0
5	L	2006	MPD	4	0
5	S	2007	MPD	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.