



wwPDB X-ray Structure Validation Summary Report ⓘ

May 28, 2020 – 08:44 pm BST

PDB ID : 1UD0
Title : CRYSTAL STRUCTURE OF THE C-TERMINAL 10-kDA SUBDOMAIN OF HSC70
Authors : Chou, C.C.; Forouhar, F.; Yeh, Y.H.; Wang, C.; Hsiao, C.D.
Deposited on : 2003-04-24
Resolution : 3.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1291 (3.52-3.40)
Clashscore	141614	1372 (3.52-3.40)
Ramachandran outliers	138981	1337 (3.52-3.40)
Sidechain outliers	138945	1338 (3.52-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	113	
1	B	113	
1	C	113	
1	D	113	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2705 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 70 kDa heat-shock-like protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	84	Total	C	N	O	S	Se	0	0	0
			672	417	114	136	2	3			
1	B	82	Total	C	N	O	S	Se	0	0	0
			664	413	112	134	2	3			
1	C	88	Total	C	N	O	S	Se	0	0	0
			702	438	118	140	2	4			
1	D	78	Total	C	N	O	S	Se	0	0	0
			641	399	108	130	2	2			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	534	LEU	-	CLONING ARTIFACT	UNP P63018
A	535	VAL	-	CLONING ARTIFACT	UNP P63018
A	536	PRO	-	CLONING ARTIFACT	UNP P63018
A	537	ARG	-	CLONING ARTIFACT	UNP P63018
A	538	GLY	-	CLONING ARTIFACT	UNP P63018
A	539	SER	-	CLONING ARTIFACT	UNP P63018
A	540	HIS	-	CLONING ARTIFACT	UNP P63018
A	541	MSE	-	CLONING ARTIFACT	UNP P63018
A	549	MSE	MET	MODIFIED RESIDUE	UNP P63018
A	617	MSE	MET	MODIFIED RESIDUE	UNP P63018
A	621	MSE	MET	MODIFIED RESIDUE	UNP P63018
B	534	LEU	-	CLONING ARTIFACT	UNP P63018
B	535	VAL	-	CLONING ARTIFACT	UNP P63018
B	536	PRO	-	CLONING ARTIFACT	UNP P63018
B	537	ARG	-	CLONING ARTIFACT	UNP P63018
B	538	GLY	-	CLONING ARTIFACT	UNP P63018
B	539	SER	-	CLONING ARTIFACT	UNP P63018
B	540	HIS	-	CLONING ARTIFACT	UNP P63018
B	541	MSE	-	CLONING ARTIFACT	UNP P63018
B	549	MSE	MET	MODIFIED RESIDUE	UNP P63018
B	617	MSE	MET	MODIFIED RESIDUE	UNP P63018

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	621	MSE	MET	MODIFIED RESIDUE	UNP P63018
C	534	LEU	-	CLONING ARTIFACT	UNP P63018
C	535	VAL	-	CLONING ARTIFACT	UNP P63018
C	536	PRO	-	CLONING ARTIFACT	UNP P63018
C	537	ARG	-	CLONING ARTIFACT	UNP P63018
C	538	GLY	-	CLONING ARTIFACT	UNP P63018
C	539	SER	-	CLONING ARTIFACT	UNP P63018
C	540	HIS	-	CLONING ARTIFACT	UNP P63018
C	541	MSE	-	CLONING ARTIFACT	UNP P63018
C	549	MSE	MET	MODIFIED RESIDUE	UNP P63018
C	617	MSE	MET	MODIFIED RESIDUE	UNP P63018
C	621	MSE	MET	MODIFIED RESIDUE	UNP P63018
D	534	LEU	-	CLONING ARTIFACT	UNP P63018
D	535	VAL	-	CLONING ARTIFACT	UNP P63018
D	536	PRO	-	CLONING ARTIFACT	UNP P63018
D	537	ARG	-	CLONING ARTIFACT	UNP P63018
D	538	GLY	-	CLONING ARTIFACT	UNP P63018
D	539	SER	-	CLONING ARTIFACT	UNP P63018
D	540	HIS	-	CLONING ARTIFACT	UNP P63018
D	541	MSE	-	CLONING ARTIFACT	UNP P63018
D	549	MSE	MET	MODIFIED RESIDUE	UNP P63018
D	617	MSE	MET	MODIFIED RESIDUE	UNP P63018
D	621	MSE	MET	MODIFIED RESIDUE	UNP P63018

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Na 1 1	0	0
2	D	2	Total Na 2 2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	8	Total O 8 8	0	0
3	B	7	Total O 7 7	0	0
3	C	5	Total O 5 5	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	3	Total	O	0	0
			3	3		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 70 kDa heat-shock-like protein




- Molecule 1: 70 kDa heat-shock-like protein



- Molecule 1: 70 kDa heat-shock-like protein



- Molecule 1: 70 kDa heat-shock-like protein



E600	A601	V602	C603	N604	P605	I606	I607	I608	A614	GLY	GLY	MSE	PRO	GLY	GLY	MSE	PRO	GLY	PHE	PRO	GLY	GLY	GLY	ALA	PRO	PRO	SER	GLY	GLY	ALA	SER	SER	GLY	PRO	THR	ILE	GLU	GLU	VAL	ASP
------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	117.48 Å 117.48 Å 163.78 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	14.93 – 3.45 14.92 – 3.45	Depositor EDS
% Data completeness (in resolution range)	90.3 (14.93-3.45) 99.0 (14.92-3.45)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.30 (at 3.48 Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.249 , 0.309 0.305 , 0.350	Depositor DCC
R_{free} test set	804 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	63.0	Xtriage
Anisotropy	0.261	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.24 , 22.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	2705	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/678	0.69	0/900
1	B	0.49	0/670	0.69	0/890
1	C	0.51	0/708	0.65	0/940
1	D	0.62	1/647 (0.2%)	0.71	1/861 (0.1%)
All	All	0.53	1/2703 (0.0%)	0.68	1/3591 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	544	SER	C-N	7.58	1.51	1.34

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	544	SER	O-C-N	-5.38	114.08	122.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	672	0	659	65	0
1	B	664	0	653	90	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	702	0	695	62	0
1	D	641	0	631	66	1
2	B	1	0	0	0	0
2	D	2	0	0	0	0
3	A	8	0	0	0	0
3	B	7	0	0	0	0
3	C	5	0	0	0	0
3	D	3	0	0	0	0
All	All	2705	0	2638	247	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 46.

The worst 5 of 247 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:537:ARG:HG2	1:D:538:GLY:H	1.15	1.08
1:B:617:MSE:H	1:B:618:PRO:HD2	1.18	1.05
1:B:562:ILE:HB	1:B:566:ASP:HB3	1.48	0.93
1:A:558:LEU:HD23	1:A:562:ILE:HB	1.53	0.91
1:A:549:MSE:HE3	1:A:549:MSE:HA	1.54	0.89

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:554:GLU:OE2	1:D:554:GLU:OE2[7_556]	1.50	0.70

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	82/113 (73%)	54 (66%)	17 (21%)	11 (13%)	0	3
1	B	80/113 (71%)	49 (61%)	23 (29%)	8 (10%)	0	6
1	C	86/113 (76%)	61 (71%)	20 (23%)	5 (6%)	1	15
1	D	76/113 (67%)	51 (67%)	20 (26%)	5 (7%)	1	12
All	All	324/452 (72%)	215 (66%)	80 (25%)	29 (9%)	1	7

5 of 29 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	618	PRO
1	B	559	GLN
1	B	563	ASN
1	B	617	MSE
1	D	539	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	74/89 (83%)	68 (92%)	6 (8%)	11	40
1	B	74/89 (83%)	71 (96%)	3 (4%)	30	62
1	C	78/89 (88%)	68 (87%)	10 (13%)	4	20
1	D	72/89 (81%)	68 (94%)	4 (6%)	21	53
All	All	298/356 (84%)	275 (92%)	23 (8%)	13	42

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	555	ASP
1	C	581	LEU
1	D	603	CYS
1	C	563	ASN
1	C	582	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	612	GLN
1	C	604	ASN
1	D	584	ASN
1	B	604	ASN
1	D	563	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.