



# Full wwPDB X-ray Structure Validation Report ⓘ

May 24, 2020 – 01:17 pm BST

PDB ID : 5UDK  
Title : IFIT1 monomeric mutant (L457E/L464E) with PPP-AAAA  
Authors : Abbas, Y.M.; Martinez-Montero, S.; Damha, M.J.; Nagar, B.  
Deposited on : 2016-12-27  
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)
RNA backbone	3102	1011 (2.36-0.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	479	<div> <div>6%</div> <div>92%</div> <div>.</div> <div>.</div> </div>
2	B	4	<div> <div>50%</div> <div>50%</div> </div>

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 8268 atoms, of which 3960 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interferon-induced protein with tetratricopeptide repeats 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	461	Total	C	H	N	O	S	0	15	0
			7690	2433	3857	668	710	22			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP P09914
A	457	GLU	LEU	engineered mutation	UNP P09914
A	464	GLU	LEU	engineered mutation	UNP P09914

- Molecule 2 is a RNA chain called RNA (5'-D\*(ATP))-R(P\*AP\*AP\*A)-3').


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	4	Total	C	H	N	O	P	0	0	0
			142	40	45	20	31	6			

- Molecule 3 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			24	6	14	4		
3	A	1	Total	C	H	O	0	1
			48	12	28	8		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			17	4	10	3		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Ca	0	0
			2	2		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	315	Total	O	0	0
			315	315		
7	B	20	Total	O	0	0
			20	20		



- Molecule 1: Interferon-induced protein with tetratricopeptide repeats 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 42 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	112.43 Å 112.43 Å 93.17 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.25 – 1.65 44.25 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.7 (44.25-1.65) 99.7 (44.25-1.65)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.01 (at 1.65 Å)	Xtriage
Refinement program	PHENIX 1.10_2142	Depositor
R, $R_{free}$	0.164 , 0.184 0.166 , 0.185	Depositor DCC
$R_{free}$ test set	3560 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.5	Xtriage
Anisotropy	0.369	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 43.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	8268	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.54% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, CA, PGE, EDO, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.58	0/3954	0.67	4/5315 (0.1%)
2	B	1.16	0/74	1.87	2/113 (1.8%)
All	All	0.60	0/4028	0.72	6/5428 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	4	A	C8-N9-C4	8.09	109.04	105.80
1	A	189	ASP	CB-CG-OD2	-7.03	111.98	118.30
1	A	93	ARG	NE-CZ-NH2	-6.60	117.00	120.30
1	A	211	ARG	NE-CZ-NH2	-6.54	117.03	120.30
2	B	3	A	C2-N3-C4	-6.40	107.40	110.60
1	A	159	ARG	NE-CZ-NH1	-5.21	117.70	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3833	3857	3834	9	0
2	B	97	45	45	0	0
3	A	30	42	42	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	7	10	10	0	0
5	A	4	6	6	0	0
6	A	2	0	0	0	0
7	A	315	0	0	2	0
7	B	20	0	0	0	0
All	All	4308	3960	3937	9	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (9) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:382:PHE:CZ	1:A:386:ARG:HD2	2.32	0.65
1:A:357:ARG:HD2	7:A:666:HOH:O	2.04	0.57
1:A:466:LEU:O	1:A:468:ALA:N	2.43	0.51
1:A:300:ILE:O	1:A:304:GLU:HG3	2.13	0.48
1:A:395:ASP:OD2	1:A:428:ARG:NH2	2.46	0.47
1:A:80:ASN:O	1:A:84:GLU:HG2	2.15	0.47
1:A:462:ARG:O	1:A:466:LEU:HD23	2.13	0.47
1:A:317:LYS:O	1:A:321[A]:MET:HG2	2.20	0.42
1:A:300:ILE:HD12	7:A:876:HOH:O	2.20	0.42

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	474/479 (99%)	469 (99%)	4 (1%)	1 (0%)	47 28

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	467	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	416/418 (100%)	415 (100%)	1 (0%)	93	89

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	339	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	2/4 (50%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PEG	A	502	-	6,6,6	0.91	0	5,5,5	0.60	0
3	PGE	A	501	-	9,9,9	0.39	0	8,8,8	0.61	0
5	EDO	A	503	-	3,3,3	0.48	0	2,2,2	0.19	0
3	PGE	A	504[B]	-	9,9,9	0.30	0	8,8,8	0.30	0
3	PGE	A	504[A]	-	9,9,9	0.27	0	8,8,8	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	502	-	-	2/4/4/4	-
3	PGE	A	501	-	-	5/7/7/7	-
5	EDO	A	503	-	-	0/1/1/1	-
3	PGE	A	504[B]	-	-	4/7/7/7	-
3	PGE	A	504[A]	-	-	3/7/7/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	502	PEG	O2-C3-C4-O4
3	A	504[B]	PGE	O3-C5-C6-O4
3	A	504[A]	PGE	O2-C3-C4-O3
3	A	504[B]	PGE	O2-C3-C4-O3
3	A	504[A]	PGE	C4-C3-O2-C2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	A	501	PGE	C6-C5-O3-C4
3	A	501	PGE	C3-C4-O3-C5
3	A	501	PGE	C4-C3-O2-C2
3	A	504[B]	PGE	C1-C2-O2-C3
3	A	504[A]	PGE	C3-C4-O3-C5
3	A	504[B]	PGE	C3-C4-O3-C5
3	A	501	PGE	O2-C3-C4-O3
4	A	502	PEG	O1-C1-C2-O2
3	A	501	PGE	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	461/479 (96%)	-0.00	27 (5%) 22 21	13, 24, 63, 136	0
2	B	3/4 (75%)	-0.51	0 100 100	16, 16, 19, 25	0
All	All	464/483 (96%)	-0.01	27 (5%) 23 22	13, 24, 63, 136	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	435	LEU	7.1
1	A	84	GLU	5.9
1	A	437	LEU	5.4
1	A	432	ARG	4.9
1	A	467	ALA	4.9
1	A	85	GLU	4.7
1	A	88	ASN	4.4
1	A	87	ASP	3.9
1	A	8	HIS	3.8
1	A	464	GLU	3.6
1	A	466	LEU	3.6
1	A	83	GLN	3.5
1	A	462	ARG	3.5
1	A	468	ALA	3.5
1	A	313	GLN	3.4
1	A	433	LYS	3.3
1	A	430	LEU	3.3
1	A	90	ALA	2.7
1	A	465	ARG	2.7
1	A	86	HIS	2.6
1	A	434	ALA	2.6
1	A	426	VAL	2.5
1	A	436	ASP	2.5
1	A	431	ARG	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	89	GLN	2.3
1	A	454	ASN	2.1
1	A	440	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	PGE	A	501	10/10	0.68	0.30	56,75,97,99	0
5	EDO	A	503	4/4	0.72	0.26	67,81,87,87	0
3	PGE	A	504[B]	10/10	0.73	0.20	24,53,64,64	24
3	PGE	A	504[A]	10/10	0.73	0.20	41,53,63,63	24
4	PEG	A	502	7/7	0.82	0.17	57,68,78,78	0
6	CA	A	506	1/1	0.95	0.07	32,32,32,32	1
6	CA	A	505	1/1	0.97	0.08	20,20,20,20	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.