



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 10, 2020 – 10:59 AM BST

PDB ID : 3UER
Title : Crystal structure of amylosucrase from *Deinococcus geothermalis* in complex with turanose
Authors : Guerin, F.; Pizzut-Serin, S.; Potocki-Veronese, G.; Guillet, V.; Mourey, L.; Remaud-Simeon, M.; Andre, I.; Tranier, S.
Deposited on : 2011-10-31
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	655	<div> <div>2%</div> <div> <div></div> <div>86%</div> <div>12%</div> <div>..</div> </div> </div>
2	B	2	<div> <div>100%</div> </div>
3	C	2	<div> <div>50%</div> <div>50%</div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5621 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Amylosucrase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	651	5181	3295	936	937	13	0	10	0

There are 5 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP Q1J0W0
A	-3	PRO	-	expression tag	UNP Q1J0W0
A	-2	LEU	-	expression tag	UNP Q1J0W0
A	-1	GLY	-	expression tag	UNP Q1J0W0
A	0	SER	-	expression tag	UNP Q1J0W0

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-3)-alpha-D-fructofuranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	B	2	Total	C	O	0	2	0
			23	12	11			

- Molecule 3 is an oligosaccharide called alpha-D-glucopyranose-(1-3)-beta-D-fructofuranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	C	2	Total	C	O	0	2	0
			23	12	11			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	394	Total 394	O 394	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Amylosucrase



- Molecule 2: alpha-D-glucopyranose-(1-3)-alpha-D-fructofuranose



- Molecule 3: alpha-D-glucopyranose-(1-3)-beta-D-fructofuranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	104.73Å 110.42Å 115.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	13.00 – 2.10 12.99 – 2.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (13.00-2.10) 99.9 (12.99-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.81 (at 2.09Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.146 , 0.207 0.148 , 0.205	Depositor DCC
R_{free} test set	1971 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	21.2	Xtriage
Anisotropy	0.040	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 48.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5621	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: Z9N, GLC, FRU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.10	6/5320 (0.1%)	0.97	19/7240 (0.3%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	405	GLU	CG-CD	7.16	1.62	1.51
1	A	433	PHE	CE1-CZ	5.64	1.48	1.37
1	A	122	TYR	CE1-CZ	5.34	1.45	1.38
1	A	405	GLU	CB-CG	5.32	1.62	1.52
1	A	352	TYR	CD1-CE1	5.12	1.47	1.39
1	A	187	TRP	CG-CD1	5.04	1.43	1.36

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	478	ARG	NE-CZ-NH1	-10.33	115.14	120.30
1	A	308	ARG	NE-CZ-NH1	7.90	124.25	120.30
1	A	341	ARG	NE-CZ-NH1	7.55	124.08	120.30
1	A	308	ARG	NE-CZ-NH2	-6.54	117.03	120.30
1	A	370	ARG	NE-CZ-NH1	-6.25	117.17	120.30
1	A	341	ARG	NE-CZ-NH2	-6.23	117.19	120.30
1	A	478	ARG	NE-CZ-NH2	5.94	123.27	120.30
1	A	558	ARG	NE-CZ-NH1	5.79	123.19	120.30
1	A	43	ARG	NE-CZ-NH1	5.58	123.09	120.30
1	A	534	ARG	NE-CZ-NH1	-5.52	117.54	120.30
1	A	299	ASN	CB-CA-C	-5.41	99.58	110.40
1	A	558	ARG	NE-CZ-NH2	-5.32	117.64	120.30
1	A	199	ARG	NE-CZ-NH1	5.24	122.92	120.30
1	A	151	ASP	CB-CG-OD2	5.20	122.98	118.30
1	A	199	ARG	NE-CZ-NH2	-5.15	117.72	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	418	ARG	NE-CZ-NH1	5.14	122.87	120.30
1	A	370	ARG	NE-CZ-NH2	5.08	122.84	120.30
1	A	557	ARG	NE-CZ-NH1	5.02	122.81	120.30
1	A	576	ASP	CB-CG-OD1	5.01	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5181	0	5009	60	0
2	B	23	0	6	3	0
3	C	23	0	12	4	0
4	A	394	0	0	7	0
All	All	5621	0	5027	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:585[A]:ASP:OD1	4:A:916:HOH:O	1.62	1.15
1:A:299:ASN:HD21	1:A:329:VAL:HG11	1.20	1.03
1:A:326:GLU:OE2	3:C:1[B]:FRU:H12	1.62	1.00
1:A:190:LYS:HD3	4:A:1149:HOH:O	1.61	0.98
1:A:335:ILE:HG13	1:A:380:PRO:HG2	1.55	0.88
1:A:34:TYR:OH	1:A:315[B]:ARG:NH1	2.07	0.86
3:C:1[B]:FRU:H62	3:C:1[B]:FRU:O2	1.76	0.83
1:A:610:ARG:HD3	1:A:615:ASP:OD1	1.83	0.79
1:A:240:GLU:O	4:A:1183:HOH:O	2.02	0.78
1:A:435[A]:ARG:NH1	4:A:1138:HOH:O	2.16	0.78
1:A:14:VAL:HG11	1:A:28[B]:LEU:HD22	1.65	0.78
1:A:591:MET:HE2	1:A:642:TRP:HB3	1.69	0.74

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:413:ASN:HD22	1:A:416:ALA:H	1.40	0.70
1:A:452:GLY:H	1:A:523:HIS:HD2	1.39	0.69
1:A:299:ASN:HD21	1:A:329:VAL:CG1	2.02	0.69
1:A:298:GLN:O	1:A:299:ASN:HB2	1.93	0.67
1:A:292:ARG:HH11	1:A:300:GLN:HE21	1.42	0.67
1:A:326:GLU:OE1	1:A:326:GLU:O	2.13	0.66
1:A:473:GLU:OE1	1:A:545:ARG:NH2	2.31	0.64
2:B:1[A]:Z9N:C1	2:B:2[A]:GLC:C1	2.80	0.60
1:A:452:GLY:H	1:A:523:HIS:CD2	2.20	0.59
1:A:299:ASN:ND2	1:A:329:VAL:HG11	2.04	0.59
1:A:610:ARG:CD	1:A:615:ASP:OD1	2.51	0.57
1:A:15[A]:ARG:NH1	1:A:25:GLU:OE1	2.38	0.56
1:A:24:ALA:O	1:A:28[B]:LEU:HD23	2.06	0.56
1:A:233:ASP:O	1:A:241:GLY:HA3	2.07	0.55
2:B:1[A]:Z9N:C6	2:B:1[A]:Z9N:O1	2.55	0.54
1:A:143[B]:GLN:NE2	4:A:960:HOH:O	2.31	0.53
1:A:298:GLN:O	1:A:299:ASN:CB	2.55	0.52
1:A:53:LEU:N	1:A:54:PRO:HD3	2.25	0.52
1:A:43:ARG:HD3	4:A:1145:HOH:O	2.11	0.51
1:A:311:ARG:O	1:A:315[B]:ARG:HG3	2.11	0.51
1:A:28[A]:LEU:O	1:A:32:GLU:HG3	2.12	0.49
1:A:596:ASN:O	1:A:638:TYR:HA	2.11	0.49
1:A:519:ASN:C	1:A:519:ASN:HD22	2.16	0.49
1:A:14:VAL:CG1	1:A:28[B]:LEU:HD22	2.40	0.48
1:A:335:ILE:CG1	1:A:380:PRO:HG2	2.37	0.48
1:A:53:LEU:N	1:A:54:PRO:CD	2.78	0.47
1:A:636[C]:GLU:H	1:A:636[C]:GLU:CD	2.16	0.47
2:B:1[A]:Z9N:C1	2:B:1[A]:Z9N:C6	2.93	0.47
1:A:208:ARG:HD3	1:A:243:TRP:CZ2	2.51	0.46
1:A:233:ASP:HB3	1:A:236:ILE:HG13	1.97	0.46
1:A:10:LEU:CD2	1:A:54:PRO:HB3	2.46	0.46
1:A:406:ASP:OD1	1:A:409:ARG:NH2	2.49	0.46
1:A:248:PHE:HE1	3:C:1[B]:FRU:O6	1.99	0.45
1:A:350:MET:HA	1:A:387:THR:O	2.17	0.45
1:A:591:MET:CE	1:A:642:TRP:HB3	2.44	0.44
1:A:10:LEU:HD21	1:A:54:PRO:HB3	1.99	0.44
1:A:433:PHE:O	1:A:459:GLY:HA2	2.17	0.44
1:A:395:HIS:O	1:A:523:HIS:HE1	2.01	0.44
1:A:240:GLU:O	1:A:241:GLY:O	2.35	0.43
1:A:37:ASP:OD2	1:A:345:GLY:HA3	2.18	0.43
1:A:547:ASN:HD21	1:A:551:ARG:HE	1.67	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:43:ARG:HA	1:A:50:VAL:HG21	2.01	0.42
1:A:599:GLU:HG3	1:A:638:TYR:CE2	2.54	0.42
1:A:468:ASP:HA	1:A:469:PRO:HD3	1.93	0.42
1:A:106:LYS:HE2	1:A:157:ASP:OD1	2.20	0.41
1:A:326:GLU:CD	3:C:1[B]:FRU:H12	2.35	0.41
1:A:143[B]:GLN:HG2	1:A:143[B]:GLN:O	2.21	0.41
1:A:271:LEU:HD21	1:A:314:ALA:HA	2.02	0.41
1:A:409:ARG:NH2	4:A:1132:HOH:O	2.53	0.41
1:A:441:TYR:HD1	1:A:448:ARG:HG2	1.85	0.41
1:A:439:PHE:HB2	1:A:519:ASN:HD21	1.86	0.41
1:A:236:ILE:O	1:A:241:GLY:HA2	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	659/655 (101%)	641 (97%)	16 (2%)	2 (0%)	41 41

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	241	GLY
1	A	624	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	517/532 (97%)	504 (98%)	13 (2%)	47 52

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	LEU
1	A	100	ARG
1	A	122	TYR
1	A	128	LEU
1	A	143[A]	GLN
1	A	143[B]	GLN
1	A	299	ASN
1	A	326	GLU
1	A	341	ARG
1	A	372	PHE
1	A	519	ASN
1	A	538	GLU
1	A	558	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	63	HIS
1	A	299	ASN
1	A	300	GLN
1	A	413	ASN
1	A	519	ASN
1	A	523	HIS
1	A	547	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	Z9N	B	1[A]	2	11,12,12	4.03	7 (63%)	10,18,18	3.30	6 (60%)
2	GLC	B	2[A]	2	11,11,12	2.45	3 (27%)	15,15,17	2.34	3 (20%)
3	FRU	C	1[B]	3	11,12,12	4.11	5 (45%)	10,18,18	3.49	5 (50%)
3	GLC	C	2[B]	3	11,11,12	2.47	3 (27%)	15,15,17	2.32	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	Z9N	B	1[A]	2	-	3/5/24/24	0/1/1/1
2	GLC	B	2[A]	2	-	0/2/19/22	0/1/1/1
3	FRU	C	1[B]	3	-	3/5/24/24	0/1/1/1
3	GLC	C	2[B]	3	-	0/2/19/22	0/1/1/1

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1[B]	FRU	O3-C3	-10.29	1.22	1.42
2	B	1[A]	Z9N	O3-C3	-9.46	1.23	1.42
3	C	2[B]	GLC	O5-C1	-6.46	1.33	1.43
2	B	2[A]	GLC	O5-C1	-6.34	1.33	1.43
2	B	1[A]	Z9N	O5-C2	-6.15	1.33	1.43
3	C	1[B]	FRU	O5-C5	-5.58	1.31	1.43
3	C	1[B]	FRU	O5-C2	-5.46	1.34	1.43
2	B	1[A]	Z9N	O5-C5	-4.22	1.34	1.43
2	B	1[A]	Z9N	O1-C1	-3.45	1.30	1.42
3	C	2[B]	GLC	O3-C3	-2.95	1.36	1.43
3	C	1[B]	FRU	C6-C5	-2.95	1.42	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2[A]	GLC	O3-C3	-2.94	1.36	1.43
2	B	2[A]	GLC	O2-C2	-2.64	1.37	1.43
3	C	2[B]	GLC	O2-C2	-2.59	1.37	1.43
2	B	1[A]	Z9N	C1-C2	-2.53	1.48	1.52
3	C	1[B]	FRU	O6-C6	-2.45	1.32	1.42
2	B	1[A]	Z9N	C6-C5	-2.43	1.43	1.51
2	B	1[A]	Z9N	O6-C6	-2.29	1.32	1.42

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1[A]	Z9N	C6-C5-C4	-7.60	96.78	115.09
3	C	1[B]	FRU	C6-C5-C4	-7.23	97.66	115.09
2	B	2[A]	GLC	O5-C1-C2	-5.85	101.73	110.77
3	C	2[B]	GLC	C1-O5-C5	5.60	119.78	112.19
3	C	2[B]	GLC	O5-C1-C2	-5.58	102.16	110.77
2	B	2[A]	GLC	C1-O5-C5	5.52	119.67	112.19
3	C	1[B]	FRU	O3-C3-C4	-4.36	98.25	113.32
3	C	1[B]	FRU	O2-C2-O5	4.35	117.91	109.50
2	B	1[A]	Z9N	O3-C3-C4	-4.16	98.96	113.32
2	B	1[A]	Z9N	O5-C5-C6	-4.07	97.52	108.85
3	C	1[B]	FRU	O5-C5-C6	-4.01	97.68	108.85
3	C	1[B]	FRU	O6-C6-C5	-3.25	100.13	111.29
2	B	1[A]	Z9N	O6-C6-C5	-2.43	102.96	111.29
2	B	1[A]	Z9N	O2-C2-O5	-2.28	105.09	109.50
3	C	2[B]	GLC	O5-C5-C6	-2.21	103.75	107.20
2	B	1[A]	Z9N	O5-C5-C4	2.10	110.72	105.49
2	B	2[A]	GLC	O5-C5-C6	-2.01	104.05	107.20

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	1[B]	FRU	O1-C1-C2-C3
3	C	1[B]	FRU	O1-C1-C2-O2
2	B	1[A]	Z9N	O1-C1-C2-O2
2	B	1[A]	Z9N	O1-C1-C2-O5
2	B	1[A]	Z9N	O1-C1-C2-C3
3	C	1[B]	FRU	O1-C1-C2-O5


There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2[A]	GLC	1	0
3	C	1[B]	FRU	4	0
2	B	1[A]	Z9N	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	651/655 (99%)	-0.49	13 (1%) 65 69	9, 18, 38, 53	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	343	HIS	5.2
1	A	239	GLY	4.6
1	A	646	GLY	4.1
1	A	240	GLU	3.8
1	A	625	ALA	3.4
1	A	624	SER	3.1
1	A	444	VAL	2.4
1	A	629	ASP	2.4
1	A	238	GLU	2.3
1	A	4	ASP	2.2
1	A	344	HIS	2.1
1	A	645	ALA	2.1
1	A	615	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	FRU	C	1[B]	12/12	0.85	0.21	24,29,31,33	12
2	Z9N	B	1[A]	12/12	0.87	0.21	31,36,38,39	12
2	GLC	B	2[A]	11/12	0.95	0.10	17,22,25,26	11
3	GLC	C	2[B]	11/12	0.95	0.10	10,16,19,19	11

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.