



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 11:35 am BST

PDB ID : 3UG1  
Title : Crystal structure of the mutated EGFR kinase domain (G719S/T790M) in the apo form  
Authors : Parker, L.J.; Handa, N.; Yoshikawa, S.; Kukimoto-Niino, M.; Shirouzu, M.; Yokoyama, S.  
Deposited on : 2011-11-02  
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1235 (2.78-2.74)
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	<div> <div>3%</div> <div>51%</div> <div>32%</div> <div>5%</div> <div>11%</div> </div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2329 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Epidermal growth factor receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	296	2308	1486	386	420	16	0	1	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	689	GLY	-	EXPRESSION TAG	UNP P00533
A	690	ALA	-	EXPRESSION TAG	UNP P00533
A	691	MET	-	EXPRESSION TAG	UNP P00533
A	692	GLY	-	EXPRESSION TAG	UNP P00533
A	693	ILE	-	EXPRESSION TAG	UNP P00533
A	694	ARG	-	EXPRESSION TAG	UNP P00533
A	719	SER	GLY	ENGINEERED MUTATION	UNP P00533
A	790	MET	THR	ENGINEERED MUTATION	UNP P00533

- Molecule 2 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	9	Total	O	0	0
			9	9		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Epidermal growth factor receptor



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	141.33Å 141.33Å 141.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.75 49.97 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.8 (50.00-2.75) 99.9 (49.97-2.75)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.71 (at 2.77Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.196 , 0.243 0.194 , 0.241	Depositor DCC
$R_{free}$ test set	617 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	58.0	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 45.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.032 for -l,-k,-h	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2329	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.23% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.84	0/2360	0.88	1/3205 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	837	ASP	CB-CG-OD1	6.16	123.84	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2308	0	2276	90	0
2	A	12	0	12	3	0
3	A	9	0	0	0	0
All	All	2329	0	2288	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:725:THR:HG22	1:A:727:TYR:HE1	1.08	1.12
1:A:725:THR:HG22	1:A:727:TYR:CE1	1.97	0.98
1:A:783:THR:HG22	1:A:785:THR:O	1.69	0.92
1:A:878:ILE:HG23	1:A:886:ILE:HD12	1.57	0.87
1:A:918:ILE:HG23	1:A:919:PRO:HD2	1.59	0.83
1:A:783:THR:CG2	1:A:785:THR:O	2.32	0.77
1:A:765:VAL:HG13	1:A:833:LEU:HD11	1.67	0.77
1:A:697:GLU:HA	1:A:697:GLU:OE1	1.87	0.75
1:A:878:ILE:HG23	1:A:886:ILE:CD1	2.18	0.74
1:A:753:PRO:HA	1:A:756:ASN:HD22	1.52	0.73
1:A:834:VAL:HG12	1:A:836:ARG:HG3	1.72	0.71
1:A:892:THR:H	1:A:895:SER:HB2	1.55	0.71
1:A:919:PRO:HG2	1:A:922:GLU:CB	2.21	0.70
1:A:765:VAL:CG1	1:A:833:LEU:HD11	2.23	0.69
1:A:880:TRP:HZ3	2:A:10:MES:H82	1.57	0.69
1:A:926:ILE:CG2	1:A:931:GLU:HB3	2.26	0.66
1:A:771:ASN:HD22	1:A:773:HIS:H	1.43	0.65
1:A:880:TRP:CD1	1:A:902:VAL:HG22	2.32	0.64
1:A:771:ASN:ND2	1:A:773:HIS:H	1.97	0.62
1:A:918:ILE:CG2	1:A:919:PRO:HD2	2.28	0.62
1:A:877:PRO:O	1:A:881:MET:HG3	2.02	0.60
1:A:793:MET:HG3	1:A:844:LEU:HD23	1.83	0.60
1:A:820:GLN:HE22	1:A:851:VAL:H	1.50	0.59
1:A:820:GLN:NE2	1:A:851:VAL:H	2.01	0.58
1:A:812:GLN:NE2	1:A:975:PRO:CG	2.66	0.58
1:A:717:VAL:HG12	1:A:719:SER:O	2.03	0.57
1:A:887:LEU:HB3	1:A:888:HIS:CD2	2.38	0.57
1:A:697:GLU:CA	1:A:697:GLU:OE1	2.52	0.57
1:A:780:ILE:HG22	1:A:788:LEU:HD12	1.86	0.57
1:A:811:SER:HB2	1:A:975:PRO:HB3	1.88	0.56
1:A:772:PRO:O	1:A:852:LYS:HE3	2.07	0.54
1:A:726:VAL:C	1:A:727:TYR:HD1	2.10	0.54
1:A:708:LYS:HE3	1:A:710:THR:HG22	1.89	0.54
1:A:836:ARG:HG2	1:A:891:TYR:CD1	2.44	0.53
1:A:926:ILE:HG23	1:A:931:GLU:HB3	1.91	0.53
1:A:812:GLN:NE2	1:A:975:PRO:HG3	2.24	0.53
1:A:880:TRP:HZ3	2:A:10:MES:C8	2.22	0.53
1:A:965:ILE:O	1:A:969:SER:HB2	2.09	0.52
1:A:833:LEU:HD21	1:A:859:ALA:HB1	1.90	0.52
1:A:1018:ILE:HD12	1:A:1018:ILE:O	2.10	0.52
1:A:717:VAL:CG1	1:A:719:SER:O	2.58	0.51
1:A:714:LYS:HD3	1:A:727:TYR:CD2	2.46	0.51

Continued on next page...



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:811:SER:HB2	1:A:975:PRO:CB	2.41	0.51
1:A:812:GLN:NE2	1:A:975:PRO:HG2	2.26	0.51
1:A:883:LEU:HA	1:A:886:ILE:HG22	1.93	0.51
1:A:700:ASN:HD21	1:A:768:SER:HB3	1.76	0.50
1:A:805:HIS:O	1:A:806:LYS:C	2.50	0.50
1:A:882:ALA:O	1:A:886:ILE:HG22	2.12	0.49
1:A:878:ILE:N	2:A:10:MES:O1S	2.45	0.49
1:A:820:GLN:HE21	1:A:851:VAL:HG22	1.78	0.49
1:A:791:GLN:NE2	1:A:1012:ASP:OD2	2.45	0.48
1:A:1017:LEU:O	1:A:1018:ILE:HB	2.13	0.48
1:A:806:LYS:HG3	1:A:910:PHE:HB3	1.96	0.48
1:A:825:MET:HE2	1:A:853:ILE:HG21	1.95	0.48
1:A:713:LYS:HE3	1:A:713:LYS:HB2	1.44	0.48
1:A:922:GLU:O	1:A:923:ILE:C	2.52	0.47
1:A:825:MET:HE2	1:A:853:ILE:HD13	1.95	0.47
1:A:885:SER:O	1:A:889:ARG:HA	2.14	0.47
1:A:829:GLU:HG2	1:A:893:HIS:CG	2.50	0.47
1:A:775:CYS:HB2	1:A:853:ILE:O	2.15	0.46
1:A:954:ASP:O	1:A:957:SER:HB2	2.16	0.46
1:A:727:TYR:N	1:A:727:TYR:CD1	2.84	0.46
1:A:829:GLU:HG2	1:A:893:HIS:CD2	2.51	0.46
1:A:781:CYS:HB3	1:A:787:GLN:HB2	1.98	0.45
1:A:725:THR:CG2	1:A:727:TYR:HE1	2.01	0.45
1:A:736:GLU:O	1:A:738:VAL:HG13	2.16	0.45
1:A:766:MET:HE2	1:A:790:MET:HG2	1.98	0.45
1:A:1018:ILE:CD1	1:A:1018:ILE:O	2.64	0.45
1:A:764:TYR:O	1:A:768:SER:OG	2.35	0.44
1:A:922:GLU:O	1:A:925:SER:N	2.51	0.44
1:A:766:MET:HE1	1:A:790:MET:SD	2.58	0.44
1:A:1003:ASP:HB2	1:A:1004:GLU:H	1.53	0.43
1:A:935:GLN:HA	1:A:944:TYR:CE1	2.53	0.43
1:A:898:TRP:HE3	1:A:958:ARG:NH2	2.17	0.43
1:A:892:THR:H	1:A:895:SER:CB	2.28	0.43
1:A:756:ASN:HA	1:A:759:ILE:HG22	2.01	0.43
1:A:746:GLU:HA	1:A:787:GLN:HG2	2.00	0.43
1:A:825:MET:CE	1:A:853:ILE:HG21	2.48	0.43
1:A:945:MET:O	1:A:949:LYS:HG3	2.18	0.43
1:A:894:GLN:O	1:A:897:VAL:HB	2.19	0.42
1:A:913:LYS:HB3	1:A:913:LYS:HE3	1.72	0.42
1:A:880:TRP:CD1	1:A:902:VAL:CG2	3.01	0.42
1:A:855:ASP:C	1:A:855:ASP:OD1	2.58	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:963:GLU:O	1:A:967:GLU:HG2	2.20	0.42
1:A:936:PRO:HA	1:A:937:PRO:HD2	1.80	0.41
1:A:961:PHE:O	1:A:965:ILE:HG13	2.21	0.41
1:A:714:LYS:HD3	1:A:727:TYR:CE2	2.56	0.41
1:A:882:ALA:HA	1:A:898:TRP:CD2	2.55	0.40
1:A:766:MET:CE	1:A:790:MET:SD	3.09	0.40
1:A:923:ILE:HG22	1:A:927:LEU:HD22	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	289/334 (86%)	265 (92%)	18 (6%)	6 (2%)	<b>7</b> <b>12</b>

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	864	ALA
1	A	940	THR
1	A	1004	GLU
1	A	1005	GLU
1	A	806	LYS
1	A	923	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	245/292 (84%)	209 (85%)	36 (15%)	<b>3</b> <b>4</b>

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	697	GLU
1	A	710	THR
1	A	713	LYS
1	A	718	LEU
1	A	720	SER
1	A	732	ILE
1	A	745	LYS
1	A	766	MET
1	A	768	SER
1	A	783	THR
1	A	784	SER
1	A	790	MET
1	A	799	LEU
1	A	802	VAL
1	A	806	LYS
1	A	811	SER
1	A	814	LEU
1	A	844	LEU
1	A	849	GLN
1	A	858	LEU
1	A	861	LEU
1	A	862	LEU
1	A	876	VAL
1	A	902	VAL
1	A	921	SER
1	A	927	LEU
1	A	938	ILE
1	A	941	ILE
1	A	952	MET
1	A	953	ILE
1	A	962	ARG
1	A	964	LEU
1	A	969	SER
1	A	1003	ASP
1	A	1008	ASP
1	A	1018	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	756	ASN
1	A	771	ASN
1	A	787	GLN
1	A	812	GLN
1	A	820	GLN
1	A	888	HIS
1	A	976	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	MES	A	10	-	12,12,12	2.47	1 (8%)	14,16,16	2.90	6 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MES	A	10	-	-	2/6/14/14	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	10	MES	C8-S	-8.35	1.65	1.77

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	10	MES	C5-N4-C3	7.22	125.07	108.83
2	A	10	MES	O1S-S-C8	5.81	113.92	106.92
2	A	10	MES	C7-N4-C5	2.71	118.17	111.23
2	A	10	MES	O2S-S-C8	2.44	109.85	106.92
2	A	10	MES	O2S-S-O1S	-2.18	106.39	113.95
2	A	10	MES	C7-N4-C3	2.15	116.74	111.23

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	10	MES	N4-C7-C8-S
2	A	10	MES	C8-C7-N4-C3

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	10	MES	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	296/334 (88%)	0.07	9 (3%) 50 59	31, 49, 78, 91	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	874	GLY	3.8
1	A	919	PRO	3.4
1	A	718	LEU	2.8
1	A	725	THR	2.6
1	A	727	TYR	2.5
1	A	720	SER	2.4
1	A	918	ILE	2.3
1	A	715	ILE	2.1
1	A	714	LYS	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MES	A	10	12/12	0.97	0.21	51,52,54,55	12

## 6.5 Other polymers ⓘ

There are no such residues in this entry.