



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 8, 2020 – 10:08 PM BST

PDB ID : 6UG9  
Title : Complex of ch28/11 Fab and SSEA-4 (hexagonal form)  
Authors : Soliman, C.; Ramsland, P.A.  
Deposited on : 2019-09-26  
Resolution : 2.74 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1271 (2.76-2.72)
Clashscore	141614	1322 (2.76-2.72)
Ramachandran outliers	138981	1297 (2.76-2.72)
Sidechain outliers	138945	1298 (2.76-2.72)
RSRZ outliers	127900	1243 (2.76-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	213	<div> <div>92%</div> <div>8%</div> <div>.</div> </div>
1	K	213	<div> <div>91%</div> <div>8%</div> <div>..</div> </div>
1	L	213	<div> <div>91%</div> <div>7%</div> <div>..</div> </div>
2	A	218	<div> <div>85%</div> <div>11%</div> <div>.</div> </div>
2	H	218	<div> <div>88%</div> <div>8%</div> <div>..</div> </div>
2	J	218	<div> <div>%</div> <div>90%</div> <div>7%</div> <div>..</div> </div>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	C	6	 17% 83%
3	D	6	 17% 83%
3	E	6	 100%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9965 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ch28/11 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	211	Total	C	N	O	S	0	0	0
			1609	1007	267	327	8			
1	B	211	Total	C	N	O	S	0	0	0
			1609	1007	267	327	8			
1	K	211	Total	C	N	O	S	0	0	0
			1609	1007	267	327	8			

- Molecule 2 is a protein called ch28/11 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	212	Total	C	N	O	S	0	0	0
			1582	1005	264	308	5			
2	A	212	Total	C	N	O	S	0	0	0
			1582	1005	264	308	5			
2	J	216	Total	C	N	O	S	0	0	0
			1605	1017	268	315	5			

- Molecule 3 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-3)-alpha-D-galactopyranose-(1-4)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	6	Total	C	N	O	0	0	0
			79	43	2	34			
3	D	6	Total	C	N	O	0	0	0
			79	43	2	34			
3	E	6	Total	C	N	O	0	0	0
			79	43	2	34			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	L	23	Total 23	O 23	0	0
4	H	34	Total 34	O 34	0	0
4	B	19	Total 19	O 19	0	0
4	A	18	Total 18	O 18	0	0
4	K	20	Total 20	O 20	0	0
4	J	18	Total 18	O 18	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ch28/11 Fab light chain

Chain L:  91% 7% ..



- Molecule 1: ch28/11 Fab light chain

Chain B:  92% 8% .





- Molecule 1: ch28/11 Fab light chain

Chain K:  91% 8% ..





- Molecule 2: ch28/11 Fab heavy chain

Chain H:  88% 8% ..




- Molecule 2: ch28/11 Fab heavy chain

Chain A:  85% 11% .



- Molecule 2: ch28/11 Fab heavy chain

Chain J:  90% 7% ..



- Molecule 3: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-3)-alpha-D-galactopyranose-(1-4)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranose

Chain C: 17% 83%



- Molecule 3: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-3)-alpha-D-galactopyranose-(1-4)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranose

Chain D: 17% 83%



- Molecule 3: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-3)-alpha-D-galactopyranose-(1-4)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranose

Chain E: 100%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	169.95Å 169.95Å 95.44Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.74 48.06 – 2.74	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-2.74) 100.0 (48.06-2.74)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.64 (at 2.73Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, $R_{free}$	0.170 , 0.247 0.176 , 0.248	Depositor DCC
$R_{free}$ test set	1992 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	52.7	Xtriage
Anisotropy	0.270	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 32.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.029 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9965	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.37% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GAL, GLA, SIA, BGC, NGA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	B	0.60	0/1645	0.76	0/2231
1	K	0.60	0/1645	0.77	0/2231
1	L	0.63	0/1645	0.78	0/2231
2	A	0.65	0/1622	0.79	0/2214
2	H	0.65	0/1622	0.83	2/2214 (0.1%)
2	J	0.62	0/1645	0.80	2/2245 (0.1%)
All	All	0.63	0/9824	0.79	4/13366 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	J	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	66	ARG	NE-CZ-NH1	6.59	123.60	120.30
2	J	80	LEU	CA-CB-CG	5.88	128.81	115.30
2	J	182	LEU	CA-CB-CG	5.63	128.25	115.30
2	H	66	ARG	NE-CZ-NH2	-5.28	117.66	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	J	64	MET	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1609	0	1558	6	0
1	K	1609	0	1558	4	0
1	L	1609	0	1558	6	0
2	A	1582	0	1565	8	0
2	H	1582	0	1565	5	0
2	J	1605	0	1585	11	0
3	C	79	0	67	0	0
3	D	79	0	67	1	0
3	E	79	0	67	0	0
4	A	18	0	0	1	0
4	B	19	0	0	0	0
4	H	34	0	0	2	0
4	J	18	0	0	0	0
4	K	20	0	0	1	0
4	L	23	0	0	1	0
All	All	9965	0	9590	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 40 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:104:GLU:OE1	1:B:172:TYR:OH	2.11	0.68
1:B:154:GLN:HE21	1:B:157:ASN:HD21	1.44	0.64
1:L:90:ALA:HA	1:L:95:LEU:HD22	1.82	0.62
2:J:90:THR:HG23	2:J:114:THR:HA	1.82	0.62
1:B:105:LEU:HD23	1:B:170:SER:OG	2.05	0.57

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	209/213 (98%)	197 (94%)	11 (5%)	1 (0%)	29	48
1	K	209/213 (98%)	199 (95%)	10 (5%)	0	100	100
1	L	209/213 (98%)	199 (95%)	10 (5%)	0	100	100
2	A	208/218 (95%)	201 (97%)	7 (3%)	0	100	100
2	H	208/218 (95%)	199 (96%)	6 (3%)	3 (1%)	11	20
2	J	212/218 (97%)	200 (94%)	10 (5%)	2 (1%)	17	32
All	All	1255/1293 (97%)	1195 (95%)	54 (4%)	6 (0%)	29	48

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	65	SER
2	J	65	SER
2	H	64	MET
1	B	210	ARG
2	H	16	GLN

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	183/185 (99%)	178 (97%)	5 (3%)	44	65
1	K	183/185 (99%)	170 (93%)	13 (7%)	14	26
1	L	183/185 (99%)	175 (96%)	8 (4%)	28	47

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	180/185 (97%)	167 (93%)	13 (7%)	14	25
2	H	180/185 (97%)	168 (93%)	12 (7%)	16	29
2	J	183/185 (99%)	174 (95%)	9 (5%)	25	43
All	All	1092/1110 (98%)	1032 (94%)	60 (6%)	21	37

5 of 60 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	96	THR
2	A	197	THR
2	J	119	SER
2	A	165	SER
2	A	205	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	154	GLN
1	K	30	ASN
1	K	137	ASN
1	K	146	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

18 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	BGC	C	1	3	12,12,12	0.54	0	17,17,17	1.27	2 (11%)
3	GAL	C	2	3	11,11,12	0.40	0	15,15,17	0.97	0
3	GLA	C	3	3	11,11,12	0.46	0	15,15,17	1.77	4 (26%)
3	NGA	C	4	3	14,14,15	0.65	0	17,19,21	1.23	2 (11%)
3	GAL	C	5	3	11,11,12	0.74	0	15,15,17	0.89	1 (6%)
3	SIA	C	6	3	17,20,21	0.89	1 (5%)	21,28,31	1.16	2 (9%)
3	BGC	D	1	3	12,12,12	0.71	0	17,17,17	1.48	4 (23%)
3	GAL	D	2	3	11,11,12	0.54	0	15,15,17	0.81	0
3	GLA	D	3	3	11,11,12	0.35	0	15,15,17	1.44	2 (13%)
3	NGA	D	4	3	14,14,15	0.52	0	17,19,21	0.96	0
3	GAL	D	5	3	11,11,12	0.71	0	15,15,17	1.58	3 (20%)
3	SIA	D	6	3	17,20,21	0.74	0	21,28,31	1.31	3 (14%)
3	BGC	E	1	3	12,12,12	0.64	0	17,17,17	1.07	1 (5%)
3	GAL	E	2	3	11,11,12	0.26	0	15,15,17	1.39	2 (13%)
3	GLA	E	3	3	11,11,12	0.53	0	15,15,17	1.13	2 (13%)
3	NGA	E	4	3	14,14,15	0.45	0	17,19,21	1.19	2 (11%)
3	GAL	E	5	3	11,11,12	0.46	0	15,15,17	1.22	1 (6%)
3	SIA	E	6	3	17,20,21	0.59	1 (5%)	21,28,31	1.57	5 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BGC	C	1	3	-	1/2/22/22	0/1/1/1
3	GAL	C	2	3	-	0/2/19/22	0/1/1/1
3	GLA	C	3	3	-	1/2/19/22	0/1/1/1
3	NGA	C	4	3	-	0/6/23/26	0/1/1/1
3	GAL	C	5	3	-	0/2/19/22	0/1/1/1
3	SIA	C	6	3	-	0/14/34/38	0/1/1/1
3	BGC	D	1	3	-	0/2/22/22	0/1/1/1
3	GAL	D	2	3	-	2/2/19/22	0/1/1/1
3	GLA	D	3	3	-	2/2/19/22	0/1/1/1
3	NGA	D	4	3	-	0/6/23/26	0/1/1/1
3	GAL	D	5	3	-	0/2/19/22	0/1/1/1

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SIA	D	6	3	-	0/14/34/38	0/1/1/1
3	BGC	E	1	3	-	2/2/22/22	0/1/1/1
3	GAL	E	2	3	-	2/2/19/22	0/1/1/1
3	GLA	E	3	3	-	2/2/19/22	0/1/1/1
3	NGA	E	4	3	-	1/6/23/26	0/1/1/1
3	GAL	E	5	3	-	0/2/19/22	0/1/1/1
3	SIA	E	6	3	-	1/14/34/38	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	6	SIA	C4-C5	-2.69	1.50	1.53
3	E	6	SIA	C7-C6	2.06	1.55	1.53

The worst 5 of 36 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	6	SIA	C3-C4-C5	4.29	116.64	111.46
3	C	3	GLA	O5-C5-C6	3.70	113.00	107.20
3	C	4	NGA	C1-O5-C5	3.55	117.00	112.19
3	D	5	GAL	C1-C2-C3	3.52	114.00	109.67
3	C	6	SIA	C6-O6-C2	3.48	118.78	111.34

There are no chirality outliers.

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	6	SIA	O6-C6-C7-O7
3	D	3	GLA	O5-C5-C6-O6
3	E	1	BGC	O5-C5-C6-O6
3	E	1	BGC	C4-C5-C6-O6
3	E	2	GAL	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	2	GAL	1	0

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	B	211/213 (99%)	-0.34	0	100	100	39, 59, 81, 88	0
1	K	211/213 (99%)	-0.33	0	100	100	34, 52, 75, 99	0
1	L	211/213 (99%)	-0.39	0	100	100	32, 48, 74, 95	0
2	A	212/218 (97%)	-0.32	0	100	100	36, 49, 70, 84	0
2	H	212/218 (97%)	-0.38	0	100	100	33, 48, 73, 83	0
2	J	216/218 (99%)	-0.16	3 (1%)	75	80	41, 60, 84, 104	0
All	All	1273/1293 (98%)	-0.32	3 (0%)	95	97	32, 52, 78, 104	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	J	135	THR	3.7
2	J	136	SER	2.4
2	J	132	SER	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	BGC	D	1	12/12	0.91	0.16	66,81,84,85	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	BGC	E	1	12/12	0.92	0.18	80,93,99,100	0
3	BGC	C	1	12/12	0.93	0.14	60,70,86,89	0
3	GLA	E	3	11/12	0.95	0.17	62,70,77,78	0
3	GAL	E	2	11/12	0.96	0.18	61,69,73,74	0
3	SIA	E	6	20/21	0.96	0.16	58,72,80,90	0
3	GAL	C	5	11/12	0.96	0.13	43,47,50,52	0
3	GAL	E	5	11/12	0.96	0.15	50,56,61,63	0
3	GAL	D	5	11/12	0.96	0.13	45,50,51,55	0
3	GAL	C	2	11/12	0.97	0.13	44,48,52,55	0
3	NGA	E	4	14/15	0.97	0.16	58,68,72,74	0
3	GLA	D	3	11/12	0.97	0.14	44,46,53,60	0
3	SIA	C	6	20/21	0.97	0.11	52,55,59,61	0
3	NGA	D	4	14/15	0.97	0.15	45,48,64,67	0
3	SIA	D	6	20/21	0.97	0.10	42,55,61,62	0
3	NGA	C	4	14/15	0.98	0.15	36,43,48,49	0
3	GAL	D	2	11/12	0.98	0.12	47,59,67,72	0
3	GLA	C	3	11/12	0.98	0.14	45,48,54,55	0

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.