



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 4, 2022 – 12:31 PM EDT

PDB ID : 7UGH
Title : Crystal Structure of enolase family protein from Naegleria fowleri with bound 2-phosphoglyceric acid
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2022-03-24
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : **FAILED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.27


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain
1	A	477	
1	B	477	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7525 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphopyruvate hydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	436	Total	C	N	O	S	0	12	0
			3407	2137	599	663	8			
1	B	434	Total	C	N	O	S	0	17	0
			3411	2144	592	667	8			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	36	MET	-	initiating methionine	UNP A0A6A5BXC3
A	37	ALA	-	expression tag	UNP A0A6A5BXC3
A	38	HIS	-	expression tag	UNP A0A6A5BXC3
A	39	HIS	-	expression tag	UNP A0A6A5BXC3
A	40	HIS	-	expression tag	UNP A0A6A5BXC3
A	41	HIS	-	expression tag	UNP A0A6A5BXC3
A	42	HIS	-	expression tag	UNP A0A6A5BXC3
A	43	HIS	-	expression tag	UNP A0A6A5BXC3
B	36	MET	-	initiating methionine	UNP A0A6A5BXC3
B	37	ALA	-	expression tag	UNP A0A6A5BXC3
B	38	HIS	-	expression tag	UNP A0A6A5BXC3
B	39	HIS	-	expression tag	UNP A0A6A5BXC3
B	40	HIS	-	expression tag	UNP A0A6A5BXC3
B	41	HIS	-	expression tag	UNP A0A6A5BXC3
B	42	HIS	-	expression tag	UNP A0A6A5BXC3
B	43	HIS	-	expression tag	UNP A0A6A5BXC3

- Molecule 2 is 2-PHOSPHOGLYCERIC ACID (three-letter code: 2PG) (formula: C₃H₇O₇P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
2	A	1	Total	C	O	P	0	0
			11	3	7	1		
2	B	1	Total	C	O	P	0	0
			11	3	7	1		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	2	Total Na 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	2	Total 2	Na 2	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	319	Total 320	O 320	0	1
6	B	321	Total 321	O 321	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS failed to run properly.


- Molecule 1: Phosphopyruvate hydratase

Chain A:  86% 5% 9%



- Molecule 1: Phosphopyruvate hydratase

Chain B:  84% 7% 9%



4 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	96.86Å 96.86Å 216.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.26 – 1.95	Depositor
% Data completeness (in resolution range)	99.0 (47.26-1.95)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.94 (at 1.95Å)	Xtrriage
Refinement program	PHENIX 4487	Depositor
R, R_{free}	0.153 , 0.191	Depositor
Wilson B-factor (Å ²)	32.5	Xtrriage
Anisotropy	0.015	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	7525	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, NA, 2PG, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/3490	0.61	1/4707 (0.0%)
1	B	0.44	0/3510	0.60	0/4740
All	All	0.44	0/7000	0.61	1/9447 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	403	LEU	CA-CB-CG	-5.25	103.23	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3407	0	3437	14	0
1	B	3411	0	3432	18	0
2	A	11	0	4	0	0
2	B	11	0	4	0	0
3	A	12	0	18	0	0
3	B	12	0	18	2	0
4	A	8	0	6	0	0
4	B	8	0	6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	2	0	0	0	0
5	B	2	0	0	0	0
6	A	320	0	0	3	0
6	B	321	0	0	3	0
All	All	7525	0	6925	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:319[A]:LYS:NZ	6:A:703:HOH:O	2.23	0.71
1:A:240:ASN:ND2	6:A:702:HOH:O	2.20	0.69
3:B:605:EDO:O2	6:B:702:HOH:O	2.10	0.68
1:A:417:LYS:NZ	1:A:421[A]:GLU:OE2	2.30	0.61
1:B:240:ASN:ND2	6:B:704:HOH:O	2.38	0.54
1:B:473:VAL:HG23	1:B:478:ALA:HB3	1.94	0.50
1:A:302[B]:ASN:OD1	6:A:701:HOH:O	2.20	0.49
1:B:253:GLU:HB2	1:B:333:ASP:HB3	1.96	0.48
1:B:178:ASP:OD2	1:B:183:LYS:HA	2.14	0.48
1:B:251[A]:ILE:HG23	1:B:334:ALA:HA	1.95	0.47
1:A:120:GLY:HA3	1:A:460:ARG:NH2	2.29	0.47
1:B:156:GLY:O	1:B:160:GLU:HG3	2.14	0.47
1:A:253:GLU:HB2	1:A:333:ASP:HB3	1.97	0.47
1:B:354:GLU:OE1	3:B:606:EDO:H21	2.15	0.47
1:A:94:ARG:HD3	1:A:295:ASP:OD2	2.16	0.46
1:B:394:THR:HA	1:B:403:LEU:CD1	2.45	0.46
1:B:239[B]:LEU:HG	1:B:278:LEU:HD21	1.97	0.46
1:B:286:ALA:HB3	1:B:290:ALA:HB2	1.99	0.45
1:B:391:LYS:O	1:B:395:GLU:HG3	2.18	0.44
1:A:380:ASP:OD2	1:A:406:ASP:HB3	2.18	0.43
1:B:394:THR:HA	1:B:403:LEU:HD13	2.01	0.42
1:B:443:ASN:O	1:B:447:MET:HG3	2.19	0.42
1:A:156:GLY:O	1:A:160:GLU:HG3	2.19	0.42
1:B:419:GLY:HA2	1:B:424:TRP:CD2	2.55	0.42
1:A:463:GLU:HB2	1:A:490:GLU:HG3	2.01	0.41
1:B:463:GLU:HB2	1:B:490:GLU:HG3	2.02	0.41
1:A:230[B]:ARG:HG3	1:A:231:LEU:N	2.33	0.41
1:A:487:ALA:O	1:A:488:ARG:HB2	2.19	0.41
1:B:446[A]:ARG:HD2	6:B:946:HOH:O	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:183:LYS:HE2	1:B:436:GLY:HA3	2.03	0.41
1:A:94:ARG:HH22	1:A:121:ALA:HA	1.85	0.41
1:B:89:GLU:HG3	1:B:144:VAL:HG22	2.03	0.40
1:A:283:ALA:HB2	1:A:291:LYS:HE2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	446/477 (94%)	435 (98%)	11 (2%)	0	100	100
1	B	449/477 (94%)	438 (98%)	11 (2%)	0	100	100
All	All	895/954 (94%)	873 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	359/384 (94%)	356 (99%)	3 (1%)	81	80
1	B	362/384 (94%)	359 (99%)	3 (1%)	81	80
All	All	721/768 (94%)	715 (99%)	6 (1%)	84	80

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	178	ASP
1	A	231	LEU
1	A	459	HIS
1	B	304[A]	GLU
1	B	304[B]	GLU
1	B	459	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 4 are monoatomic - leaving 12 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

6.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

6.4 Ligands

EDS failed to run properly - this section is therefore empty.

6.5 Other polymers

EDS failed to run properly - this section is therefore empty.