



Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 12:10 am BST

PDB ID : 6UHP
Title : Crystal Structure of C148 mGFP-ncDNA-1
Authors : Winegar, P.W.; Hayes, O.G.; McMillan, J.R.; Figg, C.A.; Focia, P.J.; Mirkin, C.A.
Deposited on : 2019-09-27
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	272	
1	B	272	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 3308 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called C148 mGFP-ncDNA-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	225	Total	C	N	O	S	0	0	0
			1731	1110	292	324	5			
1	B	225	Total	C	N	O	S	0	0	0
			1577	999	266	307	5			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: C148 mGFP-ncDNA-1



• Molecule 1: C148 mGFP-ncDNA-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	59.05Å 51.60Å 100.41Å 90.00° 106.97° 90.00°	Depositor
Resolution (Å)	56.48 – 2.90 56.48 – 2.90	Depositor EDS
% Data completeness (in resolution range)	98.9 (56.48-2.90) 98.8 (56.48-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.15	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.69 (at 2.91Å)	Xtriage
Refinement program	REFMAC 5.8.0257	Depositor
R, R_{free}	0.340 , 0.373 0.333 , 0.356	Depositor DCC
R_{free} test set	598 reflections (4.61%)	wwPDB-VP
Wilson B-factor (Å ²)	54.6	Xtriage
Anisotropy	0.327	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 61.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.37$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtriage
Reported twinning fraction	0.853 for H, K, L 0.147 for -H, -K, H+L	Depositor
Outliers	17 of 12971 reflections (0.131%)	Xtriage
F_o, F_c correlation	0.64	EDS
Total number of atoms	3308	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CRO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.70	0/1748	0.88	0/2373
1	B	0.72	0/1589	0.87	1/2174 (0.0%)
All	All	0.71	0/3337	0.87	1/4547 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	197	PRO	N-CA-CB	-5.44	96.61	102.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1731	0	1631	44	0
1	B	1577	0	1318	24	0
All	All	3308	0	2949	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:57:PRO:O	1:A:60:THR:HG22	1.13	1.23
1:A:57:PRO:O	1:A:60:THR:CG2	2.03	1.06
1:A:57:PRO:C	1:A:60:THR:HG22	1.97	0.84
1:B:38:ALA:O	1:B:41:GLY:N	2.20	0.70
1:A:119:THR:O	1:A:121:VAL:HG23	1.95	0.67
1:B:197:PRO:HB2	1:B:200:HIS:CE1	2.29	0.66
1:B:49:LEU:HD11	1:B:54:LEU:CB	2.26	0.65
1:B:61:LEU:O	1:B:64:THR:N	2.30	0.65
1:A:60:THR:OG1	1:A:107:TYR:OH	2.15	0.62
1:A:19:LEU:HD23	1:A:124:ILE:HB	1.82	0.61
1:A:24:ASN:ND2	1:A:138:LEU:HD11	2.17	0.59
1:B:197:PRO:HB2	1:B:200:HIS:HE1	1.67	0.59
1:A:57:PRO:C	1:A:60:THR:CG2	2.65	0.57
1:A:148:CYS:SG	1:A:205:GLN:HG2	2.45	0.56
1:A:17:VAL:HG13	1:A:122:ASN:HB3	1.88	0.55
1:B:45:LEU:HD12	1:B:66:CRO:CG1	2.37	0.55
1:B:122:ASN:ND2	1:B:124:ILE:HD11	2.22	0.54
1:A:42:LYS:HG3	1:A:224:PHE:CE1	2.45	0.52
1:B:150:ASN:HB3	1:B:201:TYR:CE2	2.44	0.52
1:A:119:THR:O	1:A:121:VAL:CG2	2.58	0.52
1:A:169:ARG:HG2	1:A:177:VAL:HG11	1.91	0.52
1:A:151:VAL:O	1:A:201:TYR:HA	2.10	0.51
1:A:174:ASP:OD1	1:A:175:GLY:N	2.42	0.51
1:A:62:VAL:HG13	1:A:221:LEU:HD22	1.93	0.51
1:B:178:GLN:NE2	1:B:179:LEU:O	2.42	0.50
1:B:57:PRO:O	1:B:60:THR:OG1	2.26	0.49
1:A:116:GLU:OE1	1:A:123:ARG:NH2	2.45	0.49
1:B:61:LEU:O	1:B:64:THR:HB	2.12	0.49
1:B:45:LEU:HD12	1:B:66:CRO:HG11	1.92	0.49
1:B:45:LEU:CD1	1:B:66:CRO:HG13	2.43	0.48
1:A:57:PRO:HD2	1:A:60:THR:CG2	2.44	0.48
1:A:59:PRO:O	1:A:61:LEU:N	2.48	0.47
1:A:111:ALA:HA	1:A:123:ARG:O	2.14	0.47
1:A:229:GLY:O	1:A:230:ILE:HG23	2.15	0.47
1:A:16:LEU:O	1:A:121:VAL:HA	2.15	0.47
1:A:45:LEU:HD13	1:A:47:PHE:CZ	2.50	0.46
1:B:198:ASP:O	1:B:200:HIS:ND1	2.48	0.46
1:A:30:VAL:HG23	1:A:49:LEU:HD23	1.97	0.46
1:A:122:ASN:ND2	1:A:124:ILE:HD11	2.31	0.46
1:A:57:PRO:HD2	1:A:60:THR:HG21	1.97	0.45
1:B:29:SER:N	1:B:51:THR:OG1	2.49	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:19:LEU:O	1:A:29:SER:HA	2.16	0.45
1:A:34:GLY:HA3	1:A:45:LEU:HD23	1.98	0.45
1:B:155:ALA:HB2	1:B:197:PRO:O	2.16	0.45
1:A:59:PRO:HA	1:A:62:VAL:HG23	1.99	0.45
1:A:9:PHE:O	1:A:38:ALA:HB3	2.18	0.44
1:A:19:LEU:CD2	1:A:124:ILE:HB	2.46	0.44
1:B:95:GLN:HG3	1:B:186:ASN:ND2	2.32	0.43
1:A:152:TYR:CD1	1:A:201:TYR:HB3	2.53	0.43
1:A:24:ASN:HD21	1:A:131:PHE:HB2	1.82	0.43
1:B:99:ILE:HA	1:B:181:ASP:O	2.18	0.43
1:B:95:GLN:HG3	1:B:186:ASN:HD21	1.82	0.43
1:B:45:LEU:CD1	1:B:66:CRO:CG1	2.96	0.43
1:B:49:LEU:CD1	1:B:54:LEU:CB	2.95	0.43
1:A:106:ASN:N	1:A:106:ASN:HD22	2.18	0.42
1:A:172:ILE:HB	1:A:174:ASP:OD1	2.19	0.42
1:A:97:ARG:NE	1:A:184:GLN:OE1	2.52	0.42
1:A:229:GLY:O	1:A:230:ILE:CG2	2.67	0.42
1:A:23:VAL:HG22	1:A:128:GLY:HA3	2.01	0.41
1:A:34:GLY:N	1:A:45:LEU:HD23	2.36	0.41
1:B:19:LEU:HA	1:B:124:ILE:O	2.20	0.41
1:A:97:ARG:HG2	1:A:184:GLN:HB2	2.03	0.41
1:B:34:GLY:HA3	1:B:44:THR:O	2.21	0.41
1:A:79:MET:HE1	1:A:228:ALA:HA	2.03	0.41
1:A:54:LEU:HD12	1:A:55:PRO:HD2	2.03	0.41
1:A:57:PRO:CD	1:A:60:THR:HG21	2.52	0.40
1:A:28:PHE:HB2	1:A:49:LEU:HD21	2.04	0.40
1:B:120:LEU:HD13	1:B:121:VAL:N	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	220/272 (81%)	192 (87%)	25 (11%)	3 (1%)	11	36
1	B	220/272 (81%)	190 (86%)	27 (12%)	3 (1%)	11	36
All	All	440/544 (81%)	382 (87%)	52 (12%)	6 (1%)	11	36

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	60	THR
1	A	117	GLY
1	B	39	THR
1	B	176	SER
1	A	136	ASN
1	B	77	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	174/236 (74%)	156 (90%)	18 (10%)	7	22
1	B	135/236 (57%)	122 (90%)	13 (10%)	8	25
All	All	309/472 (66%)	278 (90%)	31 (10%)	7	23

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	GLU
1	A	26	HIS
1	A	49	LEU
1	A	50	THR
1	A	60	THR
1	A	87	SER
1	A	97	ARG
1	A	104	ASP
1	A	106	ASN
1	A	109	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	115	PHE
1	A	120	LEU
1	A	123	ARG
1	A	146	TYR
1	A	162	ILE
1	A	187	THR
1	A	204	THR
1	A	213	ASN
1	B	29	SER
1	B	48	ILE
1	B	77	ASP
1	B	80	LYS
1	B	87	SER
1	B	115	PHE
1	B	129	ILE
1	B	181	ASP
1	B	197	PRO
1	B	201	TYR
1	B	206	SER
1	B	209	SER
1	B	226	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	ASN
1	A	106	ASN
1	B	147	ASN
1	B	150	ASN
1	B	186	ASN
1	B	213	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CRO	A	66	1	23,23,24	3.21	4 (17%)	30,32,34	2.41	5 (16%)
1	CRO	B	66	1	23,23,24	3.23	5 (21%)	30,32,34	2.41	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRO	A	66	1	-	0/12/31/32	0/2/2/2
1	CRO	B	66	1	-	0/12/31/32	0/2/2/2

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	66	CRO	CB2-CA2	12.27	1.45	1.35
1	A	66	CRO	CB2-CA2	12.21	1.45	1.35
1	B	66	CRO	CA2-C2	-6.20	1.42	1.48
1	A	66	CRO	CA2-C2	-6.15	1.42	1.48
1	B	66	CRO	C1-N2	4.95	1.39	1.32
1	A	66	CRO	C1-N2	4.90	1.39	1.32
1	B	66	CRO	C2-N3	-2.62	1.33	1.39
1	A	66	CRO	C2-N3	-2.61	1.33	1.39
1	B	66	CRO	CE1-CZ	2.01	1.42	1.38

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	CRO	O2-C2-CA2	-7.28	126.87	130.96
1	B	66	CRO	O2-C2-CA2	-7.19	126.92	130.96
1	B	66	CRO	CA2-C2-N3	7.18	106.76	103.37
1	A	66	CRO	CA2-C2-N3	7.11	106.73	103.37
1	B	66	CRO	CA2-N2-C1	-5.18	101.96	105.77
1	A	66	CRO	CA2-N2-C1	-5.13	101.99	105.77
1	B	66	CRO	C1-CA1-N1	-3.23	104.72	109.96

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	66	CRO	C1-CA1-N1	-3.22	104.74	109.96
1	B	66	CRO	O3-C3-CA3	-2.98	117.40	126.39
1	A	66	CRO	O3-C3-CA3	-2.98	117.40	126.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	66	CRO	4	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	224/272 (82%)	0.52	8 (3%) 42 37	5, 42, 69, 91	0
1	B	224/272 (82%)	1.13	36 (16%) 1 1	5, 68, 111, 150	0
All	All	448/544 (82%)	0.83	44 (9%) 7 5	5, 52, 98, 150	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	103	ASP	6.3
1	B	141	LYS	5.2
1	B	131	PHE	4.8
1	B	217	ASP	4.7
1	A	99	ILE	4.6
1	B	134	ASP	4.5
1	B	88	ALA	4.3
1	B	160	ASN	3.7
1	B	173	GLU	3.6
1	B	192	GLY	3.6
1	B	181	ASP	3.4
1	B	109	THR	3.1
1	B	55	PRO	3.1
1	B	170	HIS	3.1
1	B	183	TYR	3.1
1	B	52	GLY	3.1
1	B	126	LEU	3.0
1	B	228	ALA	2.9
1	B	59	PRO	2.9
1	B	174	ASP	2.9
1	B	161	GLY	2.8
1	B	182	HIS	2.8
1	B	186	ASN	2.7
1	B	105	GLY	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	101	PHE	2.5
1	B	201	TYR	2.5
1	A	56	VAL	2.5
1	B	128	GLY	2.3
1	B	49	LEU	2.3
1	B	142	LEU	2.3
1	B	190	GLY	2.3
1	B	107	TYR	2.2
1	B	162	ILE	2.2
1	B	70	GLN	2.2
1	B	50	THR	2.1
1	B	10	THR	2.1
1	A	51	THR	2.1
1	B	180	ALA	2.1
1	A	140	HIS	2.1
1	A	36	GLY	2.1
1	A	136	ASN	2.1
1	B	101	PHE	2.0
1	A	103	ASP	2.0
1	B	85	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CRO	B	66	22/23	0.84	0.27	6,7,10,12	0
1	CRO	A	66	22/23	0.88	0.24	6,7,10,12	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.