



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 05:51 am BST

PDB ID : 5UIJ
Title : X-ray structure of The FdtF N-formyltransferase from Salmonella enteric O60
in complex with TDP
Authors : Thoden, J.B.; Woodford, C.R.; Holden, H.M.
Deposited on : 2017-01-14
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	405	<div> <div style="width: 83%; background-color: green;"></div> <div style="width: 12%; background-color: yellow;"></div> <div style="width: 5%; background-color: orange;"></div> <div style="width: 2%; background-color: red;"></div> <div style="width: 0%; background-color: grey;"></div> </div> <div>83% 12% ...</div>
1	B	405	<div> <div style="width: 89%; background-color: green;"></div> <div style="width: 9%; background-color: yellow;"></div> <div style="width: 2%; background-color: orange;"></div> <div style="width: 0%; background-color: red;"></div> <div style="width: 0%; background-color: grey;"></div> </div> <div>89% 9% .</div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7136 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Formyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	398	Total	C	N	O	S	0	2	0
			3242	2105	514	611	12			
1	B	397	Total	C	N	O	S	0	1	0
			3230	2097	512	609	12			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	GLY	-	expression tag	UNP U3GK13
A	-5	GLY	-	expression tag	UNP U3GK13
A	-4	GLY	-	expression tag	UNP U3GK13
A	-3	GLY	-	expression tag	UNP U3GK13
A	-2	GLY	-	expression tag	UNP U3GK13
A	-1	GLY	-	expression tag	UNP U3GK13
A	0	HIS	-	expression tag	UNP U3GK13
A	395	ALA	GLU	engineered mutation	UNP U3GK13
B	-6	GLY	-	expression tag	UNP U3GK13
B	-5	GLY	-	expression tag	UNP U3GK13
B	-4	GLY	-	expression tag	UNP U3GK13
B	-3	GLY	-	expression tag	UNP U3GK13
B	-2	GLY	-	expression tag	UNP U3GK13
B	-1	GLY	-	expression tag	UNP U3GK13
B	0	HIS	-	expression tag	UNP U3GK13
B	395	ALA	GLU	engineered mutation	UNP U3GK13

- Molecule 2 is THYMIDINE-5'-DIPHOSPHATE (three-letter code: TYD) (formula: $C_{10}H_{16}N_2O_{11}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			25	10	2	11	2		
2	A	1	Total	C	N	O	P	0	0
			25	10	2	11	2		
2	B	1	Total	C	N	O	P	0	0
			25	10	2	11	2		
2	B	1	Total	C	N	O	P	0	0
			25	10	2	11	2		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	2	Total Na 2 2	0	0
4	A	3	Total Na 3 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	271	Total O 271 271	0	0
5	B	284	Total O 284 284	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Formyltransferase



- Molecule 1: Formyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	73.91Å 64.76Å 97.76Å 90.00° 90.54° 90.00°	Depositor
Resolution (Å)	97.75 – 1.90 73.91 – 1.90	Depositor EDS
% Data completeness (in resolution range)	95.6 (97.75-1.90) 95.6 (73.91-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.8.0124	Depositor
R, R_{free}	0.181 , 0.231 0.190 , 0.239	Depositor DCC
R_{free} test set	3552 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	20.9	Xtriage
Anisotropy	0.037	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.019 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7136	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: TYD, NA, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	0/3314	1.02	13/4470 (0.3%)
1	B	0.55	0/3299	0.99	9/4451 (0.2%)
All	All	0.56	0/6613	1.00	22/8921 (0.2%)

There are no bond length outliers.

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	214	ARG	NE-CZ-NH1	13.90	127.25	120.30
1	A	214	ARG	NE-CZ-NH2	-13.15	113.72	120.30
1	B	214	ARG	NE-CZ-NH1	-9.80	115.40	120.30
1	A	93	ILE	CG1-CB-CG2	-9.60	90.28	111.40
1	B	214	ARG	NE-CZ-NH2	9.22	124.91	120.30
1	A	239	SER	N-CA-CB	-7.12	99.81	110.50
1	B	67	ASP	CB-CG-OD1	6.87	124.48	118.30
1	B	353	ASP	CB-CG-OD2	6.72	124.35	118.30
1	A	240	ARG	NE-CZ-NH1	6.60	123.60	120.30
1	B	36	ASP	CB-CG-OD2	6.40	124.06	118.30
1	B	251	ARG	NE-CZ-NH2	-6.22	117.19	120.30
1	A	353	ASP	CB-CG-OD2	6.10	123.79	118.30
1	A	366	ASP	CB-CG-OD1	6.03	123.72	118.30
1	A	284	SER	CB-CA-C	-5.84	99.01	110.10
1	A	10	ASP	CB-CG-OD1	5.70	123.43	118.30
1	A	240	ARG	NE-CZ-NH2	-5.48	117.56	120.30
1	B	179	GLU	OE1-CD-OE2	-5.47	116.74	123.30
1	A	73	SER	N-CA-CB	-5.25	102.63	110.50
1	A	124	LYS	CD-CE-NZ	-5.23	99.68	111.70
1	B	366	ASP	CB-CG-OD1	5.12	122.91	118.30
1	A	366	ASP	CB-CG-OD2	-5.10	113.71	118.30
1	B	17	ARG	NE-CZ-NH1	-5.03	117.78	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3242	0	3283	28	0
1	B	3230	0	3265	12	0
2	A	50	0	26	1	0
2	B	50	0	26	1	0
3	A	4	0	6	0	0
4	A	3	0	0	0	0
4	B	2	0	0	0	0
5	A	271	0	0	2	0
5	B	284	0	0	1	0
All	All	7136	0	6606	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73:SER:HB3	1:A:92:ASN:HD22	1.34	0.88
1:A:239:SER:OG	1:A:264:ASP:OD1	1.94	0.84
1:A:93:ILE:HD13	1:A:163:VAL:CG2	2.20	0.71
1:A:19:LEU:HD21	1:A:167:ILE:HD13	1.75	0.68
1:A:17:ARG:HG2	1:A:49:TYR:OH	1.95	0.65
1:A:93:ILE:HD13	1:A:163:VAL:HG23	1.77	0.65
1:B:312:ALA:HA	1:B:320:ILE:HD11	1.78	0.65
1:A:146:GLU:O	1:A:214:ARG:HD2	2.02	0.59
1:A:312:ALA:HA	1:A:320:ILE:HD11	1.89	0.55
1:A:213:LEU:HD11	1:A:267:LEU:HD22	1.90	0.53
1:B:252:ASN:O	1:B:269:LYS:NZ	2.35	0.53
1:A:23:LYS:HB3	1:A:26:ILE:HG13	1.91	0.52
1:A:19:LEU:HD21	1:A:167:ILE:CD1	2.39	0.52
1:A:210:ILE:HD12	1:A:236:VAL:HG23	1.91	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:93:ILE:HD12	1:A:122:LEU:HD13	1.92	0.51
1:A:294:LYS:HE3	1:A:326:LYS:O	2.10	0.51
1:A:302:ILE:HD11	1:A:303:TYR:CZ	2.47	0.50
1:B:75:GLU:OE1	2:B:401:TYD:O3B	2.30	0.50
1:B:210:ILE:HD12	1:B:236:VAL:CG2	2.42	0.49
1:A:127:HIS:HB3	1:B:242:ILE:CD1	2.42	0.49
1:A:75:GLU:OE1	2:A:401:TYD:O1B	2.31	0.48
1:A:302:ILE:HD12	1:A:303:TYR:N	2.29	0.48
1:A:302:ILE:CD1	1:A:303:TYR:CE1	2.98	0.47
1:B:60:LEU:HD11	1:B:80:VAL:HG22	1.97	0.46
1:A:1:MET:HA	1:A:1:MET:CE	2.46	0.46
1:A:78:LYS:NZ	5:A:503:HOH:O	2.37	0.46
1:B:124:LYS:O	1:B:132:GLY:HA3	2.16	0.44
1:A:21:LYS:HB3	1:A:22:LYS:HD2	1.99	0.44
1:A:319:LEU:HD22	1:A:323:LEU:HG	2.00	0.44
1:B:210:ILE:HD12	1:B:236:VAL:HG23	2.00	0.43
1:A:149:LYS:NZ	1:A:220:GLU:OE2	2.52	0.42
1:A:207:PHE:CE1	1:B:184:TYR:HA	2.55	0.42
1:B:23:LYS:HB3	1:B:26:ILE:HG12	2.02	0.42
1:B:1:MET:HG2	5:B:725:HOH:O	2.19	0.41
1:A:93:ILE:CD1	1:A:163:VAL:HG23	2.48	0.41
1:A:261:ILE:HD12	1:B:129:ILE:HD12	2.03	0.41
1:A:318:GLU:HB3	5:A:580:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	398/405 (98%)	383 (96%)	14 (4%)	1 (0%)	41	31
1	B	396/405 (98%)	383 (97%)	12 (3%)	1 (0%)	41	31

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	794/810 (98%)	766 (96%)	26 (3%)	2 (0%)	41	31

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	284	SER
1	B	24	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	360/359 (100%)	333 (92%)	27 (8%)	13	5
1	B	358/359 (100%)	345 (96%)	13 (4%)	35	26
All	All	718/718 (100%)	678 (94%)	40 (6%)	21	11

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	21	LYS
1	A	23	LYS
1	A	24	LYS
1	A	73	SER
1	A	83[A]	SER
1	A	83[B]	SER
1	A	93	ILE
1	A	124	LYS
1	A	138	LYS
1	A	173	SER
1	A	174	GLU
1	A	191	LYS
1	A	218	PHE
1	A	239	SER
1	A	243	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	244	LYS
1	A	253	ASP
1	A	261	ILE
1	A	291	ILE
1	A	307	PRO
1	A	319	LEU
1	A	345	LYS
1	A	349	LEU
1	A	374	ASP
1	A	383	LYS
1	A	387	LYS
1	B	2	LYS
1	B	24	LYS
1	B	42	PHE
1	B	54	LEU
1	B	218	PHE
1	B	220	GLU
1	B	273	LYS
1	B	276	LEU
1	B	292	ARG
1	B	336	LYS
1	B	345	LYS
1	B	370	THR
1	B	383	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	GLN
1	A	92	ASN
1	B	92	ASN
1	B	143	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 5 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	TYD	B	401	-	23,26,26	2.25	3 (13%)	33,40,40	2.02	9 (27%)
2	TYD	B	402	-	23,26,26	2.28	2 (8%)	33,40,40	2.17	11 (33%)
3	EDO	A	403	-	3,3,3	0.36	0	2,2,2	0.75	0
2	TYD	A	402	-	23,26,26	2.18	2 (8%)	33,40,40	2.27	12 (36%)
2	TYD	A	401	-	23,26,26	2.16	2 (8%)	33,40,40	2.43	13 (39%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TYD	B	401	-	-	0/16/28/28	0/2/2/2
2	TYD	B	402	-	-	6/16/28/28	0/2/2/2
3	EDO	A	403	-	-	1/1/1/1	-
2	TYD	A	402	-	-	7/16/28/28	0/2/2/2
2	TYD	A	401	-	-	2/16/28/28	0/2/2/2

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	402	TYD	C6-N1	-9.57	1.34	1.46
2	B	401	TYD	C6-N1	-9.43	1.34	1.46
2	A	402	TYD	C6-N1	-9.03	1.35	1.46
2	A	401	TYD	C6-N1	-8.82	1.35	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	TYD	C6-C5	-4.09	1.39	1.51
2	B	402	TYD	C6-C5	-4.02	1.39	1.51
2	A	402	TYD	C6-C5	-3.82	1.39	1.51
2	B	401	TYD	C6-C5	-3.66	1.40	1.51
2	B	401	TYD	C1'-N1	2.23	1.48	1.45

All (45) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	TYD	C4-N3-C2	-6.21	119.27	126.86
2	B	402	TYD	N3-C2-N1	5.86	122.85	116.65
2	B	402	TYD	C4-N3-C2	-5.58	120.03	126.86
2	A	401	TYD	C5-C6-N1	5.23	121.40	111.11
2	B	401	TYD	C4-N3-C2	-4.91	120.86	126.86
2	B	401	TYD	C5-C6-N1	4.84	120.63	111.11
2	A	402	TYD	C4-N3-C2	-4.74	121.06	126.86
2	A	401	TYD	N3-C2-N1	4.74	121.67	116.65
2	B	402	TYD	C5-C6-N1	4.71	120.38	111.11
2	B	401	TYD	C5M-C5-C6	4.64	122.25	112.34
2	A	402	TYD	C5-C6-N1	4.50	119.98	111.11
2	A	402	TYD	C5M-C5-C6	4.24	121.39	112.34
2	A	401	TYD	O2-C2-N1	-3.84	118.29	123.11
2	A	402	TYD	PA-O3A-PB	-3.75	119.96	132.83
2	B	401	TYD	N3-C2-N1	3.73	120.60	116.65
2	A	402	TYD	C2'-C1'-N1	-3.68	111.11	115.61
2	A	402	TYD	O2-C2-N1	-3.56	118.64	123.11
2	A	401	TYD	C5M-C5-C6	3.54	119.90	112.34
2	A	401	TYD	PA-O3A-PB	-3.53	120.71	132.83
2	A	402	TYD	N3-C2-N1	3.49	120.34	116.65
2	A	401	TYD	C1'-N1-C2	2.94	122.40	118.50
2	A	402	TYD	O3'-C3'-C2'	-2.83	100.76	110.90
2	B	401	TYD	C1'-N1-C2	2.73	122.12	118.50
2	B	402	TYD	C6-C5-C4	2.73	119.36	111.53
2	A	401	TYD	O2B-PB-O1B	2.72	121.32	110.68
2	A	402	TYD	C2'-C3'-C4'	2.64	108.27	102.76
2	B	402	TYD	O4'-C1'-N1	2.59	111.71	108.41
2	A	401	TYD	O3A-PB-O1B	-2.58	96.90	111.19
2	A	402	TYD	C6-C5-C4	2.56	118.88	111.53
2	A	401	TYD	C3'-C2'-C1'	2.46	108.69	102.54
2	A	401	TYD	O2A-PA-O1A	2.45	124.36	112.24
2	B	402	TYD	O2-C2-N3	-2.45	116.94	121.50
2	B	401	TYD	O3B-PB-O1B	2.44	120.22	110.68

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	402	TYD	O3A-PB-O1B	-2.43	97.72	111.19
2	B	402	TYD	O2-C2-N1	-2.31	120.20	123.11
2	B	401	TYD	C6-C5-C4	2.29	118.11	111.53
2	A	402	TYD	O3B-PB-O3A	-2.29	96.97	104.64
2	B	402	TYD	C5M-C5-C6	2.27	117.19	112.34
2	A	401	TYD	C6-C5-C4	2.25	117.97	111.53
2	B	401	TYD	O3B-PB-O2B	2.19	116.01	107.64
2	B	402	TYD	O3'-C3'-C2'	2.15	118.60	110.90
2	A	401	TYD	C2'-C1'-N1	2.13	118.21	115.61
2	B	402	TYD	O3B-PB-O2B	2.06	115.50	107.64
2	B	402	TYD	C2'-C1'-N1	-2.04	113.11	115.61
2	B	401	TYD	C3'-C2'-C1'	2.02	107.61	102.54

There are no chirality outliers.

All (16) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	402	TYD	C5'-O5'-PA-O1A
2	A	402	TYD	C5'-O5'-PA-O2A
2	A	402	TYD	C5'-O5'-PA-O3A
2	B	402	TYD	PA-O3A-PB-O3B
2	A	402	TYD	O4'-C4'-C5'-O5'
2	A	402	TYD	C3'-C4'-C5'-O5'
3	A	403	EDO	O1-C1-C2-O2
2	B	402	TYD	PB-O3A-PA-O5'
2	B	402	TYD	PA-O3A-PB-O2B
2	A	401	TYD	PB-O3A-PA-O1A
2	B	402	TYD	C2'-C1'-N1-C6
2	A	402	TYD	PB-O3A-PA-O1A
2	A	402	TYD	PB-O3A-PA-O2A
2	A	401	TYD	PB-O3A-PA-O2A
2	B	402	TYD	O4'-C1'-N1-C2
2	B	402	TYD	PA-O3A-PB-O1B



There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	TYD	1	0
2	A	401	TYD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	398/405 (98%)	-0.36	2 (0%) 91 92	14, 23, 42, 71	0
1	B	397/405 (98%)	-0.35	0 100 100	13, 23, 44, 57	0
All	All	795/810 (98%)	-0.35	2 (0%) 94 94	13, 23, 43, 71	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	145	PHE	2.5
1	A	1	MET	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	TYD	A	402	25/25	0.93	0.11	26,33,50,53	0
3	EDO	A	403	4/4	0.93	0.11	31,31,35,35	0
2	TYD	B	402	25/25	0.95	0.08	22,28,34,36	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	NA	B	403	1/1	0.95	0.09	29,29,29,29	0
2	TYD	A	401	25/25	0.95	0.10	16,23,43,46	0
4	NA	A	406	1/1	0.96	0.17	32,32,32,32	0
2	TYD	B	401	25/25	0.96	0.09	17,22,36,46	0
4	NA	A	404	1/1	0.97	0.08	25,25,25,25	0
4	NA	A	405	1/1	0.99	0.07	27,27,27,27	0
4	NA	B	404	1/1	0.99	0.08	23,23,23,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around TYD B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around TYD A 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around TYD B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.