



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 28, 2020 – 11:35 AM EDT

PDB ID : 6UK1
Title : Crystal structure of nucleotide-binding domain 2 (NBD2) of the human Cystic Fibrosis Transmembrane Conductance Regulator (CFTR)
Authors : Wang, C.; Vorobiev, S.M.; Vernon, R.M.; Khazanov, N.; Senderowitz, H.; Forman-Kay, J.D.; Hunt, J.F.
Deposited on : 2019-10-03
Resolution : 2.69 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.6
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.6


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	229	<div> <div>3%</div> <div>90%</div> <div>10%</div> </div>
1	B	229	<div> <div>10%</div> <div>88%</div> <div>12%</div> </div>
1	C	229	<div> <div>5%</div> <div>87%</div> <div>13%</div> </div>
1	D	229	<div> <div>3%</div> <div>86%</div> <div>13%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7231 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cystic fibrosis transmembrane conductance regulator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	229	Total	C	N	O	S	0	0	0
			1767	1129	299	331	8			
1	B	228	Total	C	N	O	S	0	0	0
			1757	1122	295	332	8			
1	C	229	Total	C	N	O	S	0	0	0
			1756	1122	297	329	8			
1	D	226	Total	C	N	O	S	0	1	0
			1749	1115	299	327	8			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1255	LEU	SER	engineered mutation	UNP P13569
A	1280	GLU	GLN	engineered mutation	UNP P13569
A	1292	ASP	LYS	engineered mutation	UNP P13569
A	1307	ASN	TYR	engineered mutation	UNP P13569
A	1334	GLY	LYS	engineered mutation	UNP P13569
A	1359	ALA	SER	engineered mutation	UNP P13569
A	1402	ALA	HIS	engineered mutation	UNP P13569
A	1411	ASP	GLN	engineered mutation	UNP P13569
B	1255	LEU	SER	engineered mutation	UNP P13569
B	1280	GLU	GLN	engineered mutation	UNP P13569
B	1292	ASP	LYS	engineered mutation	UNP P13569
B	1307	ASN	TYR	engineered mutation	UNP P13569
B	1334	GLY	LYS	engineered mutation	UNP P13569
B	1359	ALA	SER	engineered mutation	UNP P13569
B	1402	ALA	HIS	engineered mutation	UNP P13569
B	1411	ASP	GLN	engineered mutation	UNP P13569
C	1255	LEU	SER	engineered mutation	UNP P13569
C	1280	GLU	GLN	engineered mutation	UNP P13569
C	1292	ASP	LYS	engineered mutation	UNP P13569
C	1307	ASN	TYR	engineered mutation	UNP P13569
C	1334	GLY	LYS	engineered mutation	UNP P13569

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	1359	ALA	SER	engineered mutation	UNP P13569
C	1402	ALA	HIS	engineered mutation	UNP P13569
C	1411	ASP	GLN	engineered mutation	UNP P13569
D	1255	LEU	SER	engineered mutation	UNP P13569
D	1280	GLU	GLN	engineered mutation	UNP P13569
D	1292	ASP	LYS	engineered mutation	UNP P13569
D	1307	ASN	TYR	engineered mutation	UNP P13569
D	1334	GLY	LYS	engineered mutation	UNP P13569
D	1359	ALA	SER	engineered mutation	UNP P13569
D	1402	ALA	HIS	engineered mutation	UNP P13569
D	1411	ASP	GLN	engineered mutation	UNP P13569

- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	C	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	D	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Mg 1	0	0
3	A	1	Total 1	Mg 1	0	0
3	D	1	Total 1	Mg 1	0	0
3	C	1	Total 1	Mg 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	25	Total 25	O 25	0	0
4	B	18	Total 18	O 18	0	0
4	C	15	Total 15	O 15	0	0
4	D	16	Total 16	O 16	0	0



3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Cystic fibrosis transmembrane conductance regulator




- Molecule 1: Cystic fibrosis transmembrane conductance regulator




- Molecule 1: Cystic fibrosis transmembrane conductance regulator



- Molecule 1: Cystic fibrosis transmembrane conductance regulator





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	95.06Å 60.13Å 105.16Å 90.00° 99.57° 90.00°	Depositor
Resolution (Å)	64.41 – 2.69 64.41 – 2.69	Depositor EDS
% Data completeness (in resolution range)	98.1 (64.41-2.69) 98.1 (64.41-2.69)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 2.69Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.273 , 0.304 0.273 , 0.303	Depositor DCC
R_{free} test set	1552 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	47.0	Xtriage
Anisotropy	0.190	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 59.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7231	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 84.51 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.6312e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.23	0/1799	0.41	0/2439
1	B	0.24	0/1789	0.40	0/2426
1	C	0.23	0/1788	0.40	0/2427
1	D	0.23	0/1784	0.41	0/2419
All	All	0.24	0/7160	0.40	0/9711

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1767	0	1761	13	0
1	B	1757	0	1743	14	0
1	C	1756	0	1737	14	0
1	D	1749	0	1735	17	0
2	A	31	0	12	0	0
2	B	31	0	12	0	0
2	C	31	0	12	1	0
2	D	31	0	12	0	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	25	0	0	0	0
4	B	18	0	0	1	0
4	C	15	0	0	1	0
4	D	16	0	0	0	0
All	All	7231	0	7024	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1216:THR:HB	1:C:1264:GLU:HB2	1.73	0.69
2:C:1501:ATP:O1B	4:C:1601:HOH:O	2.11	0.68
1:A:1209:GLN:HB2	1:A:1236:PRO:HG3	1.82	0.62
1:B:1209:GLN:HB2	1:B:1236:PRO:HG3	1.83	0.60
1:C:1277:ILE:HD11	1:C:1282:TRP:HE3	1.65	0.60
1:B:1277:ILE:HD11	1:B:1282:TRP:HE3	1.68	0.59
1:D:1412:GLN:OE1	1:D:1423:GLN:NE2	2.31	0.57
1:D:1356:LEU:HD22	1:D:1384:ILE:HD13	1.86	0.57
1:A:1405:GLU:HG3	1:B:1408:LEU:HD11	1.87	0.56
1:C:1321:GLU:O	1:C:1386:ARG:NH2	2.38	0.56
1:C:1409:GLU:HG3	1:D:1409:GLU:HG3	1.87	0.56
1:A:1277:ILE:HD11	1:A:1282:TRP:HE3	1.72	0.55
1:A:1412:GLN:OE1	1:A:1423:GLN:NE2	2.37	0.55
1:B:1251:SER:OG	1:B:1370:ASP:OD2	2.24	0.55
1:D:1288:VAL:O	1:D:1291:GLN:NE2	2.37	0.54
1:A:1215:LEU:HD22	1:A:1267:ILE:HD11	1.89	0.54
1:D:1209:GLN:HB2	1:D:1236:PRO:HG3	1.89	0.53
1:D:1238:GLN:OE1	1:D:1412:GLN:NE2	2.40	0.53
1:B:1405:GLU:HA	1:B:1408:LEU:HD21	1.91	0.53
1:A:1346:LEU:O	1:A:1351:LYS:NZ	2.42	0.53
1:B:1213:LYS:O	1:B:1231:SER:OG	2.17	0.53
1:B:1310:TRP:NE1	1:B:1361:LEU:O	2.41	0.52
1:D:1216:THR:HB	1:D:1264:GLU:HB2	1.92	0.52
1:D:1321:GLU:O	1:D:1386:ARG:NH1	2.44	0.51
1:D:1243:LEU:HB2	1:D:1407:MET:HE1	1.93	0.50
1:C:1258:LEU:HD11	1:C:1288:VAL:HG21	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1230:ILE:HD12	1:B:1421:VAL:HG23	1.94	0.49
1:A:1405:GLU:HA	1:A:1408:LEU:HD21	1.93	0.49
1:D:1258:LEU:HD11	1:D:1288:VAL:HG21	1.94	0.49
1:A:1296:PHE:N	1:A:1303:ASN:OD1	2.43	0.49
1:B:1289:ILE:O	1:B:1369:LEU:HA	2.13	0.49
1:C:1289:ILE:HG22	1:C:1376:LEU:HD11	1.94	0.48
1:D:1290:PRO:HB3	1:D:1376:LEU:HD21	1.94	0.48
1:D:1380:THR:O	1:D:1384:ILE:HG12	2.14	0.48
1:B:1215:LEU:HD22	1:B:1267:ILE:HD11	1.96	0.47
1:C:1324:LEU:O	1:C:1328:ILE:HG12	2.14	0.47
1:D:1324:LEU:O	1:D:1328:ILE:HG12	2.14	0.47
1:A:1290:PRO:HB3	1:A:1376:LEU:HD21	1.96	0.47
1:C:1416:ILE:HG12	1:C:1421:VAL:HG22	1.96	0.47
1:C:1296:PHE:N	1:C:1303:ASN:OD1	2.48	0.46
1:A:1322:VAL:HG21	1:A:1357:ALA:HB2	1.97	0.46
1:B:1289:ILE:HG22	1:B:1376:LEU:HD11	1.98	0.46
1:D:1277:ILE:HD11	1:D:1282:TRP:HE3	1.81	0.44
1:D:1296:PHE:N	1:D:1303:ASN:OD1	2.51	0.43
1:B:1372:PRO:HD2	1:B:1401:GLU:OE2	2.17	0.43
1:C:1372:PRO:HD2	1:C:1401:GLU:OE2	2.18	0.43
1:A:1239:ARG:NH2	1:A:1395:CYS:O	2.52	0.42
1:A:1255:LEU:HB3	1:A:1261:LEU:HD13	2.00	0.42
1:D:1284:LYS:HB2	1:D:1284:LYS:HE3	1.76	0.42
1:D:1278:THR:HB	1:D:1281:GLN:HG2	2.02	0.42
1:C:1281:GLN:HB3	1:C:1281:GLN:HE21	1.68	0.41
1:C:1300:PHE:CE1	1:C:1339:LEU:HD11	2.55	0.41
1:A:1317:LYS:HD2	1:A:1317:LYS:HA	1.81	0.41
1:B:1427:ILE:HD12	1:B:1427:ILE:H	1.85	0.41
1:C:1296:PHE:HE2	1:C:1306:PRO:HB3	1.85	0.41
1:B:1250:LYS:HE2	4:B:1601:HOH:O	2.21	0.40
1:C:1382:GLN:HG2	1:C:1385:ARG:HH22	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	227/229 (99%)	219 (96%)	8 (4%)	0	100	100
1	B	226/229 (99%)	213 (94%)	13 (6%)	0	100	100
1	C	227/229 (99%)	216 (95%)	11 (5%)	0	100	100
1	D	225/229 (98%)	219 (97%)	6 (3%)	0	100	100
All	All	905/916 (99%)	867 (96%)	38 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	190/197 (96%)	189 (100%)	1 (0%)	88	96
1	B	189/197 (96%)	185 (98%)	4 (2%)	53	80
1	C	187/197 (95%)	183 (98%)	4 (2%)	53	80
1	D	187/197 (95%)	186 (100%)	1 (0%)	88	96
All	All	753/788 (96%)	743 (99%)	10 (1%)	69	87

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1274	TRP
1	B	1274	TRP
1	B	1292	ASP
1	B	1360	VAL
1	B	1429	LYS
1	C	1245	ARG
1	C	1274	TRP
1	C	1284	LYS
1	C	1429	LYS
1	D	1274	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ATP	C	1501	3	26,33,33	3.09	7 (26%)	31,52,52	3.72	7 (22%)
2	ATP	D	1501	3	26,33,33	3.10	7 (26%)	31,52,52	3.72	7 (22%)
2	ATP	A	1501	3	26,33,33	3.10	8 (30%)	31,52,52	3.76	8 (25%)
2	ATP	B	1501	3	26,33,33	3.10	8 (30%)	31,52,52	3.77	8 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	C	1501	3	-	5/18/38/38	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	D	1501	3	-	7/18/38/38	0/3/3/3
2	ATP	A	1501	3	-	6/18/38/38	0/3/3/3
2	ATP	B	1501	3	-	3/18/38/38	0/3/3/3

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1501	ATP	C3'-C4'	-9.57	1.28	1.53
2	B	1501	ATP	C3'-C4'	-9.55	1.28	1.53
2	A	1501	ATP	C3'-C4'	-9.55	1.28	1.53
2	D	1501	ATP	C3'-C4'	-9.55	1.28	1.53
2	D	1501	ATP	O4'-C4'	7.90	1.62	1.45
2	B	1501	ATP	O4'-C4'	7.87	1.62	1.45
2	A	1501	ATP	O4'-C4'	7.87	1.62	1.45
2	C	1501	ATP	O4'-C4'	7.84	1.62	1.45
2	B	1501	ATP	O4'-C1'	-6.78	1.31	1.41
2	D	1501	ATP	O4'-C1'	-6.78	1.31	1.41
2	C	1501	ATP	O4'-C1'	-6.73	1.31	1.41
2	A	1501	ATP	O4'-C1'	-6.72	1.31	1.41
2	A	1501	ATP	O3'-C3'	3.37	1.50	1.43
2	B	1501	ATP	O3'-C3'	3.37	1.50	1.43
2	D	1501	ATP	O3'-C3'	3.31	1.50	1.43
2	C	1501	ATP	O3'-C3'	3.30	1.50	1.43
2	D	1501	ATP	C2-N3	3.29	1.37	1.32
2	A	1501	ATP	C2-N3	3.26	1.37	1.32
2	B	1501	ATP	C2-N3	3.26	1.37	1.32
2	C	1501	ATP	C2-N3	3.21	1.37	1.32
2	A	1501	ATP	O2'-C2'	-2.37	1.37	1.43
2	D	1501	ATP	O2'-C2'	-2.37	1.37	1.43
2	C	1501	ATP	O2'-C2'	-2.35	1.37	1.43
2	B	1501	ATP	O2'-C2'	-2.32	1.37	1.43
2	C	1501	ATP	C2-N1	2.28	1.38	1.33
2	D	1501	ATP	C2-N1	2.26	1.38	1.33
2	A	1501	ATP	C2-N1	2.24	1.38	1.33
2	B	1501	ATP	C2-N1	2.23	1.38	1.33
2	B	1501	ATP	C6-N6	2.02	1.41	1.34
2	A	1501	ATP	C6-N6	2.01	1.41	1.34

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1501	ATP	C5-C6-N6	14.28	142.05	120.35
2	C	1501	ATP	C5-C6-N6	14.24	142.00	120.35
2	B	1501	ATP	C5-C6-N6	14.24	141.99	120.35
2	D	1501	ATP	C5-C6-N6	14.19	141.92	120.35
2	A	1501	ATP	N6-C6-N1	-9.88	98.06	118.57
2	B	1501	ATP	N6-C6-N1	-9.85	98.13	118.57
2	C	1501	ATP	N6-C6-N1	-9.83	98.17	118.57
2	D	1501	ATP	N6-C6-N1	-9.81	98.22	118.57
2	C	1501	ATP	C1'-N9-C4	-8.09	112.43	126.64
2	D	1501	ATP	C1'-N9-C4	-7.99	112.61	126.64
2	B	1501	ATP	C1'-N9-C4	-7.96	112.66	126.64
2	A	1501	ATP	C1'-N9-C4	-7.93	112.71	126.64
2	C	1501	ATP	N3-C2-N1	-5.57	119.98	128.68
2	A	1501	ATP	N3-C2-N1	-5.51	120.06	128.68
2	D	1501	ATP	N3-C2-N1	-5.50	120.08	128.68
2	B	1501	ATP	N3-C2-N1	-5.50	120.08	128.68
2	B	1501	ATP	C3'-C2'-C1'	3.77	106.65	100.98
2	A	1501	ATP	C3'-C2'-C1'	3.53	106.29	100.98
2	D	1501	ATP	PA-O3A-PB	-3.05	122.36	132.83
2	C	1501	ATP	PA-O3A-PB	-3.00	122.53	132.83
2	B	1501	ATP	PB-O3B-PG	-2.82	123.14	132.83
2	A	1501	ATP	PA-O3A-PB	-2.74	123.43	132.83
2	D	1501	ATP	C3'-C2'-C1'	2.57	104.85	100.98
2	B	1501	ATP	C2'-C3'-C4'	2.45	107.39	102.64
2	B	1501	ATP	PA-O3A-PB	-2.37	124.69	132.83
2	A	1501	ATP	C2'-C3'-C4'	2.25	107.01	102.64
2	C	1501	ATP	PB-O3B-PG	-2.23	125.18	132.83
2	A	1501	ATP	PB-O3B-PG	-2.22	125.22	132.83
2	C	1501	ATP	C3'-C2'-C1'	2.20	104.29	100.98
2	D	1501	ATP	PB-O3B-PG	-2.08	125.70	132.83

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	1501	ATP	C5'-O5'-PA-O3A
2	D	1501	ATP	PB-O3B-PG-O2G
2	D	1501	ATP	PB-O3B-PG-O3G
2	D	1501	ATP	C5'-O5'-PA-O3A
2	A	1501	ATP	PB-O3B-PG-O2G
2	A	1501	ATP	C5'-O5'-PA-O3A
2	B	1501	ATP	PB-O3B-PG-O3G
2	C	1501	ATP	O4'-C4'-C5'-O5'

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	C	1501	ATP	C3'-C4'-C5'-O5'
2	D	1501	ATP	O4'-C4'-C5'-O5'
2	D	1501	ATP	C3'-C4'-C5'-O5'
2	A	1501	ATP	O4'-C4'-C5'-O5'
2	C	1501	ATP	C5'-O5'-PA-O2A
2	D	1501	ATP	C5'-O5'-PA-O2A
2	A	1501	ATP	C5'-O5'-PA-O1A
2	A	1501	ATP	C3'-C4'-C5'-O5'
2	B	1501	ATP	PB-O3A-PA-O2A
2	C	1501	ATP	C5'-O5'-PA-O1A
2	D	1501	ATP	C5'-O5'-PA-O1A
2	A	1501	ATP	C5'-O5'-PA-O2A
2	B	1501	ATP	C5'-O5'-PA-O1A


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1501	ATP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	229/229 (100%)	0.28	6 (2%) 56 57	21, 42, 67, 88	0
1	B	228/229 (99%)	0.86	24 (10%) 6 4	48, 62, 83, 103	0
1	C	229/229 (100%)	0.61	11 (4%) 30 28	42, 62, 78, 90	0
1	D	226/229 (98%)	0.18	6 (2%) 54 55	15, 40, 66, 84	0
All	All	912/916 (99%)	0.48	47 (5%) 27 25	15, 52, 80, 103	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1417	GLU	6.7
1	B	1294	PHE	4.7
1	D	1223	GLY	4.1
1	C	1416	ILE	4.0
1	A	1347	SER	3.8
1	A	1417	GLU	3.2
1	C	1294	PHE	3.1
1	B	1289	ILE	3.0
1	A	1430	LEU	2.9
1	C	1288	VAL	2.9
1	A	1416	ILE	2.8
1	B	1346	LEU	2.8
1	C	1422	ARG	2.7
1	A	1289	ILE	2.7
1	C	1362	SER	2.6
1	C	1220	THR	2.6
1	B	1247	GLY	2.6
1	B	1248	SER	2.5
1	D	1425	ASP	2.4
1	D	1416	ILE	2.4
1	B	1347	SER	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	1287	GLY	2.4
1	A	1420	LYS	2.4
1	B	1227	LEU	2.3
1	C	1253	LEU	2.3
1	C	1217	ALA	2.3
1	B	1428	GLN	2.3
1	B	1416	ILE	2.3
1	B	1327	VAL	2.3
1	B	1297	SER	2.3
1	B	1231	SER	2.2
1	B	1290	PRO	2.2
1	C	1256	ALA	2.2
1	B	1268	GLN	2.1
1	B	1271	GLY	2.1
1	B	1257	PHE	2.1
1	B	1420	LYS	2.1
1	B	1421	VAL	2.1
1	D	1249	GLY	2.1
1	B	1354	MET	2.1
1	C	1252	THR	2.1
1	B	1228	GLU	2.1
1	D	1418	GLU	2.1
1	B	1293	VAL	2.1
1	B	1361	LEU	2.0
1	B	1315	ILE	2.0
1	D	1347	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MG	B	1502	1/1	0.22	0.38	69,69,69,69	0
3	MG	C	1502	1/1	0.85	0.17	42,42,42,42	0
3	MG	A	1502	1/1	0.92	0.09	20,20,20,20	0
2	ATP	B	1501	31/31	0.94	0.17	24,45,53,60	0
2	ATP	C	1501	31/31	0.94	0.17	12,41,60,77	0
3	MG	D	1502	1/1	0.96	0.14	13,13,13,13	0
2	ATP	D	1501	31/31	0.96	0.18	8,30,43,54	0
2	ATP	A	1501	31/31	0.97	0.16	5,28,41,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ATP B 1501:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ATP C 1501:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ATP D 1501:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ATP A 1501:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.