



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 28, 2020 – 12:17 PM EDT

PDB ID : 6UK6
Title : Crystal structure of human GAC in complex with inhibitor UPGL00018
Authors : Huang, Q.; Cerione, R.A.
Deposited on : 2019-10-04
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.6
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.6


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	527	<div> <div> <div></div> <div>48%</div> <div>26%</div> <div>•</div> <div>23%</div> </div> </div>
1	B	527	<div> <div> <div></div> <div>46%</div> <div>27%</div> <div>•</div> <div>24%</div> </div> </div>
1	C	527	<div> <div> <div></div> <div>48%</div> <div>26%</div> <div>•</div> <div>24%</div> </div> </div>
1	D	527	<div> <div> <div></div> <div>49%</div> <div>23%</div> <div>•</div> <div>25%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 12524 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glutaminase kidney isoform, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	404	Total	C	N	O	S	1	0	0
			3151	2006	532	585	28			
1	B	398	Total	C	N	O	S	1	0	0
			3108	1983	525	572	28			
1	C	402	Total	C	N	O	S	1	0	0
			3128	1988	530	582	28			
1	D	397	Total	C	N	O	S	1	0	0
			3099	1975	524	572	28			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	268	ALA	VAL	conflict	UNP O94925
B	268	ALA	VAL	conflict	UNP O94925
C	268	ALA	VAL	conflict	UNP O94925
D	268	ALA	VAL	conflict	UNP O94925


- Molecule 2 is 5-{4-[(5-amino-1,3,4-thiadiazol-2-yl)oxy]piperidin-1-yl}-1,3,4-thiadiazol-2-amine (three-letter code: Q9S) (formula: C₉H₁₃N₇OS₂) (labeled as "Ligand of Interest" by author).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			19	9	7	1	2		
2	D	1	Total	C	N	O	S	0	0
			19	9	7	1	2		

- Molecule 1: Glutaminase kidney isoform, mitochondrial







4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.44Å 138.79Å 179.05Å 90.00° 89.99° 90.00°	Depositor
Resolution (Å)	22.05 – 2.90 22.05 – 2.90	Depositor EDS
% Data completeness (in resolution range)	95.7 (22.05-2.90) 95.7 (22.05-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.86 (at 2.89Å)	Xtriage
Refinement program	PHENIX 1.15.2_3472	Depositor
R, R_{free}	0.239 , 0.308 0.239 , 0.308	Depositor DCC
R_{free} test set	1603 reflections (2.84%)	wwPDB-VP
Wilson B-factor (Å ²)	60.8	Xtriage
Anisotropy	1.005	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 34.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.55$, $\langle L^2 \rangle = 0.40$	Xtriage
Estimated twinning fraction	0.458 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12524	wwPDB-VP
Average B, all atoms (Å ²)	98.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: Q9S

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.62	3/3221 (0.1%)	0.97	21/4347 (0.5%)
1	B	0.59	1/3177 (0.0%)	1.07	24/4284 (0.6%)
1	C	0.56	0/3196	0.93	15/4311 (0.3%)
1	D	0.62	2/3167 (0.1%)	1.15	23/4270 (0.5%)
All	All	0.60	6/12761 (0.0%)	1.03	83/17212 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	3
1	C	0	2
All	All	0	7

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	154	ILE	CA-C	-10.62	1.25	1.52
1	A	307	ARG	CG-CD	6.11	1.67	1.51
1	D	145	TYR	CE2-CZ	-5.52	1.31	1.38
1	D	317	ARG	CD-NE	5.48	1.55	1.46
1	B	241	GLN	CA-CB	-5.45	1.42	1.53

The worst 5 of 83 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	199	LEU	CB-CG-CD2	-20.77	75.69	111.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	387	ARG	NE-CZ-NH2	-19.98	110.31	120.30
1	D	317	ARG	NE-CZ-NH1	-18.89	110.86	120.30
1	A	272	ARG	NE-CZ-NH1	-18.18	111.21	120.30
1	D	387	ARG	NE-CZ-NH1	17.95	129.28	120.30

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	248	ASP	Peptide
1	A	272	ARG	Sidechain
1	B	137	PRO	Peptide
1	B	138	SER	Peptide
1	B	258	PRO	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3151	0	3122	113	0
1	B	3108	0	3086	120	0
1	C	3128	0	3098	124	0
1	D	3099	0	3070	122	0
2	A	19	0	0	2	0
2	D	19	0	0	0	0
All	All	12524	0	12376	446	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 446 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:272:ARG:NH1	1:A:429:VAL:HG22	1.53	1.24
1:C:139:LEU:CD1	1:C:208:ILE:HG12	1.75	1.16

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:139:LEU:HD13	1:C:208:ILE:CG1	1.79	1.12
1:C:139:LEU:HD13	1:C:208:ILE:HG12	1.23	1.07
1:B:535:HIS:HD2	1:C:439:PHE:CD1	1.76	1.03

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	400/527 (76%)	377 (94%)	20 (5%)	3 (1%)	19	51
1	B	392/527 (74%)	369 (94%)	20 (5%)	3 (1%)	19	51
1	C	398/527 (76%)	375 (94%)	20 (5%)	3 (1%)	19	51
1	D	391/527 (74%)	370 (95%)	19 (5%)	2 (0%)	29	61
All	All	1581/2108 (75%)	1491 (94%)	79 (5%)	11 (1%)	22	54

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	259	ASP
1	D	259	ASP
1	A	193	VAL
1	B	545	GLU
1	C	193	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	350/451 (78%)	332 (95%)	18 (5%)	24	56
1	B	344/451 (76%)	328 (95%)	16 (5%)	26	59
1	C	347/451 (77%)	334 (96%)	13 (4%)	34	68
1	D	343/451 (76%)	324 (94%)	19 (6%)	21	53
All	All	1384/1804 (77%)	1318 (95%)	66 (5%)	25	58

5 of 66 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	380	SER
1	C	240	LYS
1	D	384	SER
1	B	398	LYS
1	B	539	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	529	ASN
1	B	535	HIS
1	C	510	ASN
1	B	461	HIS
1	D	213	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	Q9S	D	601	-	13,21,21	3.54	6 (46%)	8,29,29	1.82	3 (37%)
2	Q9S	A	601	-	13,21,21	3.39	6 (46%)	8,29,29	2.35	4 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	Q9S	D	601	-	-	2/4/18/18	0/3/3/3
2	Q9S	A	601	-	-	2/4/18/18	0/3/3/3

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	601	Q9S	C23-N03	8.51	1.45	1.32
2	A	601	Q9S	C23-N03	7.91	1.44	1.32
2	D	601	Q9S	C02-N03	5.20	1.54	1.46
2	A	601	Q9S	C02-N03	5.01	1.54	1.46
2	A	601	Q9S	C04-N03	4.35	1.53	1.46

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	Q9S	C02-C01-C06	3.96	114.67	110.32
2	A	601	Q9S	C01-C02-N03	3.51	118.34	111.10
2	D	601	Q9S	C01-C02-N03	3.46	118.24	111.10
2	D	601	Q9S	C02-C01-C06	3.04	113.66	110.32
2	A	601	Q9S	C04-C05-C06	-2.65	107.40	110.32

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	601	Q9S	S37-C23-N03-C02
2	D	601	Q9S	S37-C23-N03-C04
2	A	601	Q9S	S37-C23-N03-C02
2	A	601	Q9S	S37-C23-N03-C04

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	Q9S	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	404/527 (76%)	0.06	5 (1%) 79 79	50, 88, 159, 246	0
1	B	398/527 (75%)	0.06	7 (1%) 68 67	52, 86, 163, 206	0
1	C	402/527 (76%)	0.06	9 (2%) 62 59	46, 88, 160, 230	0
1	D	397/527 (75%)	0.03	7 (1%) 68 67	51, 87, 165, 232	0
All	All	1601/2108 (75%)	0.05	28 (1%) 70 69	46, 87, 163, 246	0

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	318	PHE	5.6
1	A	545	GLU	5.0
1	B	249	TYR	4.3
1	B	546	GLY	4.0
1	B	152	GLU	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	Q9S	A	601	19/19	0.95	0.22	58,84,103,110	0
2	Q9S	D	601	19/19	0.96	0.19	48,79,113,117	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Q9S A 601:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Q9S D 601:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.