



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 5, 2020 – 10:37 AM EDT

PDB ID : 6UN5  
Title : Crystal structure of green fluorescent protein (GFP); S65T, Y66(2,3,5-F3Y);  
ih circular permutant (50-51)  
Authors : Lin, C.-Y.; Boxer, S.G.  
Deposited on : 2019-10-10  
Resolution : 1.36 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.14.6  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.14.6


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1509 (1.38-1.34)
Clashscore	141614	1551 (1.38-1.34)
Ramachandran outliers	138981	1530 (1.38-1.34)
Sidechain outliers	138945	1530 (1.38-1.34)
RSRZ outliers	127900	1487 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	252	
1	B	252	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4106 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein, Green fluorescent protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	226	Total	C	F	N	O	S	0	18	0
			1874	1196	3	308	363	4			
1	B	228	Total	C	F	N	O	S	0	20	0
			1903	1213	3	320	362	5			

There are 70 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	MET	-	initiating methionine	UNP A0A059PIQ0
A	-9	GLY	-	expression tag	UNP A0A059PIQ0
A	-8	HIS	-	expression tag	UNP A0A059PIQ0
A	-7	HIS	-	expression tag	UNP A0A059PIQ0
A	-6	HIS	-	expression tag	UNP A0A059PIQ0
A	-5	HIS	-	expression tag	UNP A0A059PIQ0
A	-4	HIS	-	expression tag	UNP A0A059PIQ0
A	-3	HIS	-	expression tag	UNP A0A059PIQ0
A	-2	SER	-	expression tag	UNP A0A059PIQ0
A	-1	SER	-	expression tag	UNP A0A059PIQ0
A	0	GLY	-	expression tag	UNP A0A059PIQ0
A	15	QC4	THR	chromophore	UNP A0A059PIQ0
A	15	QC4	TYR	chromophore	UNP A0A059PIQ0
A	15	QC4	GLY	chromophore	UNP A0A059PIQ0
A	22	SER	ALA	engineered mutation	UNP A0A059PIQ0
A	30	ARG	GLN	engineered mutation	UNP A0A059PIQ0
A	55	LYS	THR	engineered mutation	UNP A0A059PIQ0
A	61	VAL	GLU	engineered mutation	UNP A0A059PIQ0
A	78	THR	ILE	engineered mutation	UNP A0A059PIQ0
A	116	THR	LYS	engineered mutation	UNP A0A059PIQ0
A	117	VAL	ILE	engineered mutation	UNP A0A059PIQ0
A	155	THR	SER	engineered mutation	UNP A0A059PIQ0
A	156	VAL	ALA	engineered mutation	UNP A0A059PIQ0
A	188	GLY	-	linker	UNP A0A059PIQ0
A	189	GLY	-	linker	UNP A0A059PIQ0

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	190	THR	-	linker	UNP A0A059PIQ0
A	191	GLY	-	linker	UNP A0A059PIQ0
A	192	GLY	-	linker	UNP A0A059PIQ0
A	193	SER	-	linker	UNP A0A059PIQ0
A	194	ALA	-	linker	UNP A0A059PIQ0
A	195	SER	-	linker	UNP A0A059PIQ0
A	196	GLN	-	linker	UNP A0A059PIQ0
A	223	ARG	SER	engineered mutation	UNP P42212
A	232	ILE	TYR	engineered mutation	UNP P42212
A	241	SER	CYS	engineered mutation	UNP P42212
B	-10	MET	-	initiating methionine	UNP A0A059PIQ0
B	-9	GLY	-	expression tag	UNP A0A059PIQ0
B	-8	HIS	-	expression tag	UNP A0A059PIQ0
B	-7	HIS	-	expression tag	UNP A0A059PIQ0
B	-6	HIS	-	expression tag	UNP A0A059PIQ0
B	-5	HIS	-	expression tag	UNP A0A059PIQ0
B	-4	HIS	-	expression tag	UNP A0A059PIQ0
B	-3	HIS	-	expression tag	UNP A0A059PIQ0
B	-2	SER	-	expression tag	UNP A0A059PIQ0
B	-1	SER	-	expression tag	UNP A0A059PIQ0
B	0	GLY	-	expression tag	UNP A0A059PIQ0
B	15	QC4	THR	chromophore	UNP A0A059PIQ0
B	15	QC4	TYR	chromophore	UNP A0A059PIQ0
B	15	QC4	GLY	chromophore	UNP A0A059PIQ0
B	22	SER	ALA	engineered mutation	UNP A0A059PIQ0
B	30	ARG	GLN	engineered mutation	UNP A0A059PIQ0
B	55	LYS	THR	engineered mutation	UNP A0A059PIQ0
B	61	VAL	GLU	engineered mutation	UNP A0A059PIQ0
B	78	THR	ILE	engineered mutation	UNP A0A059PIQ0
B	116	THR	LYS	engineered mutation	UNP A0A059PIQ0
B	117	VAL	ILE	engineered mutation	UNP A0A059PIQ0
B	155	THR	SER	engineered mutation	UNP A0A059PIQ0
B	156	VAL	ALA	engineered mutation	UNP A0A059PIQ0
B	188	GLY	-	linker	UNP A0A059PIQ0
B	189	GLY	-	linker	UNP A0A059PIQ0
B	190	THR	-	linker	UNP A0A059PIQ0
B	191	GLY	-	linker	UNP A0A059PIQ0
B	192	GLY	-	linker	UNP A0A059PIQ0
B	193	SER	-	linker	UNP A0A059PIQ0
B	194	ALA	-	linker	UNP A0A059PIQ0
B	195	SER	-	linker	UNP A0A059PIQ0
B	196	GLN	-	linker	UNP A0A059PIQ0

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	223	ARG	SER	engineered mutation	UNP P42212
B	232	ILE	TYR	engineered mutation	UNP P42212
B	241	SER	CYS	engineered mutation	UNP P42212


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	192	Total 192	O 192	0	0
2	B	137	Total 137	O 137	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Green fluorescent protein, Green fluorescent protein



- Molecule 1: Green fluorescent protein, Green fluorescent protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.39Å 68.46Å 60.60Å 90.00° 102.07° 90.00°	Depositor
Resolution (Å)	38.93 – 1.36 38.93 – 1.36	Depositor EDS
% Data completeness (in resolution range)	97.9 (38.93-1.36) 91.8 (38.93-1.36)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.31 (at 1.36Å)	Xtriage
Refinement program	PHENIX 1.13RC2_2986	Depositor
R, $R_{free}$	0.174 , 0.205 0.176 , 0.208	Depositor DCC
$R_{free}$ test set	4078 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.8	Xtriage
Anisotropy	0.203	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 40.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4106	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.52% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: QC4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.31	0/1940	0.56	0/2629
1	B	0.30	0/1972	0.54	0/2668
All	All	0.30	0/3912	0.55	0/5297

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1874	0	1858	21	0
1	B	1903	0	1901	22	0
2	A	192	0	0	7	0
2	B	137	0	0	7	0
All	All	4106	0	3759	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:27:HIS:ND1	2:A:403:HOH:O	2.22	0.72
1:A:154:GLN:NE2	2:B:402:HOH:O	2.28	0.67
1:B:165[A]:ARG:NH2	1:B:240:ILE:HD13	2.14	0.63
1:B:165[A]:ARG:CZ	1:B:240:ILE:HG21	2.29	0.62
1:A:83:ASP:OD1	1:A:83:ASP:N	2.36	0.59
1:B:2:LYS:NZ	2:B:406:HOH:O	2.34	0.58
1:A:95:PHE:HB3	1:A:155[B]:THR:HG23	1.86	0.58
1:A:156[A]:VAL:HG11	1:B:173:PHE:CD2	2.40	0.56
1:B:65:GLU:CD	1:B:70[B]:VAL:HG21	2.26	0.56
1:B:70[B]:VAL:HG12	1:B:208:LEU:HB3	1.89	0.55
1:A:231[B]:THR:HG23	2:A:413:HOH:O	2.07	0.53
1:B:95:PHE:HB3	1:B:155[B]:THR:OG1	2.10	0.52
1:B:238:LYS:HE2	1:B:240:ILE:HD11	1.92	0.51
1:B:72:ARG:HG3	2:B:479:HOH:O	2.12	0.49
1:A:173:PHE:CD2	1:B:156[B]:VAL:HG11	2.47	0.49
1:A:171:LEU:HD11	1:A:173:PHE:HE2	1.77	0.49
1:B:65:GLU:OE1	1:B:70[B]:VAL:HG21	2.12	0.49
1:A:173:PHE:HD2	1:B:156[B]:VAL:HG11	1.78	0.48
1:A:82:GLU:CD	1:A:82:GLU:H	2.17	0.48
1:A:155[B]:THR:HG22	2:A:471:HOH:O	2.14	0.47
1:A:70[B]:VAL:HG12	1:A:208:LEU:HB3	1.96	0.47
1:A:171:LEU:HD11	1:A:173:PHE:CE2	2.51	0.46
1:A:236[A]:THR:HG23	2:A:429:HOH:O	2.14	0.46
1:A:103[B]:THR:HG23	2:A:515:HOH:O	2.15	0.45
1:A:70[B]:VAL:HG23	2:A:446:HOH:O	2.16	0.45
1:A:236[B]:THR:HG22	2:A:547:HOH:O	2.17	0.44
1:B:142:PRO:HA	2:B:466:HOH:O	2.16	0.44
1:B:81:LYS:HA	1:B:81:LYS:HD2	1.73	0.43
1:B:60:ALA:HA	1:B:72:ARG:O	2.19	0.43
1:B:193:SER:N	2:B:414:HOH:O	2.52	0.42
1:B:72:ARG:C	1:B:73:ILE:HG13	2.38	0.42
1:B:223:ARG:NE	2:B:403:HOH:O	2.29	0.42
1:A:171:LEU:HD21	1:A:234:LYS:HE3	2.01	0.42
1:B:162:ASN:ND2	2:B:401:HOH:O	2.21	0.42
1:A:65:GLU:OE2	1:A:72:ARG:NH1	2.36	0.42
1:A:13:THR:CG2	1:A:73:ILE:HG21	2.50	0.42
1:B:113:ALA:HB3	1:B:133:GLN:HB3	2.01	0.41
1:B:223:ARG:NH2	1:B:225[A]:GLU:OE2	2.53	0.41
1:A:173:PHE:HA	1:A:233:GLY:O	2.21	0.41
1:B:165[B]:ARG:HH22	1:B:242:THR:HG22	1.87	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	237/252 (94%)	235 (99%)	2 (1%)	0	100	100
1	B	241/252 (96%)	238 (99%)	3 (1%)	0	100	100
All	All	478/504 (95%)	473 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	214/215 (100%)	209 (98%)	5 (2%)	50	16
1	B	216/215 (100%)	211 (98%)	5 (2%)	50	16
All	All	430/430 (100%)	420 (98%)	10 (2%)	59	16

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3[A]	LEU
1	A	3[B]	LEU
1	A	106	LYS
1	A	241[A]	SER
1	A	241[B]	SER
1	B	81	LYS
1	B	83	ASP
1	B	108	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	236[A]	THR
1	B	236[B]	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	QC4	B	15	1	26,26,27	3.74	10 (38%)	36,38,40	1.65	10 (27%)
1	QC4	A	15	1	26,26,27	3.39	10 (38%)	36,38,40	1.84	8 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	QC4	B	15	1	-	0/12/31/32	0/2/2/2
1	QC4	A	15	1	-	0/12/31/32	0/2/2/2

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	15	QC4	CG2-CD2	10.97	1.46	1.39
1	B	15	QC4	CZ-CE1	9.15	1.50	1.39

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	15	QC4	CZ-CE1	9.04	1.49	1.39
1	A	15	QC4	CG2-CD2	8.48	1.44	1.39
1	B	15	QC4	CZ-CE2	6.53	1.49	1.38
1	A	15	QC4	CZ-CE2	6.24	1.48	1.38
1	B	15	QC4	CD1-CG2	5.56	1.50	1.40
1	A	15	QC4	CD1-CG2	5.37	1.50	1.40
1	B	15	QC4	OH-CZ	-4.80	1.25	1.37
1	A	15	QC4	OH-CZ	-4.62	1.26	1.37
1	B	15	QC4	CB2-CA2	4.52	1.38	1.35
1	B	15	QC4	C1-N3	3.69	1.43	1.37
1	A	15	QC4	CB2-CA2	3.56	1.38	1.35
1	A	15	QC4	C1-N3	3.39	1.42	1.37
1	A	15	QC4	CA1-C1	-2.95	1.47	1.51
1	B	15	QC4	CG2-CB2	-2.87	1.42	1.46
1	A	15	QC4	CG2-CB2	-2.65	1.42	1.46
1	B	15	QC4	CA1-C1	-2.24	1.48	1.51
1	A	15	QC4	CD1-CE1	2.14	1.41	1.37
1	B	15	QC4	CD1-CE1	2.01	1.41	1.37

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	15	QC4	O2-C2-CA2	5.05	133.79	130.96
1	B	15	QC4	C2-N3-C1	-4.73	105.57	107.97
1	A	15	QC4	F2-CD2-CG2	-3.74	116.28	119.68
1	A	15	QC4	C2-N3-C1	-3.51	106.19	107.97
1	A	15	QC4	N3-C1-N2	3.02	113.54	111.45
1	B	15	QC4	CA1-C1-N3	-2.99	121.16	124.75
1	B	15	QC4	N3-C1-N2	2.99	113.53	111.45
1	A	15	QC4	C1-CA1-N1	-2.86	105.32	109.96
1	A	15	QC4	CA1-C1-N3	-2.81	121.37	124.75
1	B	15	QC4	CD1-CE1-CZ	-2.77	121.62	123.79
1	B	15	QC4	C1-CA1-N1	-2.73	105.54	109.96
1	B	15	QC4	F2-CD2-CG2	-2.67	117.25	119.68
1	A	15	QC4	CD1-CE1-CZ	-2.47	121.86	123.79
1	B	15	QC4	CA3-N3-C2	2.32	129.12	123.80
1	B	15	QC4	F3-CE1-CZ	2.25	118.80	117.13
1	A	15	QC4	CG2-CD1-CE1	2.19	121.75	119.67
1	B	15	QC4	CA2-N2-C1	2.16	107.36	105.77
1	B	15	QC4	CG2-CD1-CE1	2.08	121.65	119.67

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	225/252 (89%)	0.27	11 (4%) 29 33	14, 25, 44, 84	0
1	B	227/252 (90%)	0.59	25 (11%) 5 6	18, 34, 63, 87	0
All	All	452/504 (89%)	0.43	36 (7%) 12 13	14, 29, 59, 87	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	179	ILE	4.1
1	A	194	ALA	4.0
1	B	141	GLY	3.9
1	A	195	SER	3.9
1	A	161	PRO	3.7
1	B	107	GLN	3.5
1	B	83	ASP	3.4
1	B	193	SER	3.4
1	A	193	SER	3.3
1	A	242	THR	3.3
1	B	243	THR	3.0
1	B	140	ASP	3.0
1	B	145	LEU	2.9
1	B	30[A]	ARG	2.8
1	B	33	PHE	2.8
1	A	83	ASP	2.7
1	A	179	ILE	2.7
1	B	134[A]	GLN	2.7
1	B	81	LYS	2.6
1	B	138	ILE	2.6
1	A	82	GLU	2.5
1	A	2	LYS	2.5
1	B	132	TYR	2.5
1	B	110	GLY	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	1	GLY	2.4
1	A	162	ASN	2.4
1	B	105	ASP	2.3
1	B	142	PRO	2.2
1	B	162	ASN	2.2
1	A	132	TYR	2.2
1	B	118[A]	ARG	2.2
1	B	108	LYS	2.2
1	B	143	VAL	2.2
1	B	194	ALA	2.2
1	B	26	ASP	2.1
1	B	25	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	QC4	B	15	25/26	0.94	0.08	19,23,29,34	0
1	QC4	A	15	25/26	0.96	0.08	13,17,24,33	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.