



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 5, 2020 – 11:16 AM EDT

PDB ID : 6UN7
Title : Crystal structure of green fluorescent protein (GFP); S65T, Y66(3-OMeY); ih
circular permutant (50-51)
Authors : Lin, C.-Y.; Boxer, S.G.
Deposited on : 2019-10-10
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.6
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.6


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	252	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>7%</div> <div>10%</div> </div> </div>
1	B	252	<div> <div>9%</div> <div> <div></div> <div>77%</div> <div>13%</div> <div>9%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4061 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	227	Total	C	N	O	S	0	17	0
			1876	1197	309	366	4			
1	B	229	Total	C	N	O	S	0	17	0
			1905	1217	318	365	5			

There are 70 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	MET	-	initiating methionine	UNP A0A059PIQ0
A	-9	GLY	-	expression tag	UNP A0A059PIQ0
A	-8	HIS	-	expression tag	UNP A0A059PIQ0
A	-7	HIS	-	expression tag	UNP A0A059PIQ0
A	-6	HIS	-	expression tag	UNP A0A059PIQ0
A	-5	HIS	-	expression tag	UNP A0A059PIQ0
A	-4	HIS	-	expression tag	UNP A0A059PIQ0
A	-3	HIS	-	expression tag	UNP A0A059PIQ0
A	-2	SER	-	expression tag	UNP A0A059PIQ0
A	-1	SER	-	expression tag	UNP A0A059PIQ0
A	0	GLY	-	expression tag	UNP A0A059PIQ0
A	15	QCD	THR	chromophore	UNP A0A059PIQ0
A	?	-	TYR	chromophore	UNP A0A059PIQ0
A	?	-	GLY	chromophore	UNP A0A059PIQ0
A	22	SER	ALA	engineered mutation	UNP A0A059PIQ0
A	30	ARG	GLN	engineered mutation	UNP A0A059PIQ0
A	55	LYS	THR	engineered mutation	UNP A0A059PIQ0
A	61	VAL	GLU	engineered mutation	UNP A0A059PIQ0
A	78	THR	ILE	engineered mutation	UNP A0A059PIQ0
A	116	THR	LYS	engineered mutation	UNP A0A059PIQ0
A	117	VAL	ILE	engineered mutation	UNP A0A059PIQ0
A	155	THR	SER	engineered mutation	UNP A0A059PIQ0
A	156	VAL	ALA	engineered mutation	UNP A0A059PIQ0
A	188	GLY	-	linker	UNP A0A059PIQ0
A	189	GLY	-	linker	UNP A0A059PIQ0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	190	THR	-	linker	UNP A0A059PIQ0
A	191	GLY	-	linker	UNP A0A059PIQ0
A	192	GLY	-	linker	UNP A0A059PIQ0
A	193	SER	-	linker	UNP A0A059PIQ0
A	194	ALA	-	linker	UNP A0A059PIQ0
A	195	SER	-	linker	UNP A0A059PIQ0
A	196	GLN	-	linker	UNP A0A059PIQ0
A	223	ARG	SER	conflict	UNP P42212
A	232	ILE	TYR	conflict	UNP P42212
A	241	SER	CYS	conflict	UNP P42212
B	-10	MET	-	initiating methionine	UNP A0A059PIQ0
B	-9	GLY	-	expression tag	UNP A0A059PIQ0
B	-8	HIS	-	expression tag	UNP A0A059PIQ0
B	-7	HIS	-	expression tag	UNP A0A059PIQ0
B	-6	HIS	-	expression tag	UNP A0A059PIQ0
B	-5	HIS	-	expression tag	UNP A0A059PIQ0
B	-4	HIS	-	expression tag	UNP A0A059PIQ0
B	-3	HIS	-	expression tag	UNP A0A059PIQ0
B	-2	SER	-	expression tag	UNP A0A059PIQ0
B	-1	SER	-	expression tag	UNP A0A059PIQ0
B	0	GLY	-	expression tag	UNP A0A059PIQ0
B	15	QCD	THR	chromophore	UNP A0A059PIQ0
B	?	-	TYR	chromophore	UNP A0A059PIQ0
B	?	-	GLY	chromophore	UNP A0A059PIQ0
B	22	SER	ALA	engineered mutation	UNP A0A059PIQ0
B	30	ARG	GLN	engineered mutation	UNP A0A059PIQ0
B	55	LYS	THR	engineered mutation	UNP A0A059PIQ0
B	61	VAL	GLU	engineered mutation	UNP A0A059PIQ0
B	78	THR	ILE	engineered mutation	UNP A0A059PIQ0
B	116	THR	LYS	engineered mutation	UNP A0A059PIQ0
B	117	VAL	ILE	engineered mutation	UNP A0A059PIQ0
B	155	THR	SER	engineered mutation	UNP A0A059PIQ0
B	156	VAL	ALA	engineered mutation	UNP A0A059PIQ0
B	188	GLY	-	linker	UNP A0A059PIQ0
B	189	GLY	-	linker	UNP A0A059PIQ0
B	190	THR	-	linker	UNP A0A059PIQ0
B	191	GLY	-	linker	UNP A0A059PIQ0
B	192	GLY	-	linker	UNP A0A059PIQ0
B	193	SER	-	linker	UNP A0A059PIQ0
B	194	ALA	-	linker	UNP A0A059PIQ0
B	195	SER	-	linker	UNP A0A059PIQ0
B	196	GLN	-	linker	UNP A0A059PIQ0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	223	ARG	SER	engineered mutation	UNP P42212
B	232	ILE	TYR	engineered mutation	UNP P42212
B	241	SER	CYS	engineered mutation	UNP P42212


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	169	Total 169	O 169	0	0
2	B	111	Total 111	O 111	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Green fluorescent protein



- Molecule 1: Green fluorescent protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.88Å 68.60Å 60.19Å 90.00° 101.64° 90.00°	Depositor
Resolution (Å)	36.14 – 1.50 36.14 – 1.50	Depositor EDS
% Data completeness (in resolution range)	97.7 (36.14-1.50) 88.9 (36.14-1.50)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.89 (at 1.50Å)	Xtriage
Refinement program	PHENIX 1.13RC2_2986	Depositor
R, R_{free}	0.167 , 0.191 0.170 , 0.191	Depositor DCC
R_{free} test set	3173 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	19.8	Xtriage
Anisotropy	0.442	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 45.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4061	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: QCD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/1946	0.62	0/2637
1	B	0.35	0/1974	0.57	0/2667
All	All	0.36	0/3920	0.59	0/5304

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1876	0	1861	17	0
1	B	1905	0	1893	32	0
2	A	169	0	0	3	0
2	B	111	0	0	4	0
All	All	4061	0	3754	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:40:GLU:OE2	1:B:139:GLY:HA3	1.82	0.80
1:A:156[B]:VAL:HG23	1:B:156:VAL:HG13	1.70	0.73
1:A:238:LYS:HE2	1:A:240:ILE:HD11	1.70	0.73
1:B:171:LEU:HD23	1:B:236:THR:HG22	1.69	0.73
1:A:76:LYS:HE3	2:A:433:HOH:O	1.90	0.71
1:B:172:GLU:HG2	1:B:235[B]:LEU:HB2	1.72	0.70
1:B:63:LYS:HE2	1:B:65:GLU:HG3	1.79	0.65
1:A:172:GLU:HG2	1:A:235:LEU:HB2	1.79	0.64
1:B:47:THR:HG23	1:B:57:LYS:HE2	1.88	0.56
1:A:156[A]:VAL:HG13	1:B:156:VAL:HG13	1.86	0.56
1:A:156[B]:VAL:HG11	1:B:173:PHE:HD2	1.72	0.54
1:B:214:ASP:OD1	2:B:402:HOH:O	2.19	0.54
1:B:134[A]:GLN:NE2	2:B:408:HOH:O	2.42	0.52
1:A:156[A]:VAL:HG12	1:B:154:GLN:HB3	1.92	0.51
1:B:210[B]:GLU:OE2	1:B:223:ARG:NH1	2.44	0.50
1:A:70[B]:VAL:HG12	1:A:208:LEU:HB3	1.96	0.48
1:B:59:ARG:NH2	2:B:405:HOH:O	2.24	0.48
1:A:232:ILE:HG22	1:B:161:PRO:HG3	1.96	0.47
1:B:59:ARG:HB3	1:B:74[B]:GLU:CG	2.45	0.47
1:A:103[B]:THR:HG23	1:A:148:ASN:OD1	2.15	0.47
1:A:154:GLN:HB3	1:B:156:VAL:HG12	1.97	0.46
1:B:226:GLY:HA3	1:B:237:LEU:HD23	1.98	0.46
1:B:202:THR:O	2:B:403:HOH:O	2.21	0.46
1:B:221[A]:SER:HB2	1:B:242:THR:HG1	1.81	0.45
1:B:47:THR:OG1	1:B:57:LYS:HE2	2.16	0.45
1:B:171:LEU:HD23	1:B:236:THR:CG2	2.41	0.45
1:A:74[B]:GLU:OE1	2:A:401:HOH:O	2.21	0.45
1:A:156[A]:VAL:HG21	1:B:171:LEU:HD12	1.99	0.45
1:B:73:ILE:HB	1:B:211[B]:LEU:HD13	1.99	0.44
1:B:111:ILE:HG13	1:B:135:ASN:HB2	2.00	0.43
1:A:171:LEU:HD23	1:B:171:LEU:HD12	2.00	0.43
1:B:81:LYS:HD3	1:B:81:LYS:HA	1.75	0.43
1:B:47:THR:CG2	1:B:57:LYS:HE2	2.50	0.42
1:A:83:ASP:OD1	1:A:90:LYS:HE3	2.20	0.41
1:B:171:LEU:HD22	1:B:172:GLU:N	2.35	0.41
1:B:18:VAL:HG23	1:B:235[B]:LEU:HD11	2.01	0.41
1:A:156[A]:VAL:HG21	1:B:171:LEU:CD1	2.51	0.40
1:B:71:ASN:ND2	1:B:73:ILE:HD11	2.37	0.40
1:A:70[B]:VAL:HG23	2:A:424:HOH:O	2.22	0.40
1:B:6:PRO:HD3	1:B:86:ILE:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	239/252 (95%)	236 (99%)	3 (1%)	0	100	100
1	B	240/252 (95%)	235 (98%)	5 (2%)	0	100	100
All	All	479/504 (95%)	471 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	215/215 (100%)	213 (99%)	2 (1%)	78	61
1	B	216/215 (100%)	215 (100%)	1 (0%)	88	78
All	All	431/430 (100%)	428 (99%)	3 (1%)	81	69

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	76	LYS
1	A	107	GLN
1	B	59	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	QCD	B	15	1	25,25,26	2.58	6 (24%)	33,35,37	2.52	11 (33%)
1	QCD	A	15	1	25,25,26	2.47	5 (20%)	33,35,37	2.34	10 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	QCD	B	15	1	-	1/14/33/34	0/2/2/2
1	QCD	A	15	1	-	0/14/33/34	0/2/2/2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	15	QCD	CB2-CA2	8.79	1.42	1.35
1	B	15	QCD	CB2-CA2	8.63	1.42	1.35
1	B	15	QCD	O4-CE1	5.98	1.46	1.37
1	A	15	QCD	O4-CE1	5.55	1.46	1.37
1	B	15	QCD	CA1-C1	-4.55	1.44	1.51
1	A	15	QCD	CA1-C1	-3.58	1.46	1.51
1	B	15	QCD	CB1-CA1	-3.41	1.40	1.53
1	A	15	QCD	CB1-CA1	-3.15	1.41	1.53
1	B	15	QCD	CA1-N1	-2.55	1.40	1.47
1	B	15	QCD	OG1-CB1	-2.47	1.36	1.43
1	A	15	QCD	CA1-N1	-2.20	1.41	1.47

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	15	QCD	O2-C2-CA2	-9.15	125.82	130.96
1	A	15	QCD	CA2-C2-N3	5.68	106.06	103.37
1	A	15	QCD	O2-C2-CA2	-5.57	127.83	130.96
1	B	15	QCD	CA2-C2-N3	5.10	105.78	103.37
1	A	15	QCD	CA2-N2-C1	5.01	109.46	105.77
1	A	15	QCD	C2-CA2-N2	-4.72	105.63	108.93
1	B	15	QCD	C4-O4-CE1	-4.42	110.86	117.53
1	A	15	QCD	C4-O4-CE1	-3.75	111.87	117.53
1	B	15	QCD	CA2-N2-C1	3.64	108.46	105.77
1	B	15	QCD	C2-CA2-N2	-3.55	106.45	108.93
1	A	15	QCD	OG1-CB1-CA1	2.87	115.19	109.04
1	B	15	QCD	O3-C3-CA3	-2.51	118.80	126.39
1	A	15	QCD	O3-C3-CA3	-2.40	119.15	126.39
1	B	15	QCD	C1-CA1-N1	-2.39	106.08	109.96
1	A	15	QCD	C1-CA1-N1	-2.13	106.50	109.96
1	B	15	QCD	CG2-CB2-CA2	-2.12	127.34	129.94
1	A	15	QCD	CD2-CE2-CZ	-2.10	118.34	120.50
1	A	15	QCD	N3-C1-N2	-2.10	110.00	111.45
1	B	15	QCD	OG1-CB1-CA1	2.07	113.47	109.04
1	B	15	QCD	CA1-C1-N3	-2.07	122.27	124.75
1	B	15	QCD	O2-C2-N3	2.04	128.40	124.35

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	15	QCD	C3-CA3-N3-C2

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	226/252 (89%)	-0.11	5 (2%) 62 67	19, 28, 52, 80	0
1	B	228/252 (90%)	0.42	22 (9%) 8 8	23, 41, 75, 98	0
All	All	454/504 (90%)	0.16	27 (5%) 22 24	19, 34, 64, 98	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	145	LEU	5.1
1	B	180	THR	4.3
1	B	182	GLY	4.2
1	B	196	GLN	3.8
1	B	1	GLY	3.8
1	B	181[A]	HIS	3.5
1	B	33	PHE	3.5
1	A	242	THR	3.3
1	B	30	ARG	3.3
1	B	143	VAL	3.2
1	B	109	ASN	3.0
1	B	37	ALA	2.8
1	B	67	ASP	2.6
1	B	26	ASP	2.5
1	B	136	THR	2.5
1	B	146	PRO	2.5
1	B	144	LEU	2.4
1	A	192	GLY	2.4
1	B	138	ILE	2.3
1	B	243	THR	2.2
1	A	107	GLN	2.2
1	B	29	LYS	2.2
1	B	134[A]	GLN	2.2
1	A	30	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	156	VAL	2.1
1	B	199	GLU	2.1
1	A	117	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	QCD	B	15	24/25	0.97	0.07	24,27,30,32	0
1	QCD	A	15	24/25	0.98	0.07	18,23,25,30	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.