



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 03:59 pm BST

PDB ID : 3UO3
Title : Jac1 co-chaperone from *Saccharomyces cerevisiae*, 5-182 clone
Authors : Osipiuk, J.; Bigelow, L.; Mulligan, R.; Feldmann, B.; Babnigg, G.; Marszalek, J.; Craig, E.A.; Dutkiewicz, R.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2011-11-16
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	181	<div> <div>22%</div> <div>86%</div> <div>9%</div> <div>..</div> </div>
1	B	181	<div> <div>22%</div> <div>57%</div> <div>22%</div> <div>•</div> <div>19%</div> </div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2907 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called J-type co-chaperone JAC1, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	174	Total	C	N	O	S	0	10	0
			1496	950	258	284	4			
1	B	147	Total	C	N	O	S	0	3	0
			1244	797	210	234	3			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	SER	-	EXPRESSION TAG	UNP P53193
A	3	ASN	-	EXPRESSION TAG	UNP P53193
A	4	ALA	-	EXPRESSION TAG	UNP P53193
B	2	SER	-	EXPRESSION TAG	UNP P53193
B	3	ASN	-	EXPRESSION TAG	UNP P53193
B	4	ALA	-	EXPRESSION TAG	UNP P53193

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	154	Total	O	0	0
			154	154		
3	B	9	Total	O	0	0
			9	9		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: J-type co-chaperone JAC1, mitochondrial

Chain A: 



- Molecule 1: J-type co-chaperone JAC1, mitochondrial

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.79Å 60.45Å 108.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.50 – 1.85 40.45 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.8 (40.50-1.85) 99.8 (40.45-1.85)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.63 (at 1.86Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.192 , 0.238 0.192 , 0.236	Depositor DCC
R_{free} test set	1736 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	33.3	Xtriage
Anisotropy	0.637	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 48.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.018 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2907	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.07	5/1555 (0.3%)	0.96	3/2097 (0.1%)
1	B	0.63	2/1276 (0.2%)	0.73	0/1720
All	All	0.90	7/2831 (0.2%)	0.86	3/3817 (0.1%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	174	TRP	CD2-CE2	6.33	1.49	1.41
1	A	139	GLU	CD-OE2	-5.91	1.19	1.25
1	A	29	TRP	CD2-CE2	5.67	1.48	1.41
1	B	164	TRP	CD2-CE2	5.49	1.48	1.41
1	A	151	TYR	CE1-CZ	5.45	1.45	1.38
1	A	165	TYR	CE1-CZ	5.34	1.45	1.38
1	B	29	TRP	CD2-CE2	5.25	1.47	1.41

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	106	LEU	CA-CB-CG	-6.01	101.47	115.30
1	A	109	LEU	CA-CB-CG	5.68	128.37	115.30
1	A	165	TYR	CD1-CE1-CZ	-5.49	114.86	119.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1496	0	1518	9	0
1	B	1244	0	1259	25	0
2	A	4	0	3	1	0
3	A	154	0	0	2	0
3	B	9	0	0	0	0
All	All	2907	0	2780	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:25[B]:LYS:HD2	3:A:188:HOH:O	1.44	1.15
1:B:86:ILE:HG23	1:B:93:THR:HG21	1.38	1.03
1:B:86:ILE:CG2	1:B:93:THR:HG21	2.08	0.82
1:B:18:PHE:O	1:B:21:THR:O	2.04	0.74
1:B:40:TYR:O	1:B:44:GLN:HG3	1.90	0.71
1:A:171:PHE:O	2:A:501:ACT:H2	1.91	0.69
1:B:13:THR:O	1:B:17:LEU:HD22	1.95	0.65
1:B:21:THR:HG22	1:B:39:GLU:OE1	1.98	0.63
1:B:21:THR:HG21	1:B:36:LEU:HA	1.81	0.62
1:B:99:THR:O	1:B:101:ASP:N	2.34	0.60
1:B:79:MET:O	1:B:81:LYS:N	2.32	0.59
1:B:43:LEU:HD23	1:B:62:LEU:HD11	1.84	0.58
1:B:41:ARG:HA	1:B:44:GLN:OE1	2.05	0.57
1:B:121:GLU:O	1:B:125:LYS:HG2	2.05	0.56
1:A:109:LEU:HD13	1:B:109:LEU:CD2	2.37	0.55
1:B:21:THR:CG2	1:B:39:GLU:OE1	2.55	0.54
1:A:109:LEU:HD13	1:B:109:LEU:HG	1.92	0.52
1:B:24:LYS:HE3	1:B:26:LEU:HB2	1.93	0.51
1:B:29:TRP:CD1	1:B:143:GLY:HA2	2.49	0.48
1:A:20:LYS:HD3	1:A:35[B]:ARG:NH1	2.30	0.47
1:B:80:LEU:HD22	1:B:146:TYR:OH	2.15	0.47
1:B:20[B]:LYS:HD3	1:B:39:GLU:OE2	2.16	0.45
1:B:131:ASN:OD1	1:B:134:ARG:NH2	2.50	0.44
1:B:104:LEU:O	1:B:108:VAL:HG23	2.18	0.43
1:B:24:LYS:HB3	1:B:24:LYS:HE2	1.74	0.43
1:A:42:GLN:NE2	3:A:196:HOH:O	2.51	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:181:GLU:OE2	1:B:172:LYS:NZ	2.45	0.42
1:A:132:LYS:O	1:A:136:GLN:HG3	2.19	0.42
1:A:124:VAL:HG21	1:A:174:TRP:CE2	2.56	0.41
1:B:44:GLN:C	1:B:46:GLN:H	2.25	0.40
1:B:35:ARG:O	1:B:35:ARG:HG2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	182/181 (101%)	182 (100%)	0	0	100	100
1	B	144/181 (80%)	137 (95%)	6 (4%)	1 (1%)	22	9
All	All	326/362 (90%)	319 (98%)	6 (2%)	1 (0%)	41	26

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	100	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/165 (102%)	164 (97%)	5 (3%)	41	24
1	B	138/165 (84%)	126 (91%)	12 (9%)	10	2
All	All	307/330 (93%)	290 (94%)	17 (6%)	20	7

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	85	ASN
1	A	100	SER
1	A	106	LEU
1	A	109	LEU
1	A	149	LYS
1	B	17	LEU
1	B	21	THR
1	B	22	PHE
1	B	44	GLN
1	B	80	LEU
1	B	89	THR
1	B	91	GLU
1	B	103	GLN
1	B	105	LEU
1	B	106	LEU
1	B	136	GLN
1	B	144	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	42	GLN
1	B	130	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACT	A	501	-	1,3,3	2.37	1 (100%)	0,3,3	0.00	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	ACT	CH3-C	2.37	1.51	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	174/181 (96%)	0.20	0 100 100	24, 37, 68, 98	0
1	B	147/181 (81%)	1.24	39 (26%) 0 0	51, 76, 112, 135	0
All	All	321/362 (88%)	0.67	39 (12%) 4 4	24, 55, 104, 135	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	94	SER	9.9
1	B	82	LEU	7.0
1	B	152	ALA	5.9
1	B	93	THR	5.5
1	B	151	TYR	4.5
1	B	30	THR	4.5
1	B	62	LEU	4.1
1	B	84	ARG	3.8
1	B	46	GLN	3.8
1	B	176	PRO	3.8
1	B	26	LEU	3.7
1	B	86	ILE	3.5
1	B	12	SER	3.2
1	B	69	LEU	3.1
1	B	150	ASP	3.1
1	B	83	LEU	3.1
1	B	88	LEU	3.0
1	B	153	ALA	3.0
1	B	23	PRO	2.9
1	B	24	LYS	2.9
1	B	40	TYR	2.8
1	B	92	GLN	2.6
1	B	45	ALA	2.5
1	B	100	SER	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	44	GLN	2.5
1	B	138	ILE	2.5
1	B	85	ASN	2.4
1	B	175	ALA	2.4
1	B	81	LYS	2.4
1	B	144	GLN	2.4
1	B	25	LYS	2.3
1	B	155	VAL	2.3
1	B	80	LEU	2.3
1	B	149	LYS	2.3
1	B	156	LYS	2.3
1	B	22	PHE	2.2
1	B	154	ALA	2.2
1	B	36	LEU	2.1
1	B	42	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ACT	A	501	4/4	0.93	0.08	44,59,63,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.