



wwPDB X-ray Structure Validation Summary Report ⓘ

May 17, 2020 – 12:15 pm BST

PDB ID : 3UO9
Title : Crystal Structure of Human GAC in Complex with Glutamate and BPTES
Authors : DeLaBarre, B.; Gross, S.; Cheng, F.; Gao, Y.; Jha, A.; Jiang, F.; Song, J.J.; Wei, W.; Hurov, J.B.
Deposited on : 2011-11-16
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	534	<div> <div>9%</div> <div>68% 8% • 23%</div> </div>
1	B	534	<div> <div>8%</div> <div>67% 6% • 25%</div> </div>
1	C	534	<div> <div>9%</div> <div>67% 7% •• 24%</div> </div>
1	D	534	<div> <div>6%</div> <div>67% 8% • 24%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	2	-	-	X	-
2	GOL	D	1	-	-	X	-
3	04A	B	2	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13319 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutaminase kidney isoform, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	411	Total	C	N	O	S	0	0	0
			3208	2044	541	595	28			
1	B	401	Total	C	N	O	S	0	0	0
			3134	2000	527	579	28			
1	C	404	Total	C	N	O	S	0	0	0
			3161	2019	531	583	28			
1	D	408	Total	C	N	O	S	0	0	0
			3191	2034	538	591	28			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	640	HIS	-	EXPRESSION TAG	UNP O94925
A	641	HIS	-	EXPRESSION TAG	UNP O94925
A	642	HIS	-	EXPRESSION TAG	UNP O94925
A	643	HIS	-	EXPRESSION TAG	UNP O94925
A	644	HIS	-	EXPRESSION TAG	UNP O94925
A	645	HIS	-	EXPRESSION TAG	UNP O94925
B	640	HIS	-	EXPRESSION TAG	UNP O94925
B	641	HIS	-	EXPRESSION TAG	UNP O94925
B	642	HIS	-	EXPRESSION TAG	UNP O94925
B	643	HIS	-	EXPRESSION TAG	UNP O94925
B	644	HIS	-	EXPRESSION TAG	UNP O94925
B	645	HIS	-	EXPRESSION TAG	UNP O94925
C	640	HIS	-	EXPRESSION TAG	UNP O94925
C	641	HIS	-	EXPRESSION TAG	UNP O94925
C	642	HIS	-	EXPRESSION TAG	UNP O94925
C	643	HIS	-	EXPRESSION TAG	UNP O94925
C	644	HIS	-	EXPRESSION TAG	UNP O94925
C	645	HIS	-	EXPRESSION TAG	UNP O94925
D	640	HIS	-	EXPRESSION TAG	UNP O94925
D	641	HIS	-	EXPRESSION TAG	UNP O94925
D	642	HIS	-	EXPRESSION TAG	UNP O94925

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	643	HIS	-	EXPRESSION TAG	UNP O94925
D	644	HIS	-	EXPRESSION TAG	UNP O94925
D	645	HIS	-	EXPRESSION TAG	UNP O94925

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is N,N'-[sulfanediylbis(ethane-2,1-diyl-1,3,4-thiadiazole-5,2-diyl)]bis(2-phenylacetamide) (three-letter code: 04A) (formula: $C_{24}H_{24}N_6O_2S_3$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	115	Total O 115 115	0	0
4	B	132	Total O 132 132	0	0
4	C	145	Total O 145 145	0	0
4	D	139	Total O 139 139	0	0

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Chain A:
-
- 9% 68% 8% 23%
- ILE LYS GLN GLY LEU LEU P137 S138 L139 E140 D141 L142 F144 Y145 T146 I147 A148 E149 Q150 Q151 E152 K153 I154 H157 L175 K176 E177 R183 L184 T185 L186 T189 S190 D191 G192 V193 M194 L195 D196 K197 D198 L199 F200 K201 K202 Q205 F215 R216 R217 Q241 E247
- D248 V265 R272 R311 L316 R317 F318 R319 R320 L321 F322 R325 D326 Q360 F361 D366 R387 M407 C424 A432 T433 L434 F441 R446 V453 L459 C463 Y466 G477 A480 T488 V491 V492 M496 G497 M498 M499 C500
- VAL TYR ARG MET GLU SER F527 H528 N529 H532 L533 R534 H535 F536 A537 K538 G546 ASP GLN ARG HIS SER PHE GLU TYR LEU ASP TYR SER LEU GLN GLU LEU ALA LYS GLU THR VAL TRP LYS VAL SER PRO GLU ASN GLU ASP ILE SER THR THR VAL

- [illegible]

- Molecule 1: Glutaminase kidney isoform, mitochondrial



- Molecule 1: Glutaminase kidney isoform, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.12Å 139.43Å 177.71Å 90.00° 93.73° 90.00°	Depositor
Resolution (Å)	45.09 – 2.30 46.21 – 2.28	Depositor EDS
% Data completeness (in resolution range)	99.0 (45.09-2.30) 99.0 (46.21-2.28)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.70 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.195 , 0.249 0.222 , 0.257	Depositor DCC
R_{free} test set	5439 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	32.4	Xtriage
Anisotropy	0.083	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 36.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	13319	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, 04A

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.96	3/3279 (0.1%)	1.06	18/4424 (0.4%)
1	B	0.93	0/3202	1.03	17/4317 (0.4%)
1	C	0.96	0/3230	1.08	20/4356 (0.5%)
1	D	0.96	1/3261 (0.0%)	1.08	22/4400 (0.5%)
All	All	0.95	4/12972 (0.0%)	1.06	77/17497 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6
1	B	0	3
1	C	0	5
1	D	0	6
All	All	0	20

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	463	CYS	CB-SG	-6.58	1.71	1.82
1	A	424	CYS	CB-SG	-6.04	1.72	1.82
1	A	500	CYS	CB-SG	-5.57	1.72	1.81
1	D	500	CYS	CB-SG	-5.51	1.72	1.81

The worst 5 of 77 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	543	ARG	NE-CZ-NH2	-13.25	113.68	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	454	ARG	NE-CZ-NH2	-9.34	115.63	120.30
1	C	534	ARG	NE-CZ-NH1	8.95	124.78	120.30
1	B	407	MET	CG-SD-CE	-8.81	86.10	100.20
1	A	216	ARG	NE-CZ-NH1	8.27	124.43	120.30

There are no chirality outliers.

5 of 20 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	183	ARG	Sidechain
1	A	317	ARG	Sidechain
1	A	387	ARG	Sidechain
1	A	446	ARG	Sidechain
1	A	534	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3208	0	3181	14	0
1	B	3134	0	3109	25	0
1	C	3161	0	3136	27	0
1	D	3191	0	3164	29	0
2	A	12	0	16	12	0
2	C	6	0	8	0	0
2	D	6	0	7	16	0
3	B	35	0	22	10	0
3	C	35	0	24	6	0
4	A	115	0	0	2	0
4	B	132	0	0	0	0
4	C	145	0	0	2	0
4	D	139	0	0	3	0
All	All	13319	0	12667	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 94 close contacts within the same asymmetric unit are listed below, sorted by their

clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:2:04A:CAE	1:D:317:ARG:HH12	1.19	1.24
3:B:2:04A:HAE	1:D:317:ARG:NH1	1.14	1.10
2:A:2:GOL:H32	1:C:532:ASN:HB2	1.28	1.07
3:B:2:04A:CAE	1:D:317:ARG:NH1	1.80	1.02
1:D:532:ASN:HD22	1:D:535:HIS:H	1.15	0.95

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	408/534 (76%)	398 (98%)	7 (2%)	3 (1%)	22	26
1	B	394/534 (74%)	379 (96%)	12 (3%)	3 (1%)	19	23
1	C	397/534 (74%)	385 (97%)	10 (2%)	2 (0%)	29	35
1	D	405/534 (76%)	390 (96%)	10 (2%)	5 (1%)	13	14
All	All	1604/2136 (75%)	1552 (97%)	39 (2%)	13 (1%)	19	23

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	217	ARG
1	B	527	PHE
1	B	538	LYS
1	C	526	ASN
1	D	217	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	356/458 (78%)	342 (96%)	14 (4%)	32	46
1	B	348/458 (76%)	337 (97%)	11 (3%)	39	54
1	C	351/458 (77%)	334 (95%)	17 (5%)	25	36
1	D	354/458 (77%)	338 (96%)	16 (4%)	27	39
All	All	1409/1832 (77%)	1351 (96%)	58 (4%)	30	43

5 of 58 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	143	LEU
1	C	250	ILE
1	D	318	PHE
1	C	153	LYS
1	C	195	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 38 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	157	HIS
1	C	461	HIS
1	D	519	HIS
1	C	319	ASN
1	C	475	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GOL	D	1	-	5,5,5	1.30	1 (20%)	5,5,5	1.43	1 (20%)
3	04A	C	1	-	32,38,38	2.78	7 (21%)	29,49,49	1.33	4 (13%)
2	GOL	A	4	-	5,5,5	0.38	0	5,5,5	0.67	0
2	GOL	A	2	-	5,5,5	0.80	0	5,5,5	1.35	0
2	GOL	C	3	-	5,5,5	0.42	0	5,5,5	0.71	0
3	04A	B	2	-	32,38,38	2.65	7 (21%)	29,49,49	1.08	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	D	1	-	-	2/4/4/4	-
3	04A	C	1	-	-	2/18/24/24	0/4/4/4
2	GOL	A	4	-	-	1/4/4/4	-
2	GOL	A	2	-	-	4/4/4/4	-
2	GOL	C	3	-	-	2/4/4/4	-
3	04A	B	2	-	-	4/18/24/24	0/4/4/4

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1	04A	CAQ-CBD	-9.12	1.36	1.51
3	C	1	04A	CAP-CBG	7.91	1.54	1.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	2	04A	CAQ-CBD	-7.82	1.38	1.51
3	B	2	04A	CAP-CBG	7.08	1.53	1.49
3	C	1	04A	CAO-CBF	6.46	1.53	1.49

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1	04A	CBI-NAX-CBC	-3.44	120.26	129.54
3	C	1	04A	CBH-NAW-CBB	-2.77	122.07	129.54
3	B	2	04A	CBH-NAW-CBB	-2.75	122.12	129.54
3	C	1	04A	CAN-SAY-CAM	-2.51	94.25	101.87
3	B	2	04A	CBI-NAX-CBC	-2.48	122.84	129.54

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	2	GOL	O1-C1-C2-C3
2	C	3	GOL	C1-C2-C3-O3
3	B	2	04A	SAY-CAM-CAO-CBF
2	A	2	GOL	O1-C1-C2-O2
2	D	1	GOL	O1-C1-C2-C3


There are no ring outliers.


5 monomers are involved in 44 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1	GOL	16	0
3	C	1	04A	6	0
2	A	4	GOL	1	0
2	A	2	GOL	11	0
3	B	2	04A	10	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	411/534 (76%)	0.69	49 (11%) 4 6	17, 28, 70, 82	0
1	B	401/534 (75%)	0.60	41 (10%) 6 9	17, 27, 61, 81	0
1	C	404/534 (75%)	0.67	49 (12%) 4 6	16, 26, 65, 81	0
1	D	408/534 (76%)	0.59	33 (8%) 12 16	15, 26, 66, 78	0
All	All	1624/2136 (76%)	0.64	172 (10%) 6 8	15, 27, 66, 82	0

The worst 5 of 172 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	318	PHE	8.6
1	C	186	LEU	8.4
1	A	317	ARG	8.4
1	A	546	GLY	7.4
1	B	186	LEU	6.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	C	3	6/6	0.39	0.37	54,61,61,62	0
3	04A	B	2	35/35	0.78	0.26	56,69,83,84	0
2	GOL	A	4	6/6	0.81	0.16	59,60,61,64	0
2	GOL	D	1	6/6	0.88	0.24	30,34,35,37	0
3	04A	C	1	35/35	0.89	0.21	49,62,90,91	0
2	GOL	A	2	6/6	0.89	0.28	28,33,37,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.