



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 08:35 am BST

PDB ID : 5UPK
Title : CDC42 binds PAK4 via an extended GTPase-effector interface - 3 peptide:
PAK4cat, PAK4-N45, CDC42
Authors : Ha, B.H.; Boggon, T.J.
Deposited on : 2017-02-03
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	48	<div> <div>4%</div> <div> <div></div> <div>77%</div> <div>6%</div> <div>17%</div> </div> </div>
2	B	346	<div> <div>%</div> <div> <div></div> <div>80%</div> <div>••</div> <div>16%</div> </div> </div>
3	C	185	<div> <div>18%</div> <div> <div></div> <div>87%</div> <div>8%</div> <div>5%</div> </div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4070 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase PAK 4.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	40	Total	C	N	O	0	0	0
			341	215	68	58			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP O96013
A	-1	GLY	-	expression tag	UNP O96013
A	0	SER	-	expression tag	UNP O96013

- Molecule 2 is a protein called Serine/threonine-protein kinase PAK 4.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	289	Total	C	N	O	P	S	0	0	0
			2284	1456	401	412	1	14			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	246	MET	-	initiating methionine	UNP O96013
B	247	GLY	-	expression tag	UNP O96013
B	248	SER	-	expression tag	UNP O96013
B	249	SER	-	expression tag	UNP O96013
B	250	HIS	-	expression tag	UNP O96013
B	251	HIS	-	expression tag	UNP O96013
B	252	HIS	-	expression tag	UNP O96013
B	253	HIS	-	expression tag	UNP O96013
B	254	HIS	-	expression tag	UNP O96013
B	255	HIS	-	expression tag	UNP O96013
B	256	SER	-	expression tag	UNP O96013
B	257	SER	-	expression tag	UNP O96013
B	258	GLY	-	expression tag	UNP O96013

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	259	LEU	-	expression tag	UNP O96013
B	260	VAL	-	expression tag	UNP O96013
B	261	PRO	-	expression tag	UNP O96013
B	262	ARG	-	expression tag	UNP O96013
B	263	GLY	-	expression tag	UNP O96013
B	264	SER	-	expression tag	UNP O96013
B	265	HIS	-	expression tag	UNP O96013
B	266	MET	-	expression tag	UNP O96013
B	267	GLU	-	expression tag	UNP O96013
B	268	ASN	-	expression tag	UNP O96013
B	269	LEU	-	expression tag	UNP O96013
B	270	TYR	-	expression tag	UNP O96013
B	271	PHE	-	expression tag	UNP O96013
B	272	GLN	-	expression tag	UNP O96013
B	273	GLY	-	expression tag	UNP O96013

- Molecule 3 is a protein called Cell division control protein 42 homolog.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	175	Total	C	N	O	S	0	0	0
			1364	879	217	262	6			

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	61	LEU	GLN	engineered mutation	UNP P60953
C	163	ARG	LYS	conflict	UNP P60953
C	178	LEU	-	expression tag	UNP P60953
C	179	GLU	-	expression tag	UNP P60953
C	180	HIS	-	expression tag	UNP P60953
C	181	HIS	-	expression tag	UNP P60953
C	182	HIS	-	expression tag	UNP P60953
C	183	HIS	-	expression tag	UNP P60953
C	184	HIS	-	expression tag	UNP P60953
C	185	HIS	-	expression tag	UNP P60953

- Molecule 4 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

- Molecule 5 is PHOSPHOAMINOPHOSPHONIC ACID-GUANYLATE ESTER (three-letter code: GNP) (formula: $\text{C}_{10}\text{H}_{17}\text{N}_6\text{O}_{13}\text{P}_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	C	1	Total	C	N	O	P	0	0
			32	10	6	13	3		

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	1	Total	Mg	0	0
			1	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	23	Total	O	0	0
			23	23		
7	C	2	Total	O	0	0
			2	2		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Serine/threonine-protein kinase PAK 4



- Molecule 2: Serine/threonine-protein kinase PAK 4



- Molecule 3: Cell division control protein 42 homolog



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.47Å 61.62Å 85.79Å 90.00° 91.57° 90.00°	Depositor
Resolution (Å)	50.01 – 2.40 42.88 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (50.01-2.40) 99.8 (42.88-2.40)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	0.16	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.52 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.8.0151	Depositor
R, R_{free}	0.218 , 0.264 0.218 , 0.261	Depositor DCC
R_{free} test set	1097 reflections (4.56%)	wwPDB-VP
Wilson B-factor (Å ²)	51.9	Xtriage
Anisotropy	0.159	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 38.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.031 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4070	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GNP, ANP, MG, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/350	0.65	0/468
2	B	0.43	0/2321	0.68	1/3143 (0.0%)
3	C	0.43	0/1392	0.64	0/1895
All	All	0.43	0/4063	0.67	1/5506 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	555	ARG	NE-CZ-NH1	5.07	122.83	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	341	0	337	1	0
2	B	2284	0	2338	6	0
3	C	1364	0	1378	8	0
4	B	23	0	12	0	0
5	C	32	0	13	0	0
6	C	1	0	0	0	0
7	B	23	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	C	2	0	0	0	0
All	All	4070	0	4078	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:98:VAL:HG21	3:C:149:LEU:HD21	1.76	0.68
2:B:528:ARG:O	2:B:555:ARG:NH2	2.29	0.65
3:C:145:LEU:O	3:C:149:LEU:HB2	2.03	0.59
2:B:370:MET:SD	2:B:381:MET:HB2	2.48	0.54
3:C:24:THR:HG22	3:C:42:VAL:HG11	1.91	0.52
2:B:399:GLU:H	2:B:399:GLU:CD	2.16	0.48
2:B:369:ILE:HG21	2:B:459:PHE:CD1	2.49	0.47
3:C:7:VAL:HG22	3:C:56:PHE:HB2	1.99	0.45
3:C:149:LEU:N	3:C:150:LYS:HA	2.34	0.43
2:B:461:PHE:HB3	2:B:476:VAL:CG1	2.48	0.42
3:C:14:VAL:HG13	3:C:83:SER:N	2.35	0.42
2:B:448:LEU:CD2	2:B:454:VAL:HG22	2.49	0.42
3:C:8:VAL:HG22	3:C:79:LEU:HD12	2.02	0.41
1:A:22:HIS:ND1	3:C:38:ASP:OD1	2.53	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	38/48 (79%)	35 (92%)	3 (8%)	0	100	100
2	B	286/346 (83%)	277 (97%)	9 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	C	171/185 (92%)	165 (96%)	6 (4%)	0	100	100
All	All	495/579 (86%)	477 (96%)	18 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	37/43 (86%)	35 (95%)	2 (5%)	22	36
2	B	251/300 (84%)	248 (99%)	3 (1%)	71	85
3	C	155/165 (94%)	154 (99%)	1 (1%)	86	94
All	All	443/508 (87%)	437 (99%)	6 (1%)	67	82

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	ARG
1	A	37	ARG
2	B	381	MET
2	B	444	ASP
2	B	464	GLN
3	C	100	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	426	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SEP	B	474	2	8,9,10	0.73	0	8,12,14	1.17	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SEP	B	474	2	-	0/5/8/10	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	474	SEP	O3P-P-O1P	2.05	118.72	110.68

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	ANP	B	600	-	22,25,33	1.05	1 (4%)	25,38,52	1.34	4 (16%)
5	GNP	C	200	6	28,34,34	3.24	9 (32%)	30,54,54	1.32	4 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ANP	B	600	-	-	0/6/26/38	0/3/3/3
5	GNP	C	200	6	-	12/17/38/38	0/3/3/3

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	200	GNP	C4-N9	-10.84	1.33	1.47
5	C	200	GNP	PB-O1B	9.94	1.61	1.46
5	C	200	GNP	PB-N3B	4.18	1.74	1.63
5	C	200	GNP	PG-N3B	4.01	1.73	1.63
5	C	200	GNP	PB-O2B	-3.16	1.48	1.56
5	C	200	GNP	C8-N9	-2.98	1.35	1.45
4	B	600	ANP	C5-C4	2.51	1.47	1.40
5	C	200	GNP	C5-C6	-2.20	1.49	1.52
5	C	200	GNP	PB-O3A	2.10	1.61	1.59
5	C	200	GNP	C2-N1	-2.05	1.36	1.44

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	600	ANP	N3-C2-N1	-3.71	122.88	128.68
5	C	200	GNP	PA-O3A-PB	-3.32	120.92	132.62
5	C	200	GNP	C4-C5-N7	3.11	106.58	102.46
5	C	200	GNP	O1G-PG-N3B	-2.99	107.36	111.77
4	B	600	ANP	C4-C5-N7	-2.30	107.00	109.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	600	ANP	O2A-PA-O1A	2.25	119.51	110.68
5	C	200	GNP	O1B-PB-N3B	-2.07	108.72	111.77
4	B	600	ANP	C2-N1-C6	2.06	122.28	118.75

There are no chirality outliers.



All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	200	GNP	PB-N3B-PG-O1G
5	C	200	GNP	PA-O3A-PB-O1B
5	C	200	GNP	PA-O3A-PB-O2B
5	C	200	GNP	C5'-O5'-PA-O3A
5	C	200	GNP	O4'-C1'-N9-C4
5	C	200	GNP	C2'-C1'-N9-C8
5	C	200	GNP	C2'-C1'-N9-C4
5	C	200	GNP	O4'-C4'-C5'-O5'
5	C	200	GNP	C3'-C4'-C5'-O5'
5	C	200	GNP	C5'-O5'-PA-O1A
5	C	200	GNP	C5'-O5'-PA-O2A
5	C	200	GNP	PG-N3B-PB-O1B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	40/48 (83%)	0.43	2 (5%) 28 27	56, 84, 123, 139	0
2	B	288/346 (83%)	-0.00	2 (0%) 87 86	32, 54, 80, 96	0
3	C	175/185 (94%)	0.98	33 (18%) 1 1	46, 78, 146, 171	0
All	All	503/579 (86%)	0.37	37 (7%) 14 13	32, 63, 113, 171	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	129	LEU	6.6
3	C	125	THR	6.4
3	C	140	GLU	6.1
3	C	127	GLU	5.9
3	C	131	LYS	5.5
3	C	128	LYS	5.2
3	C	130	ALA	5.1
3	C	121	ASP	4.6
3	C	120	ARG	4.5
3	C	87	PRO	4.4
3	C	30	SER	4.4
3	C	29	PRO	4.1
3	C	133	LYS	4.1
3	C	112	LEU	4.0
3	C	85	VAL	3.8
3	C	147	ARG	3.7
3	C	90	PHE	3.7
3	C	146	ALA	3.7
3	C	123	PRO	3.5
3	C	132	ASN	3.5
3	C	126	ILE	3.4
1	A	30	GLN	3.2
3	C	122	ASP	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	124	SER	2.9
3	C	31	GLU	2.9
3	C	84	VAL	2.8
3	C	80	VAL	2.7
1	A	28	HIS	2.4
3	C	94	LYS	2.4
3	C	82	PHE	2.4
3	C	148	ASP	2.4
2	B	586	ARG	2.4
3	C	108	THR	2.3
3	C	149	LEU	2.2
3	C	119	LEU	2.2
2	B	467	LYS	2.1
3	C	152	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SEP	B	474	10/11	0.99	0.15	43,47,48,49	0

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	MG	C	201	1/1	0.92	0.08	72,72,72,72	0
4	ANP	B	600	23/31	0.93	0.14	47,57,87,88	0
5	GNP	C	200	32/32	0.94	0.15	65,83,93,94	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.