



wwPDB EM Validation Summary Report ⓘ

Nov 12, 2022 – 03:16 PM EST

PDB ID : 6URO
EMDB ID : EMD-20861
Title : Cryo-EM structure of human CPSF160-WDR33-CPSF30-PAS RNA-CstF77 complex
Authors : Sun, Y.; Zhang, Y.; Walz, T.; Tong, L.
Deposited on : 2019-10-23
Resolution : 3.60 Å (reported)
Based on initial models : 6DNH, 2OOE

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1443	<div> <div>10%</div> <div>65%</div> <div>18%</div> <div>•</div> <div>17%</div> </div>
2	B	587	<div> <div>9%</div> <div>46%</div> <div>18%</div> <div>•</div> <div>36%</div> </div>
3	C	250	<div> <div>29%</div> <div>44%</div> <div>•</div> <div>54%</div> </div>
4	D	47	<div> <div>•</div> <div>15%</div> <div>•</div> <div>83%</div> </div>
5	E	717	<div> <div>48%</div> <div>62%</div> <div>12%</div> <div>26%</div> </div>
5	F	717	<div> <div>44%</div> <div>62%</div> <div>12%</div> <div>26%</div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 22401 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cleavage and polyadenylation specificity factor subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	1199	Total	C	N	O	S	0	0
			9498	6100	1626	1717	55		

- Molecule 2 is a protein called pre-mRNA 3' end processing protein WDR33.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	378	Total	C	N	O	S	0	0
			3051	1926	560	545	20		

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	MET	-	expression tag	UNP Q9C0J8
B	-13	GLY	-	expression tag	UNP Q9C0J8
B	-12	SER	-	expression tag	UNP Q9C0J8
B	-11	SER	-	expression tag	UNP Q9C0J8
B	-10	HIS	-	expression tag	UNP Q9C0J8
B	-9	HIS	-	expression tag	UNP Q9C0J8
B	-8	HIS	-	expression tag	UNP Q9C0J8
B	-7	HIS	-	expression tag	UNP Q9C0J8
B	-6	HIS	-	expression tag	UNP Q9C0J8
B	-5	HIS	-	expression tag	UNP Q9C0J8
B	-4	SER	-	expression tag	UNP Q9C0J8
B	-3	SER	-	expression tag	UNP Q9C0J8
B	-2	GLY	-	expression tag	UNP Q9C0J8
B	-1	LEU	-	expression tag	UNP Q9C0J8
B	0	VAL	-	expression tag	UNP Q9C0J8

- Molecule 3 is a protein called Cleavage and polyadenylation specificity factor subunit 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	116	Total	C	N	O	S	0	0
			913	585	150	164	14		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	245	HIS	-	expression tag	UNP O95639
C	246	HIS	-	expression tag	UNP O95639
C	247	HIS	-	expression tag	UNP O95639
C	248	HIS	-	expression tag	UNP O95639
C	249	HIS	-	expression tag	UNP O95639
C	250	HIS	-	expression tag	UNP O95639

- Molecule 4 is a RNA chain called PAS RNA.


Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	8	Total	C	N	O	P	0	0
			172	78	35	51	8		

- Molecule 5 is a protein called Cleavage stimulation factor subunit 3.


Mol	Chain	Residues	Atoms					AltConf	Trace
5	E	527	Total	C	N	O	S	0	0
			4382	2815	741	803	23		
5	F	527	Total	C	N	O	S	0	0
			4382	2815	741	803	23		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
6	C	3	Total	Zn	0
			3	3	



• Molecule 5: Cleavage stimulation factor subunit 3



VAL	GLN	ASP
LEU	GLY	SER
LYS	PRO	ASP
ASP	PHE	GLU
GLU	VAL	ASP
VAL	GLN	GLU
ASP	VAL	GLU
ARG	ASP	LYS
LYS	GLU	GLY
PRO	LEU	ALA
GLU	MET	VAL
TYR	GLU	VAL
PRO	ILE	PRO
LYS	PHE	PRO
PRO	ARG	VAL
ASP	ARG	HIS
THR	CYS	ASP
GLN	LYS	ILE
GLN	ILE	TYR
MET	PRO	ARG
ILE	ASN	ALA
PRO	THR	ARG
PHE	VAL	GLN
GLN	GLU	GLN
PRO	GLU	LYS
ARG	ALA	ARG
HIS	VAL	ILE
LEU	ARG	ARG
ALA	ILE	ILE
PRO	ILE	
PRO	THR	
GLY	GLY	
LEU	GLY	
HIS	ALA	
PRO	PRO	
VAL	GLU	
PRO	LEU	
GLY	ALA	
PHE	VAL	
PRO	GLY	
ASN	GLY	
PRO	VAL	
PRO	VAL	
ALA	GLU	
ALA	SER	
VAL	ASN	
VAL	ALA	
LEU	VAL	
MET	LEU	
LYS	THR	
LEU	LYS	
ALA	ALA	
VAL	VAL	
PRO	LYS	
PRO	ARG	
ILE	PRO	
CYS	ASN	
PHE	GLU	

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	50092	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	70	Depositor
Minimum defocus (nm)	900	Depositor
Maximum defocus (nm)	2800	Depositor
Magnification	46729	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.105	Depositor
Minimum map value	-0.061	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.02	Depositor
Map size (Å)	299.6, 299.6, 299.6	wwPDB
Map dimensions	280, 280, 280	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.07, 1.07, 1.07	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/9717	0.65	0/13194
2	B	0.53	0/3136	0.67	0/4246
3	C	0.35	0/937	0.56	0/1256
4	D	0.44	0/193	0.87	0/298
5	E	0.31	0/4479	0.46	0/6030
5	F	0.31	0/4479	0.46	0/6030
All	All	0.43	0/22941	0.59	0/31054

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9498	0	9538	200	0
2	B	3051	0	2966	93	0
3	C	913	0	884	5	0
4	D	172	0	88	0	0
5	E	4382	0	4378	54	0
5	F	4382	0	4378	51	0
6	C	3	0	0	0	0
All	All	22401	0	22232	380	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 380 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1346:GLY:C	1:A:1347:LEU:HD12	1.54	1.28
2:B:58:ASP:HA	2:B:149:LEU:CD2	1.65	1.24
5:F:510:THR:HG23	5:F:513:LYS:CE	1.73	1.19
2:B:58:ASP:HA	2:B:149:LEU:HD22	1.24	1.14
1:A:1126:CYS:SG	1:A:1172:VAL:N	2.25	1.10

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1177/1443 (82%)	1057 (90%)	118 (10%)	2 (0%)	47	79
2	B	376/587 (64%)	333 (89%)	43 (11%)	0	100	100
3	C	114/250 (46%)	94 (82%)	20 (18%)	0	100	100
5	E	523/717 (73%)	510 (98%)	13 (2%)	0	100	100
5	F	523/717 (73%)	511 (98%)	12 (2%)	0	100	100
All	All	2713/3714 (73%)	2505 (92%)	206 (8%)	2 (0%)	54	83

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	256	TRP
1	A	254	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1047/1235 (85%)	1037 (99%)	10 (1%)	76	88
2	B	328/514 (64%)	324 (99%)	4 (1%)	71	87
3	C	99/217 (46%)	99 (100%)	0	100	100
5	E	465/627 (74%)	457 (98%)	8 (2%)	60	82
5	F	465/627 (74%)	457 (98%)	8 (2%)	60	82
All	All	2404/3220 (75%)	2374 (99%)	30 (1%)	72	87

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	E	162	ASN
5	F	461	ASN
5	E	433	PHE
5	F	547	LYS
5	F	339	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
5	E	299	GLN
5	F	162	ASN
5	E	461	ASN
5	F	262	ASN
2	B	215	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
4	D	7/47 (14%)	1 (14%)	0

All (1) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
4	D	2	A

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-20861. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 140



Y Index: 140




Z Index: 140


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 142



Y Index: 130



Z Index: 133

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)

6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.02. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 98 nm³; this corresponds to an approximate mass of 88 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.278 \AA^{-1}


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-20861 and PDB model 6URO. Per-residue inclusion information can be found in [section 3](#) on [page 5](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.02 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.02).

9.4 Atom inclusion [i](#)



At the recommended contour level, 64% of all backbone atoms, 52% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.02) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.5191	<div></div> 0.3480
A	<div></div> 0.6696	<div></div> 0.4370
B	<div></div> 0.6572	<div></div> 0.4220
C	<div></div> 0.3227	<div></div> 0.2660
D	<div></div> 0.5756	<div></div> 0.4170
E	<div></div> 0.2971	<div></div> 0.2020
F	<div></div> 0.3561	<div></div> 0.2650

1.0
0.0
-0.0