



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 27, 2022 – 04:33 PM EDT

PDB ID : 7URQ
Title : Crystal Structure of SARS-CoV-2 S delta variant receptor-binding domain (RBD) in complex CoV11 Fab crystal form 1
Authors : Tolbert, W.D.; Pazgier, M.
Deposited on : 2022-04-22
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.28.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.28.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	229	<div> <div>8%</div> <div> <div></div> <div>73%</div> <div>11%</div> <div>•</div> <div>14%</div> </div> </div>
2	H	221	<div> <div>2%</div> <div> <div></div> <div>87%</div> <div>9%</div> <div>••</div> </div> </div>
3	L	216	<div> <div>2%</div> <div> <div></div> <div>94%</div> <div>6%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	CIT	H	301	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5095 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	196	Total	C	N	O	S	0	0	0
			1559	999	264	288	8			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	317	GLY	-	expression tag	UNP P0DTC2
A	318	SER	-	expression tag	UNP P0DTC2
A	452	ARG	LEU	engineered mutation	UNP P0DTC2
A	478	LYS	THR	engineered mutation	UNP P0DTC2
A	538	LEU	-	expression tag	UNP P0DTC2
A	539	GLU	-	expression tag	UNP P0DTC2
A	540	HIS	-	expression tag	UNP P0DTC2
A	541	HIS	-	expression tag	UNP P0DTC2
A	542	HIS	-	expression tag	UNP P0DTC2
A	543	HIS	-	expression tag	UNP P0DTC2
A	544	HIS	-	expression tag	UNP P0DTC2
A	545	HIS	-	expression tag	UNP P0DTC2

- Molecule 2 is a protein called COV11 Fab HEAVY CHAIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	214	Total	C	N	O	S	0	1	0
			1606	1016	270	314	6			

- Molecule 3 is a protein called COV11 Fab LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	215	Total	C	N	O	S	0	1	0
			1660	1035	285	336	4			


- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:

C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is CITRIC ACID (three-letter code: CIT) (formula: C₆H₈O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	H	1	Total	C	O	0	0
			13	6	7		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	69	Total 69	O 69	0	0
6	H	86	Total 86	O 86	0	0
6	L	88	Total 88	O 88	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Spike protein S1



- Molecule 2: COV11 Fab HEAVY CHAIN



- Molecule 3: COV11 Fab LIGHT CHAIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2 ₁	Depositor
Cell constants a, b, c, α , β , γ	85.82Å 103.55Å 112.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.23 – 2.05 42.74 – 2.04	Depositor EDS
% Data completeness (in resolution range)	92.8 (41.23-2.05) 92.8 (42.74-2.04)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.11 (at 2.03Å)	Xtriage
Refinement program	PHENIX (1.19.1_4122: ???)	Depositor
R, R_{free}	0.177 , 0.208 0.176 , 0.206	Depositor DCC
R_{free} test set	3070 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å ²)	40.1	Xtriage
Anisotropy	0.299	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 42.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5095	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CIT, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.43	0/1603	0.62	0/2177
2	H	0.44	0/1641	0.64	0/2234
3	L	0.42	0/1696	0.64	0/2302
All	All	0.43	0/4940	0.64	0/6713

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1559	0	1486	19	0
2	H	1606	0	1587	11	0
3	L	1660	0	1607	5	0
4	A	14	0	13	0	0
5	H	13	0	5	2	0
6	A	69	0	0	1	0
6	H	86	0	0	3	0
6	L	88	0	0	0	0
All	All	5095	0	4698	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:516:GLU:HG2	1:A:518:LEU:CD1	2.01	0.90
1:A:516:GLU:CG	1:A:518:LEU:HD11	2.12	0.79
3:L:150:LYS:NZ	3:L:196:GLU:OE2	2.21	0.74
1:A:516:GLU:HG2	1:A:518:LEU:HD13	1.68	0.73
1:A:516:GLU:CG	1:A:518:LEU:CD1	2.68	0.70
1:A:516:GLU:CD	1:A:518:LEU:HD11	2.14	0.68
1:A:360:ASN:N	1:A:523:THR:OG1	2.27	0.62
2:H:178:LEU:C	2:H:178:LEU:HD12	2.19	0.62
2:H:75:LYS:HB3	2:H:77:THR:HG23	1.82	0.60
1:A:393:THR:O	1:A:523:THR:HG23	2.02	0.59
1:A:365:TYR:CD2	1:A:387:LEU:HB3	2.39	0.58
1:A:389:ASP:OD1	1:A:528:LYS:HB2	2.05	0.57
1:A:366:SER:HA	1:A:369:TYR:CE1	2.40	0.56
1:A:458:LYS:HG3	6:A:1145:HOH:O	2.08	0.53
1:A:518:LEU:CD1	1:A:518:LEU:N	2.73	0.52
2:H:210:LYS:HD3	2:H:212:GLU:OE2	2.10	0.51
2:H:64:LYS:HD3	6:H:454:HOH:O	2.10	0.50
1:A:392:PHE:HD1	1:A:517:LEU:HD11	1.77	0.50
1:A:520:ALA:HB1	1:A:521:PRO:HD2	1.93	0.50
1:A:335:LEU:HD12	1:A:362:VAL:O	2.12	0.49
2:H:150:VAL:HG23	2:H:178:LEU:HD21	1.95	0.49
2:H:2:VAL:HA	2:H:25:SER:O	2.13	0.48
2:H:31:ARG:NH1	6:H:403:HOH:O	2.46	0.47
1:A:354:ASN:O	1:A:398:ASP:HA	2.15	0.46
2:H:105[A]:GLN:HB3	6:H:435:HOH:O	2.16	0.46
1:A:439:ASN:O	1:A:443:SER:HB2	2.17	0.45
3:L:160:SER:HA	3:L:179:THR:O	2.17	0.44
1:A:412:PRO:HG3	1:A:429:PHE:HB3	1.99	0.44
3:L:186:ASP:HA	3:L:189:LYS:HD3	2.00	0.44
5:H:301:CIT:O2	5:H:301:CIT:H42	2.17	0.44
2:H:71:ARG:NH2	5:H:301:CIT:H41	2.33	0.43
3:L:1:GLU:OE1	3:L:96:PRO:HD3	2.18	0.43
2:H:31:ARG:HA	2:H:31:ARG:HD3	1.77	0.41
2:H:12:ILE:HG21	2:H:82(C):LEU:HD13	2.02	0.41
1:A:359:SER:HA	1:A:524:VAL:HG22	2.02	0.40
3:L:146:LYS:HB3	3:L:198:THR:HB	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	194/229 (85%)	181 (93%)	13 (7%)	0	100	100
2	H	211/221 (96%)	206 (98%)	4 (2%)	1 (0%)	29	18
3	L	214/216 (99%)	211 (99%)	3 (1%)	0	100	100
All	All	619/666 (93%)	598 (97%)	20 (3%)	1 (0%)	47	39

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	100	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/201 (84%)	162 (96%)	7 (4%)	30	23
2	H	179/184 (97%)	170 (95%)	9 (5%)	24	16
3	L	185/185 (100%)	183 (99%)	2 (1%)	73	73
All	All	533/570 (94%)	515 (97%)	18 (3%)	39	30

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	346	ARG
1	A	369	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	389	ASP
1	A	408	ARG
1	A	443	SER
1	A	494	SER
1	A	523	THR
2	H	6	GLU
2	H	53	SER
2	H	77	THR
2	H	97	GLU
2	H	105[A]	GLN
2	H	105[B]	GLN
2	H	179	SER
2	H	186	SER
2	H	214	LYS
3	L	34	LEU
3	L	170	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	394	ASN
1	A	450	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	1001	1	14,14,15	0.88	2 (14%)	17,19,21	0.59	0
5	CIT	H	301	-	3,12,12	1.49	0	3,17,17	1.37	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1001	1	-	3/6/23/26	0/1/1/1
5	CIT	H	301	-	-	6/6/16/16	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1001	NAG	O5-C1	2.26	1.47	1.43
4	A	1001	NAG	C1-C2	2.25	1.55	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	H	301	CIT	C3-C2-C1	2.22	118.53	114.98

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	H	301	CIT	C1-C2-C3-O7
5	H	301	CIT	C1-C2-C3-C4
5	H	301	CIT	C1-C2-C3-C6
4	A	1001	NAG	C8-C7-N2-C2
4	A	1001	NAG	O7-C7-N2-C2
5	H	301	CIT	C2-C3-C4-C5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	1001	NAG	O5-C5-C6-O6
5	H	301	CIT	C6-C3-C4-C5
5	H	301	CIT	O7-C3-C4-C5

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	H	301	CIT	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	196/229 (85%)	0.46	19 (9%) 7 8	31, 53, 95, 131	0
2	H	214/221 (96%)	0.00	5 (2%) 60 64	30, 45, 74, 102	0
3	L	215/216 (99%)	0.04	5 (2%) 60 64	29, 46, 67, 86	0
All	All	625/666 (93%)	0.16	29 (4%) 32 35	29, 47, 84, 131	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	518	LEU	8.5
1	A	519	HIS	6.5
1	A	369	TYR	5.8
1	A	521	PRO	4.4
1	A	517	LEU	4.2
3	L	1	GLU	4.2
1	A	523	THR	4.2
1	A	335	LEU	3.8
1	A	520	ALA	3.7
2	H	1	GLU	3.6
2	H	128	SER	3.3
1	A	525	CYS	3.0
3	L	170	ARG	2.9
1	A	365	TYR	2.8
3	L	204	SER	2.7
1	A	522	ALA	2.6
2	H	134	GLY	2.5
2	H	214	LYS	2.4
3	L	127	LYS	2.4
1	A	393	THR	2.4
1	A	447	GLY	2.3
1	A	334	ASN	2.2
1	A	386	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	L	35	ALA	2.2
1	A	394	ASN	2.2
1	A	363	ALA	2.2
1	A	385	THR	2.2
2	H	127	SER	2.2
1	A	391	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	CIT	H	301	13/13	0.59	0.42	72,86,97,109	0
4	NAG	A	1001	14/15	0.80	0.30	77,95,111,112	0

6.5 Other polymers [i](#)

There are no such residues in this entry.