



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 20, 2020 – 06:50 PM BST

PDB ID : 5USD  
Title : Crystal structure of MccF-like protein (BA\_5613) in the complex with aspartyl sulfamoyl adenylate  
Authors : Nocek, B.; Anderson, W.F.; Joachimiak, A.; Center for Structural Genomics of Infectious Diseases (CSGID)  
Deposited on : 2017-02-13  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, orange 1%, orange 97%, yellow 97%, yellow 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>97%</span> <span>.</span> </div> </div>
1	B	330	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 3%, orange 3%, orange 95%, yellow 95%, yellow 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>3%</span> <span>95%</span> <span>5%</span> </div> </div>
1	C	330	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 2%, orange 2%, orange 97%, yellow 97%, yellow 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>2%</span> <span>97%</span> <span>.</span> </div> </div>
1	D	330	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 11%, orange 11%, orange 93%, yellow 93%, yellow 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>11%</span> <span>93%</span> <span>7%</span> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 21149 atoms, of which 10170 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Peptidase S66.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	329	Total	C	H	N	O	S	Se	0	0	0
			5064	1624	2507	422	500	4	7			
1	B	330	Total	C	H	N	O	S	Se	0	1	0
			5120	1637	2540	428	504	4	7			
1	C	329	Total	C	H	N	O	S	Se	0	1	0
			5089	1630	2521	427	500	4	7			
1	D	330	Total	C	H	N	O	S	Se	0	0	0
			5092	1631	2520	426	504	4	7			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	ASN	-	expression tag	UNP Q81JT5
A	0	ALA	-	expression tag	UNP Q81JT5
B	-1	ASN	-	expression tag	UNP Q81JT5
B	0	ALA	-	expression tag	UNP Q81JT5
C	-1	ASN	-	expression tag	UNP Q81JT5
C	0	ALA	-	expression tag	UNP Q81JT5
D	-1	ASN	-	expression tag	UNP Q81JT5
D	0	ALA	-	expression tag	UNP Q81JT5

- Molecule 2 is 5'-O-(L-alpha-aspartylsulfamoyl)adenosine (three-letter code: DSZ) (formula: C<sub>14</sub>H<sub>19</sub>N<sub>7</sub>O<sub>9</sub>S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	S	0	0
			47	14	16	7	9	1		
2	B	1	Total	C	H	N	O	S	0	0
			48	14	17	7	9	1		
2	C	1	Total	C	H	N	O	S	0	0
			47	14	16	7	9	1		
2	D	1	Total	C	H	N	O	S	0	0
			48	14	17	7	9	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			14	3	8	3		
3	C	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	205	Total	O	0	0
			205	205		
4	B	165	Total	O	0	0
			165	165		
4	C	136	Total	O	0	0
			136	136		
4	D	60	Total	O	0	0
			60	60		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Peptidase S66





- Molecule 1: Peptidase S66



- Molecule 1: Peptidase S66



- Molecule 1: Peptidase S66



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.46 Å 85.81 Å 124.80 Å 90.00° 92.32° 90.00°	Depositor
Resolution (Å)	32.61 – 2.10 32.61 – 2.09	Depositor EDS
% Data completeness (in resolution range)	91.1 (32.61-2.10) 81.1 (32.61-2.09)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.21 (at 2.10 Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, $R_{free}$	0.181 , 0.210 0.183 , 0.211	Depositor DCC
$R_{free}$ test set	3795 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.0	Xtriage
Anisotropy	0.222	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 43.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.029 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	21149	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, DSZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.25	0/2603	0.46	0/3516
1	B	0.25	0/2629	0.46	0/3549
1	C	0.25	0/2617	0.46	0/3534
1	D	0.25	0/2618	0.47	0/3536
All	All	0.25	0/10467	0.46	0/14135

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2557	2507	2506	4	0
1	B	2580	2540	2538	8	0
1	C	2568	2521	2521	5	0
1	D	2572	2520	2518	11	0
2	A	31	16	18	1	0
2	B	31	17	18	1	0
2	C	31	16	18	0	0
2	D	31	17	18	1	0
3	A	6	8	8	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	6	8	8	0	0
4	A	205	0	0	2	4
4	B	165	0	0	1	0
4	C	136	0	0	0	4
4	D	60	0	0	1	0
All	All	10979	10170	10171	28	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 28 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:401:DSZ:OD1	4:A:501:HOH:O	2.11	0.68
1:D:5:LYS:NZ	1:D:328:HIS:O	2.27	0.67
1:A:101:ARG:NH2	4:A:502:HOH:O	2.21	0.66
2:B:500:DSZ:OD1	4:B:601:HOH:O	2.12	0.66
1:D:190:SER:OG	1:D:316:SER:OG	2.16	0.63

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:690:HOH:O	4:C:631:HOH:O[2_645]	1.92	0.28
4:A:674:HOH:O	4:C:633:HOH:O[2_645]	1.95	0.25
4:A:690:HOH:O	4:C:627:HOH:O[2_645]	2.15	0.05
4:A:525:HOH:O	4:C:631:HOH:O[2_645]	2.17	0.03

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	327/330 (99%)	318 (97%)	9 (3%)	0	100	100
1	B	329/330 (100%)	322 (98%)	7 (2%)	0	100	100
1	C	328/330 (99%)	320 (98%)	8 (2%)	0	100	100
1	D	328/330 (99%)	320 (98%)	8 (2%)	0	100	100
All	All	1312/1320 (99%)	1280 (98%)	32 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	283/285 (99%)	280 (99%)	3 (1%)	73	79
1	B	287/285 (101%)	284 (99%)	3 (1%)	76	82
1	C	284/285 (100%)	279 (98%)	5 (2%)	59	65
1	D	285/285 (100%)	280 (98%)	5 (2%)	59	65
All	All	1139/1140 (100%)	1123 (99%)	16 (1%)	67	73

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	156	SER
1	C	209	ASN
1	D	209	ASN
1	C	35	ARG
1	D	221	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	DSZ	D	500	-	27,33,33	2.20	6 (22%)	29,49,49	2.28	7 (24%)
2	DSZ	B	500	-	27,33,33	2.31	6 (22%)	29,49,49	1.94	5 (17%)
3	GOL	C	402	-	5,5,5	0.38	0	5,5,5	0.21	0
3	GOL	A	402	-	5,5,5	0.36	0	5,5,5	0.25	0
2	DSZ	A	401	-	27,33,33	2.29	5 (18%)	29,49,49	1.96	5 (17%)
2	DSZ	C	401	-	27,33,33	2.33	6 (22%)	29,49,49	1.91	6 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DSZ	D	500	-	-	2/16/39/39	0/3/3/3
2	DSZ	B	500	-	-	0/16/39/39	0/3/3/3
3	GOL	C	402	-	-	4/4/4/4	-
3	GOL	A	402	-	-	4/4/4/4	-
2	DSZ	A	401	-	-	0/16/39/39	0/3/3/3
2	DSZ	C	401	-	-	4/16/39/39	0/3/3/3

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	401	DSZ	OAX-SBE	7.27	1.48	1.42
2	B	500	DSZ	OAX-SBE	7.11	1.48	1.42
2	A	401	DSZ	OAX-SBE	7.07	1.48	1.42
2	A	401	DSZ	OAY-SBE	6.66	1.48	1.42
2	B	500	DSZ	OAY-SBE	6.52	1.48	1.42

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	500	DSZ	OAY-SBE-OAX	-9.00	106.74	120.76
2	A	401	DSZ	OAY-SBE-OAX	-7.62	108.88	120.76
2	C	401	DSZ	OAY-SBE-OAX	-7.41	109.22	120.76
2	B	500	DSZ	OAY-SBE-OAX	-7.24	109.48	120.76
2	D	500	DSZ	C-NAT-SBE	-4.28	117.68	124.61

There are no chirality outliers.

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	500	DSZ	C3'-C4'-C5'-O5'
3	C	402	GOL	C1-C2-C3-O3
3	A	402	GOL	C1-C2-C3-O3
3	A	402	GOL	O2-C2-C3-O3
3	C	402	GOL	O1-C1-C2-C3


There are no ring outliers.


3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	500	DSZ	1	0
2	B	500	DSZ	1	0
2	A	401	DSZ	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	322/330 (97%)	-0.12	3 (0%) 84 86	15, 24, 39, 61	0
1	B	323/330 (97%)	-0.01	9 (2%) 53 59	16, 26, 42, 75	1 (0%)
1	C	322/330 (97%)	-0.08	7 (2%) 62 66	16, 25, 37, 61	0
1	D	323/330 (97%)	0.63	35 (10%) 5 7	24, 50, 70, 86	0
All	All	1290/1320 (97%)	0.10	54 (4%) 36 42	15, 28, 61, 86	1 (0%)

The worst 5 of 54 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	179	ASN	3.9
1	D	181	LYS	3.8
1	D	113	ILE	3.8
1	D	159	PRO	3.7
1	B	-1	ASN	3.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	DSZ	D	500	31/31	0.89	0.15	41,51,67,69	0
3	GOL	C	402	6/6	0.91	0.15	33,40,43,43	0
3	GOL	A	402	6/6	0.92	0.16	30,37,39,39	0
2	DSZ	A	401	31/31	0.94	0.13	24,30,56,57	0
2	DSZ	C	401	31/31	0.94	0.12	23,29,50,53	0
2	DSZ	B	500	31/31	0.95	0.13	25,33,56,58	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around DSZ D 500:**




2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around DSZ A 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around DSZ C 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around DSZ B 500:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.