



wwPDB EM Validation Summary Report ⓘ

Oct 29, 2022 – 12:30 PM EDT

PDB ID : 7USY
EMDB ID : EMD-26743
Title : Structure of C. elegans TMC-1 complex with ARRD-6
Authors : Jeong, H.; Clark, S.; Gouaux, E.
Deposited on : 2022-04-26
Resolution : 3.54 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev43
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4.02b-467
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.9
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1285	<div> <div>10%</div> <div>37%</div> <div>9%</div> <div>53%</div> </div>
1	B	1285	<div> <div>10%</div> <div>37%</div> <div>10%</div> <div>53%</div> </div>
2	C	201	<div> <div>24%</div> <div>63%</div> <div>27%</div> <div>8%</div> </div>
2	E	201	<div> <div>20%</div> <div>75%</div> <div>16%</div> <div>8%</div> </div>
3	D	117	<div> <div>8%</div> <div>32%</div> <div>8%</div> <div>61%</div> </div>
3	F	117	<div> <div>9%</div> <div>32%</div> <div>6%</div> <div>61%</div> </div>
4	J	469	<div> <div>73%</div> <div>63%</div> <div>14%</div> <div>23%</div> </div>

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 16582 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transmembrane channel-like protein 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	602	Total	C	N	O	S	0	0
			4864	3206	816	815	27		
1	B	602	Total	C	N	O	S	0	0
			4864	3206	816	815	27		

- Molecule 2 is a protein called CALMyrin (Calcium and Integrin Binding protein) homolog.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	C	184	Total	C	N	O	S	0	0
			1536	978	256	295	7		
2	E	184	Total	C	N	O	S	0	0
			1536	978	256	295	7		

- Molecule 3 is a protein called Transmembrane inner ear expressed protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	D	46	Total	C	N	O	S	0	0
			343	228	60	50	5		
3	F	46	Total	C	N	O	S	0	0
			343	228	60	50	5		

- Molecule 4 is a protein called ARRestin Domain protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	J	361	Total	C	N	O	S	0	0
			2878	1832	500	529	17		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		AltConf
5	A	2	Total	Ca	0
			2	2	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
5	B	2	Total	Ca	0
			2	2	
5	C	2	Total	Ca	0
			2	2	
5	E	2	Total	Ca	0
			2	2	

- Molecule 6 is 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (three-letter code: PEE) (formula: $C_{41}H_{78}NO_8P$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
6	A	1	Total	C	N	O	P	0
			42	32	1	8	1	
6	B	1	Total	C	N	O	P	0
			42	32	1	8	1	

- Molecule 7 is 1,2-Distearoyl-sn-glycerophosphoethanolamine (three-letter code: 3PE) (formula: $C_{41}H_{82}NO_8P$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
7	A	1	Total	C	N	O	P	0
			35	25	1	8	1	
7	B	1	Total	C	N	O	P	0
			35	25	1	8	1	

- Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
8	A	1	Total	C	N	O	0
			14	8	1	5	

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				AltConf
8	B	1	Total	C	N	O	0
			14	8	1	5	



- Molecule 9 is PALMITIC ACID (three-letter code: PLM) (formula: C₁₆H₃₂O₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			AltConf
9	D	1	Total	C	O	0
			14	13	1	
9	F	1	Total	C	O	0
			14	13	1	




• Molecule 1: Transmembrane channel-like protein 1







- Molecule 2: CALMyrin (Calcium and Integrin Binding protein) homolog




- Molecule 2: CALMyrin (Calcium and Integrin Binding protein) homolog









• Molecule 3: Transmembrane inner ear expressed protein



• Molecule 3: Transmembrane inner ear expressed protein



• Molecule 4: ARRestin Domain protein



ASN	GLY	ASN	GLN	G368	◆	Y308
SER	VAL	ASN	VAL	GLN	◆	A309
ARG	TYR	LYS	TYR	GLU	◆	L310
ARG	ASP	GLY	ASP	GLY	◆	R311
ARG	GLU	GLY	GLY	GLU	◆	V312
SER	ILE	GLY	GLY	GLY	◆	G313
LEU	LEU	GLY	GLY	GLY	◆	M314
VAL	VAL	GLY	ILE	ILE	◆	E315
ALA	ALA	ASN	ASN	ASN	◆	D316
SER	SER	ASN	GLY	GLY	◆	E317
ASN	PRO	GLY	GLY	GLY	◆	K318
PRO	PRO	GLY	GLY	GLY	◆	G319
CYS	LEU	CYS	E362	◆	◆	N320
LEU	ALA	ALA	E363	◆	◆	E321
MET	MET	ARG	I364	◆	◆	C322
ARG	ARG	ASP	V365	◆	◆	L323
ASP	GLY	GLY	L366	◆	◆	H324
SER	SER	SER	Y367	◆	◆	I325
MET	MET	ASP	R368	◆	◆	D326
ASP	ASP	GLY	P369	◆	◆	F327
GLY	GLY	LYS	V390	◆	◆	P328
LEU	LEU	LEU	Y391	◆	◆	L329
MET	MET	MET	V392	◆	◆	T330
MET	MET	THR	K393	◆	◆	V331
THR	THR	ASN	L394	◆	◆	A332
ASN	GLY	GLY	A395	◆	◆	T333
CYS	CYS	CYS	D396	◆	◆	I334
GLY	GLY	SER	R397	◆	◆	P335
SER	GLY	GLY	R398	◆	◆	Y336
GLY	GLY	GLY	I399	◆	◆	R337
PRO	PRO	ASP	G400	◆	◆	I338
LEU	LEU	LEU	SER	◆	◆	P339
VAL	VAL	VAL	HIS	◆	◆	N340
ALA	ALA	SER	VAL	◆	◆	A341
		LYS	LYS	◆	◆	P342
		ASP	ASP	◆	◆	P343
		PHE	PHE	◆	◆	P344
		ARG	ARG	◆	◆	P345
		SER	SER	◆	◆	V346
		GLY	GLY	◆	◆	D347
		SER	SER	◆	◆	Y348
		PHE	PHE	◆	◆	D349
		THR	THR	◆	◆	F350
		ARG	ARG	◆	◆	A351
		ILE	ILE	◆	◆	S352
		ALA	ALA	◆	◆	N353
		ASP	ASP	◆	◆	H354
		SER	SER	◆	◆	V355
		LEU	LEU	◆	◆	E356
		VAL	VAL	◆	◆	G357
		THR	THR	◆	◆	G358
		GLY	GLY	◆	◆	K359
		LEU	LEU	◆	◆	Y360
		PRO	PRO	◆	◆	V361
				◆	◆	S362
				◆	◆	P363
				◆	◆	E364
				◆	◆	F365
				◆	◆	R366
				◆	◆	I367

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	99248	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2400	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	66.937	Depositor
Minimum map value	-42.398	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	1.000	Depositor
Recommended contour level	6.0	Depositor
Map size (\AA)	335.6, 335.6, 335.6	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	0.839, 0.839, 0.839	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 3PE, PLM, NAG, PEE, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/4987	0.50	0/6763
1	B	0.26	0/4987	0.47	0/6763
2	C	0.25	0/1569	0.50	0/2111
2	E	0.26	0/1569	0.52	0/2111
3	D	0.24	0/347	0.56	0/471
3	F	0.24	0/347	0.53	0/471
4	J	0.25	0/2938	0.52	0/3979
All	All	0.25	0/16744	0.50	0/22669

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4864	0	4941	84	0
1	B	4864	0	4941	90	0
2	C	1536	0	1484	39	0
2	E	1536	0	1484	20	0
3	D	343	0	377	5	0
3	F	343	0	378	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	J	2878	0	2908	40	0
5	A	2	0	0	0	0
5	B	2	0	0	0	0
5	C	2	0	0	0	0
5	E	2	0	0	0	0
6	A	42	0	58	4	0
6	B	42	0	58	4	0
7	A	35	0	44	3	0
7	B	35	0	44	0	0
8	A	14	0	13	0	0
8	B	14	0	13	1	0
9	D	14	0	22	4	0
9	F	14	0	22	1	0
All	All	16582	0	16787	278	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 278 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:44:CYS:SG	9:F:201:PLM:C1	2.38	1.11
9:D:201:PLM:C9	9:D:201:PLM:CD	2.48	0.92
1:A:868:LEU:HD23	1:A:871:LEU:HD12	1.49	0.92
9:D:201:PLM:CD	9:D:201:PLM:H92	2.05	0.85
1:A:378:VAL:HG11	1:A:446:ALA:HB1	1.64	0.79

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	598/1285 (46%)	577 (96%)	21 (4%)	0	100	100
1	B	598/1285 (46%)	579 (97%)	19 (3%)	0	100	100
2	C	182/201 (90%)	181 (100%)	1 (0%)	0	100	100
2	E	182/201 (90%)	178 (98%)	4 (2%)	0	100	100
3	D	44/117 (38%)	43 (98%)	1 (2%)	0	100	100
3	F	44/117 (38%)	42 (96%)	2 (4%)	0	100	100
4	J	357/469 (76%)	345 (97%)	12 (3%)	0	100	100
All	All	2005/3675 (55%)	1945 (97%)	60 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	517/1150 (45%)	507 (98%)	10 (2%)	57	80
1	B	517/1150 (45%)	505 (98%)	12 (2%)	50	77
2	C	171/185 (92%)	164 (96%)	7 (4%)	30	64
2	E	171/185 (92%)	166 (97%)	5 (3%)	42	72
3	D	35/96 (36%)	35 (100%)	0	100	100
3	F	35/96 (36%)	34 (97%)	1 (3%)	42	72
4	J	323/419 (77%)	315 (98%)	8 (2%)	47	76
All	All	1769/3281 (54%)	1726 (98%)	43 (2%)	51	76

5 of 43 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	188	SER
4	J	34	PHE
2	E	28	TYR
2	E	152	ASP
4	J	44	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	731	HIS
1	A	879	ASN
1	B	420	HIS
1	B	879	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	NAG	B	1305	-	14,14,15	0.46	0	19,19,21	0.73	0
7	3PE	B	1304	-	34,34,50	1.03	4 (11%)	37,39,55	1.13	2 (5%)
8	NAG	A	1305	-	14,14,15	0.47	0	19,19,21	0.73	0
6	PEE	B	1303	-	41,41,50	1.26	6 (14%)	44,46,55	1.23	2 (4%)
9	PLM	F	201	-	13,13,17	0.32	0	12,12,17	0.92	0
6	PEE	A	1303	-	41,41,50	1.25	6 (14%)	44,46,55	1.20	3 (6%)
7	3PE	A	1304	-	34,34,50	1.03	4 (11%)	37,39,55	1.16	2 (5%)
9	PLM	D	201	3	13,13,17	0.32	0	12,12,17	0.91	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	NAG	B	1305	-	-	2/6/22/26	0/1/1/1
7	3PE	B	1304	-	-	16/38/38/54	-
8	NAG	A	1305	-	-	0/6/22/26	0/1/1/1
6	PEE	B	1303	-	-	21/45/45/54	-
9	PLM	F	201	-	-	5/10/11/15	-
6	PEE	A	1303	-	-	25/45/45/54	-
7	3PE	A	1304	-	-	16/38/38/54	-
9	PLM	D	201	3	-	3/10/11/15	-

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	1303	PEE	C18-C19	3.70	1.53	1.31
6	A	1303	PEE	C18-C19	3.69	1.53	1.31
6	B	1303	PEE	C39-C38	3.69	1.53	1.31
6	A	1303	PEE	C39-C38	3.68	1.53	1.31
7	A	1304	3PE	O21-C2	-2.56	1.40	1.46

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	1303	PEE	O2-C10-C11	4.18	120.52	111.50
7	A	1304	3PE	O21-C21-C22	4.10	120.34	111.50
7	B	1304	3PE	O21-C21-C22	4.09	120.31	111.50
6	A	1303	PEE	O2-C10-C11	3.84	119.77	111.50
7	A	1304	3PE	O31-C31-C32	2.71	120.42	111.91

There are no chirality outliers.

5 of 88 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	1303	PEE	C1-O3P-P-O2P
6	A	1303	PEE	C1-O3P-P-O1P
6	A	1303	PEE	C4-O4P-P-O2P
6	A	1303	PEE	C4-O4P-P-O1P
6	A	1303	PEE	O4P-C4-C5-N



There are no ring outliers.




6 monomers are involved in 16 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	B	1305	NAG	1	0
6	B	1303	PEE	4	0
9	F	201	PLM	1	0
6	A	1303	PEE	4	0
7	A	1304	3PE	3	0
9	D	201	PLM	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-26743. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200


6.2.2 Raw map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 169




Y Index: 202




Z Index: 207


6.3.2 Raw map



X Index: 165



Y Index: 202




Z Index: 186

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 6.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.5 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 105 nm^3 ; this corresponds to an approximate mass of 94 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.282 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.282 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.54	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	4.06	8.03	4.10

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.06 differs from the reported value 3.54 by more than 10 %

9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-26743 and PDB model 7USY. Per-residue inclusion information can be found in section 3 on page 7.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 6.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (6.0).

9.4 Atom inclusion [i](#)



At the recommended contour level, 56% of all backbone atoms, 54% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (6.0) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div><div></div></div> 0.5409	<div><div></div></div> 0.3550
A	<div><div></div></div> 0.6526	<div><div></div></div> 0.3880
B	<div><div></div></div> 0.6608	<div><div></div></div> 0.3820
C	<div><div></div></div> 0.5899	<div><div></div></div> 0.3800
D	<div><div></div></div> 0.6638	<div><div></div></div> 0.4150
E	<div><div></div></div> 0.6198	<div><div></div></div> 0.3880
F	<div><div></div></div> 0.6149	<div><div></div></div> 0.4000
J	<div><div></div></div> 0.0488	<div><div></div></div> 0.2080

1.0

0.0

<0.0