



# Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 05:30 pm BST

PDB ID : 3UT2  
Title : Crystal Structure of Fungal MagKatG2  
Authors : Zamocky, M.; Garcia-Fernandez, M.Q.; Gasselhuber, B.; Jakopitsch, C.; Furtmuller, P.G.; Loewen, P.C.; Fita, I.; Obinger, C.; Carpena, X.  
Deposited on : 2011-11-24  
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2556 (1.56-1.52)
Clashscore	141614	2634 (1.56-1.52)
Ramachandran outliers	138981	2580 (1.56-1.52)
Sidechain outliers	138945	2577 (1.56-1.52)
RSRZ outliers	127900	2524 (1.56-1.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	764	<div> <div>4%</div> <div> <div></div> <div>92%</div> <div></div> </div> <div></div> </div>
1	B	764	<div> <div>2%</div> <div> <div></div> <div>92%</div> <div></div> </div> <div></div> </div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 13094 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Catalase-peroxidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	735	Total	C	N	O	S	0	9	0
			5696	3576	999	1098	23			
1	B	732	Total	C	N	O	S	0	12	0
			5697	3577	998	1100	22			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	23	MET	-	EXPRESSION TAG	UNP A4QUT2
B	23	MET	-	EXPRESSION TAG	UNP A4QUT2

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	801	Total	O	0	0
			801	801		
3	B	814	Total	O	0	0
			814	814		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Catalase-peroxidase 2



- Molecule 1: Catalase-peroxidase 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.99Å 109.61Å 132.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.55 19.98 – 1.55	Depositor EDS
% Data completeness (in resolution range)	96.9 (20.00-1.55) 96.9 (19.98-1.55)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	0.11	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.98 (at 1.55Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.197 , 0.220 0.210 , 0.232	Depositor DCC
$R_{free}$ test set	10526 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	12.5	Xtriage
Anisotropy	0.043	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 36.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	13094	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.26% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, TOX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.41	0/5841	0.59	0/7926
1	B	0.40	0/5851	0.59	0/7938
All	All	0.41	0/11692	0.59	0/15864

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5696	0	5499	25	0
1	B	5697	0	5513	25	0
2	A	43	0	30	1	0
2	B	43	0	30	0	0
3	A	801	0	0	4	0
3	B	814	0	0	11	0
All	All	13094	0	11072	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:119:GLU:HG2	3:B:2283:HOH:O	1.40	1.21
1:A:641[B]:ARG:HG2	1:A:641[B]:ARG:HH21	1.14	1.11
1:B:152[A]:ARG:NH2	3:B:1887:HOH:O	1.89	0.99
1:B:152[A]:ARG:NH1	3:B:1887:HOH:O	2.05	0.90
1:B:466[B]:THR:HG21	3:B:1534:HOH:O	1.70	0.90
1:A:466[B]:THR:HG21	3:A:2164:HOH:O	1.78	0.83
1:A:641[B]:ARG:HG2	1:A:641[B]:ARG:NH2	1.94	0.82
1:B:488[B]:ARG:NH2	3:B:1977:HOH:O	2.14	0.80
1:A:641[B]:ARG:CG	1:A:641[B]:ARG:HH21	1.97	0.77
1:B:152[A]:ARG:CZ	3:B:1887:HOH:O	2.26	0.71
1:B:400:VAL:HG23	1:B:401:ASN:HD22	1.56	0.70
1:B:721:VAL:HG13	1:B:730:GLU:HG3	1.73	0.70
1:B:523[B]:CYS:SG	1:B:541:ILE:HG21	2.34	0.68
1:A:85:GLN:HG3	1:B:85:GLN:OE1	1.96	0.65
1:A:523[C]:CYS:SG	1:A:541:ILE:HG21	2.40	0.61
1:A:497:ASP:OD2	1:A:578:LYS:HD2	2.03	0.58
1:A:721:VAL:HG13	1:A:730:GLU:HG3	1.85	0.58
1:B:488[B]:ARG:NH2	3:B:1394:HOH:O	2.34	0.58
1:A:783:LYS:O	1:A:784:VAL:CB	2.51	0.58
1:A:641[B]:ARG:CG	1:A:641[B]:ARG:NH2	2.63	0.57
1:A:591:THR:HG23	1:A:607:VAL:HG12	1.90	0.53
1:B:488[B]:ARG:NH1	3:B:2040:HOH:O	2.26	0.53
1:A:709:ALA:HB2	1:A:725:ARG:HD3	1.92	0.51
1:A:760:GLN:HG3	3:A:1707:HOH:O	2.11	0.49
1:B:523[B]:CYS:SG	1:B:541:ILE:CG2	3.01	0.49
1:A:523[C]:CYS:SG	1:A:541:ILE:CG2	3.01	0.48
1:B:51:THR:HG23	3:B:2178:HOH:O	2.13	0.48
1:B:140:TOX:H9	1:B:273:TYR:OH	2.14	0.47
1:B:523[B]:CYS:SG	1:B:586:ILE:HG21	2.55	0.47
1:B:400:VAL:HG23	1:B:401:ASN:ND2	2.28	0.46
1:A:466[B]:THR:HA	1:A:469:TYR:CD1	2.51	0.46
1:B:709:ALA:HB2	1:B:725:ARG:HD3	1.97	0.46
1:A:111:LYS:NZ	3:A:2082:HOH:O	2.30	0.45
1:B:75:GLN:NE2	3:B:1580:HOH:O	2.50	0.45
1:A:386:THR:CG2	1:A:400:VAL:HG13	2.47	0.44
1:A:432:GLU:OE2	3:A:2241:HOH:O	2.21	0.44
1:A:82:ARG:HG2	1:B:88:GLN:OE1	2.18	0.44
1:A:59:SER:HA	1:B:645:ARG:O	2.19	0.43
1:B:69[B]:HIS:CD2	3:B:1877:HOH:O	2.72	0.43
1:A:386:THR:HG21	1:A:400:VAL:CG1	2.49	0.42
1:B:304[A]:THR:HG23	1:B:440:TRP:CZ2	2.54	0.42

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:135:PHE:HB3	1:B:201:THR:HG22	2.01	0.41
1:A:508:THR:HB	1:A:511:LEU:HD12	2.01	0.41
1:A:721:VAL:CG1	1:A:730:GLU:HG3	2.50	0.41
1:A:88:GLN:OE1	1:B:82:ARG:HG2	2.20	0.41
1:A:319:THR:HG22	2:A:1500:HEM:HAA1	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	742/764 (97%)	727 (98%)	15 (2%)	0	100	100
1	B	742/764 (97%)	730 (98%)	12 (2%)	0	100	100
All	All	1484/1528 (97%)	1457 (98%)	27 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	588/605 (97%)	580 (99%)	8 (1%)	67	39
1	B	591/605 (98%)	588 (100%)	3 (0%)	88	77
All	All	1179/1210 (97%)	1168 (99%)	11 (1%)	78	60

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	THR
1	A	102	GLN
1	A	147	ARG
1	A	442	GLU
1	A	632[A]	GLN
1	A	632[B]	GLN
1	A	655	LYS
1	A	728	ARG
1	B	147	ARG
1	B	170	ASP
1	B	655	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	102	GLN
1	A	249	ASN
1	A	625	GLN
1	A	707	ASN
1	A	757	ASN
1	A	760	GLN
1	B	61	GLN
1	B	401	ASN
1	B	556	GLN
1	B	691	ASN
1	B	757	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	TOX	A	140	1	10,17,18	2.65	3 (30%)	10,23,25	1.33	1 (10%)
1	TOX	B	140	1,2	10,17,18	2.39	3 (30%)	10,23,25	1.55	2 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TOX	A	140	1	-	2/4/8/10	0/2/2/2
1	TOX	B	140	1,2	-	2/4/8/10	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	140	TOX	CD1-NE1	-5.23	1.34	1.39
1	A	140	TOX	O-C	5.14	1.40	1.19
1	B	140	TOX	O-C	4.88	1.39	1.19
1	B	140	TOX	CD1-NE1	-4.57	1.34	1.39
1	A	140	TOX	CH2-CZ2	2.53	1.42	1.36
1	B	140	TOX	CH2-CZ2	2.16	1.41	1.36

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	140	TOX	CZ3-CH2-CZ2	-2.83	116.47	120.44
1	A	140	TOX	CZ3-CH2-CZ2	-2.70	116.65	120.44
1	B	140	TOX	CB-CG-CD1	-2.68	124.66	127.97

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	140	TOX	N-CA-CB-CG
1	A	140	TOX	C-CA-CB-CG
1	B	140	TOX	N-CA-CB-CG
1	B	140	TOX	C-CA-CB-CG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	140	TOX	1	0

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	HEM	B	1500	1	27,50,50	1.96	5 (18%)	17,82,82	1.47	1 (5%)
2	HEM	A	1500	1	27,50,50	1.98	5 (18%)	17,82,82	1.74	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	B	1500	1	-	0/6/54/54	-
2	HEM	A	1500	1	-	0/6/54/54	-

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1500	HEM	C3D-C2D	4.70	1.51	1.37
2	A	1500	HEM	C3B-C2B	-4.65	1.33	1.40
2	B	1500	HEM	C3B-C2B	-4.61	1.34	1.40
2	A	1500	HEM	C3D-C2D	4.55	1.51	1.37

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1500	HEM	C3C-C2C	-4.12	1.34	1.40
2	B	1500	HEM	C3C-C2C	-3.58	1.35	1.40
2	B	1500	HEM	C3C-CAC	3.52	1.55	1.47
2	A	1500	HEM	C3B-CAB	3.25	1.54	1.47
2	A	1500	HEM	C3C-CAC	3.17	1.54	1.47
2	B	1500	HEM	C3B-CAB	2.81	1.53	1.47

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1500	HEM	CAA-CBA-CGA	-4.94	104.38	112.67
2	B	1500	HEM	CAA-CBA-CGA	-4.43	105.24	112.67
2	A	1500	HEM	C1D-C2D-C3D	-2.68	105.13	107.00
2	A	1500	HEM	CMC-C2C-C3C	2.10	128.61	124.68

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1500	HEM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	734/764 (96%)	0.22	28 (3%)	40 46	5, 11, 21, 34	0
1	B	731/764 (95%)	0.17	19 (2%)	56 62	6, 11, 21, 35	0
All	All	1465/1528 (95%)	0.20	47 (3%)	47 54	5, 11, 21, 35	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	784	VAL	10.5
1	A	50	THR	7.2
1	A	413	THR	5.4
1	A	400	VAL	5.3
1	B	573	GLY	5.1
1	B	94	ASP	4.6
1	A	412	LYS	4.4
1	A	714	GLY	4.3
1	B	728	ARG	4.2
1	A	726	LYS	4.2
1	B	51	THR	4.0
1	A	94	ASP	3.9
1	A	414	LYS	3.9
1	A	102	GLN	3.9
1	A	783	LYS	3.7
1	A	715	ALA	3.6
1	B	714	GLY	3.6
1	B	726	LYS	3.6
1	A	51	THR	3.5
1	B	93	GLY	3.3
1	A	401	ASN	3.3
1	A	728	ARG	2.9
1	B	69[A]	HIS	2.9
1	B	413	THR	2.9

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	401	ASN	2.9
1	A	69[A]	HIS	2.9
1	B	577	ASN	2.9
1	B	400	VAL	2.9
1	A	489	GLU	2.8
1	A	402	GLY	2.8
1	A	93	GLY	2.7
1	A	690	ALA	2.6
1	A	758	GLY	2.6
1	B	690	ALA	2.5
1	A	99	GLU	2.5
1	B	102	GLN	2.5
1	A	399	ALA	2.4
1	B	715	ALA	2.4
1	B	606	LYS	2.3
1	A	509	ASP	2.3
1	A	713	SER	2.3
1	A	56	ALA	2.3
1	B	555	THR	2.2
1	B	414	LYS	2.2
1	A	511	LEU	2.2
1	B	489	GLU	2.2
1	A	573	GLY	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	TOX	A	140	16/17	0.92	0.10	6,7,15,15	0
1	TOX	B	140	16/17	0.93	0.09	6,7,14,15	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	HEM	B	1500	43/43	0.97	0.09	6,8,11,12	0
2	HEM	A	1500	43/43	0.97	0.09	6,7,9,11	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




### Electron density around HEM B 1500:

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around HEM A 1500:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.