



wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 04:21 am BST

PDB ID : 2UUA
Title : Structure of the *Thermus thermophilus* 30S ribosomal subunit complexed with a Valine-ASL with cmo5U in position 34 bound to an mRNA with a GUC-codon in the A-site and paromomycin.
Authors : Weixlbaumer, A.; Murphy, F.V.; Dziergowska, A.; Malkiewicz, A.; Vendeix, F.A.P.; Agris, P.F.; Ramakrishnan, V.
Deposited on : 2007-03-01
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)
RNA backbone	3102	1007 (3.16-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1522	<div> <div>0%</div> <div> <div>41%</div> <div>47%</div> <div>9%</div> <div>..</div> </div> </div>
2	B	256	<div> <div>8%</div> <div> <div>27%</div> <div>53%</div> <div>11%</div> <div>8%</div> </div> </div>
3	C	239	<div> <div>2%</div> <div> <div>27%</div> <div>51%</div> <div>8%</div> <div>13%</div> </div> </div>
4	D	209	<div> <div>3%</div> <div> <div>47%</div> <div>46%</div> <div>6%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	162	
6	F	101	
7	G	156	
8	H	138	
9	I	128	
10	J	105	
11	K	129	
12	L	135	
13	M	126	
14	N	61	
15	O	89	
16	P	88	
17	Q	105	
18	R	88	
19	S	93	
20	T	106	
21	U	27	
22	X	6	
23	Y	17	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
25	MG	H	202	-	-	-	X
27	ZN	N	101	-	-	X	-

2 Entry composition [i](#)

There are 27 unique types of molecules in this entry. The entry contains 52344 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 16S RRNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1512	Total	C	N	O	P	0	0	0
			32489	14462	6011	10505	1511			

- Molecule 2 is a protein called 30S RIBOSOMAL PROTEIN S2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	235	Total	C	N	O	S	0	0	1
			1901	1213	342	341	5			

- Molecule 3 is a protein called 30S RIBOSOMAL PROTEIN S3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	207	Total	C	N	O	S	0	0	1
			1613	1016	315	281	1			

- Molecule 4 is a protein called 30S RIBOSOMAL PROTEIN S4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	208	Total	C	N	O	S	0	0	0
			1703	1066	339	291	7			

- Molecule 5 is a protein called 30S RIBOSOMAL PROTEIN S5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	151	Total	C	N	O	S	0	0	1
			1147	724	218	201	4			

- Molecule 6 is a protein called 30S RIBOSOMAL PROTEIN S6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	101	Total	C	N	O	S	0	0	0
			843	531	155	154	3			

- Molecule 7 is a protein called 30S RIBOSOMAL PROTEIN S7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	155	Total	C	N	O	S	0	0	0
			1257	781	252	218	6			

- Molecule 8 is a protein called 30S RIBOSOMAL PROTEIN S8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	138	Total	C	N	O	S	0	0	0
			1116	705	215	193	3			

- Molecule 9 is a protein called 30S RIBOSOMAL PROTEIN S9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	127	Total	C	N	O	S	0	0	0
			1010	639	197	174				

- Molecule 10 is a protein called 30S RIBOSOMAL PROTEIN S10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	99	Total	C	N	O	S	0	0	1
			793	498	157	137	1			

- Molecule 11 is a protein called 30S RIBOSOMAL PROTEIN S11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	119	Total	C	N	O	S	0	0	0
			885	549	168	165	3			

- Molecule 12 is a protein called 30S RIBOSOMAL PROTEIN S12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	125	Total	C	N	O	S	0	0	1
			971	611	196	163	1			

- Molecule 13 is a protein called 30S RIBOSOMAL PROTEIN S13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	M	125	Total	C	N	O	S	0	0	0
			997	617	207	171	2			

- Molecule 14 is a protein called 30S RIBOSOMAL PROTEIN S14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	N	60	Total	C	N	O	S	0	0	0
			492	312	104	72	4			

- Molecule 15 is a protein called 30S RIBOSOMAL PROTEIN S15.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
15	O	88	Total	C	N	O	S	0	0	0
			734	459	147	126	2			

- Molecule 16 is a protein called 30S RIBOSOMAL PROTEIN S16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
16	P	84	Total	C	N	O	S	0	0	1
			701	443	140	117	1			

- Molecule 17 is a protein called 30S RIBOSOMAL PROTEIN S17.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
17	Q	104	Total	C	N	O	S	0	0	0
			857	547	160	148	2			

- Molecule 18 is a protein called 30S RIBOSOMAL PROTEIN S18.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
18	R	73	Total	C	N	O	0	0	0
			598	381	118	99			

- Molecule 19 is a protein called 30S RIBOSOMAL PROTEIN S19.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
19	S	81	Total	C	N	O	S	0	0	1
			648	414	120	112	2			

- Molecule 20 is a protein called 30S RIBOSOMAL PROTEIN S20.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
20	T	99	Total	C	N	O	S	0	0	0
			763	470	162	129	2			

- Molecule 21 is a protein called 30S RIBOSOMAL PROTEIN THX.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
21	U	25	Total	C	N	O	0	0	1
			209	128	51	30			


- Molecule 22 is a RNA chain called 5'-R(*GP*UP*CP*AP*AP*AP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
22	X	5	Total	C	N	O	P	0	0	0
			104	48	20	32	4			

- Molecule 23 is a RNA chain called 5'-R(*CP*CP*UP*CP*CP*CP*UP*CM0P*AP*CP*6MZP*AP*GP*GP*AP*GP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
23	Y	13	Total	C	N	O	P	0	0	0
			277	126	48	91	12			

- Molecule 24 is PAROMOMYCIN (three-letter code: PAR) (formula: $C_{23}H_{45}N_5O_{14}$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
24	A	1	Total	C	N	O	0	0
			42	23	5	14		

- Molecule 25 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
25	G	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
25	Q	1	Total 1	Mg 1	0	0
25	D	1	Total 1	Mg 1	0	0
25	E	2	Total 2	Mg 2	0	0
25	H	2	Total 2	Mg 2	0	0
25	B	1	Total 1	Mg 1	0	0
25	A	161	Total 161	Mg 161	0	0
25	F	1	Total 1	Mg 1	0	0

- Molecule 26 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
26	A	21	Total 21	K 21	0	0
26	E	1	Total 1	K 1	0	0


- Molecule 27 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
27	D	1	Total 1	Zn 1	0	0
27	N	1	Total 1	Zn 1	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: 16S rRNA






- Molecule 3: 30S RIBOSOMAL PROTEIN S3







• Molecule 7: 30S RIBOSOMAL PROTEIN S7




• Molecule 8: 30S RIBOSOMAL PROTEIN S8




• Molecule 9: 30S RIBOSOMAL PROTEIN S9




• Molecule 10: 30S RIBOSOMAL PROTEIN S10




• Molecule 11: 30S RIBOSOMAL PROTEIN S11




- Molecule 12: 30S RIBOSOMAL PROTEIN S12



- Molecule 13: 30S RIBOSOMAL PROTEIN S13



- Molecule 14: 30S RIBOSOMAL PROTEIN S14




- Molecule 15: 30S RIBOSOMAL PROTEIN S15




- Molecule 16: 30S RIBOSOMAL PROTEIN S16

Chain P: 





• Molecule 17: 30S RIBOSOMAL PROTEIN S17

Chain Q: 




• Molecule 18: 30S RIBOSOMAL PROTEIN S18

Chain R: 




• Molecule 19: 30S RIBOSOMAL PROTEIN S19

Chain S: 



• Molecule 20: 30S RIBOSOMAL PROTEIN S20

Chain T: 



- Molecule 21: 30S RIBOSOMAL PROTEIN THX

Chain U: 





- Molecule 22: 5'-R(*GP*UP*CP*AP*AP*AP)-3'

Chain X: 



- Molecule 23: 5'-R(*CP*CP*UP*CP*CP*CP*UP*CM0P*AP*CP*6MZP*AP *GP*GP*AP*GP*G)-3'

Chain Y: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	401.92Å 401.92Å 174.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.88 – 2.90 284.20 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.3 (29.88-2.90) 99.3 (284.20-2.90)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 2.91Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.223 , 0.254 0.224 , 0.254	Depositor DCC
R_{free} test set	15908 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	63.7	Xtriage
Anisotropy	0.212	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 75.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	52344	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, PAR, K, ZN, 6MZ, CM0

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	1/36365 (0.0%)	0.70	29/56754 (0.1%)
2	B	0.41	0/1936	0.68	0/2611
3	C	0.43	0/1637	0.67	1/2207 (0.0%)
4	D	0.53	1/1733 (0.1%)	0.67	1/2318 (0.0%)
5	E	0.56	0/1163	0.77	1/1566 (0.1%)
6	F	0.40	0/856	0.65	0/1154
7	G	0.43	0/1276	0.62	0/1709
8	H	0.55	0/1136	0.81	0/1527
9	I	0.42	0/1029	0.68	1/1379 (0.1%)
10	J	0.41	0/806	0.68	1/1084 (0.1%)
11	K	0.46	0/900	0.72	0/1213
12	L	0.53	0/987	0.83	0/1322
13	M	0.42	0/1008	0.66	0/1347
14	N	0.47	0/501	0.68	0/664
15	O	0.46	0/745	0.62	0/992
16	P	0.56	0/717	0.77	0/965
17	Q	0.58	1/870 (0.1%)	0.74	0/1159
18	R	0.47	0/604	0.65	0/801
19	S	0.45	0/662	0.70	0/892
20	T	0.51	0/765	0.77	0/1007
21	U	0.58	0/213	0.73	0/279
22	X	0.54	0/116	0.73	0/179
23	Y	0.45	0/253	0.66	0/387
All	All	0.53	3/56278 (0.0%)	0.70	34/83516 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	2	17
12	L	0	1
All	All	2	18

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1531	A	O3'-P	24.47	1.90	1.61
4	D	9	CYS	CB-SG	5.49	1.91	1.82
17	Q	96	GLU	CB-CG	5.08	1.61	1.52

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1498	U	C2'-C3'-O3'	9.36	130.08	109.50
1	A	281	G	C2'-C3'-O3'	9.32	130.01	109.50
1	A	243	A	C2'-C3'-O3'	9.19	129.71	109.50
1	A	366	C	C2'-C3'-O3'	8.66	128.54	109.50
1	A	812	C	C2'-C3'-O3'	8.32	127.80	109.50

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	243	A	C3'
1	A	281	G	C3'

5 of 18 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	250	A	Sidechain
1	A	251	G	Sidechain
1	A	410	G	Sidechain
1	A	528	C	Sidechain
1	A	530	G	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	32489	0	16401	841	0
2	B	1901	0	1951	227	0
3	C	1613	0	1677	181	0
4	D	1703	0	1765	134	0
5	E	1147	0	1207	115	0
6	F	843	0	857	78	0
7	G	1257	0	1296	72	0
8	H	1116	0	1177	82	0
9	I	1010	0	1037	87	0
10	J	793	0	835	138	0
11	K	885	0	904	62	0
12	L	971	0	1057	92	0
13	M	997	0	1072	99	0
14	N	492	0	530	62	0
15	O	734	0	771	66	0
16	P	701	0	720	67	0
17	Q	857	0	928	57	0
18	R	598	0	670	47	0
19	S	648	0	673	60	0
20	T	763	0	861	74	0
21	U	209	0	221	20	0
22	X	104	0	55	3	0
23	Y	277	0	146	6	0
24	A	42	0	45	1	0
25	A	161	0	0	0	0
25	B	1	0	0	0	0
25	D	1	0	0	0	0
25	E	2	0	0	0	0
25	F	1	0	0	0	0
25	G	1	0	0	0	0
25	H	2	0	0	0	0
25	Q	1	0	0	0	0
26	A	21	0	0	0	0
26	E	1	0	0	0	0
27	D	1	0	0	1	0
27	N	1	0	0	2	0
All	All	52344	0	36856	2462	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 2462 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:14:ILE:HG22	3:C:15:THR:H	1.09	1.13
1:A:243:A:H4'	1:A:244:U:H5'	1.34	1.09
3:C:50:ALA:HB1	3:C:70:VAL:HG11	1.32	1.09
1:A:975:A:H4'	1:A:976:G:H5''	1.38	1.05
10:J:32:ALA:HB1	10:J:75:ILE:HG13	1.37	1.04

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	233/256 (91%)	157 (67%)	55 (24%)	21 (9%)	1	1
3	C	205/239 (86%)	151 (74%)	41 (20%)	13 (6%)	1	4
4	D	206/209 (99%)	170 (82%)	34 (16%)	2 (1%)	15	45
5	E	149/162 (92%)	134 (90%)	14 (9%)	1 (1%)	22	54
6	F	99/101 (98%)	84 (85%)	12 (12%)	3 (3%)	4	17
7	G	153/156 (98%)	121 (79%)	29 (19%)	3 (2%)	7	27
8	H	136/138 (99%)	119 (88%)	12 (9%)	5 (4%)	3	13
9	I	125/128 (98%)	97 (78%)	19 (15%)	9 (7%)	1	3
10	J	97/105 (92%)	73 (75%)	13 (13%)	11 (11%)	0	1
11	K	117/129 (91%)	95 (81%)	18 (15%)	4 (3%)	3	15
12	L	123/135 (91%)	96 (78%)	17 (14%)	10 (8%)	1	2
13	M	123/126 (98%)	88 (72%)	23 (19%)	12 (10%)	0	1
14	N	58/61 (95%)	47 (81%)	5 (9%)	6 (10%)	0	1
15	O	86/89 (97%)	67 (78%)	17 (20%)	2 (2%)	6	23
16	P	82/88 (93%)	73 (89%)	8 (10%)	1 (1%)	13	40
17	Q	102/105 (97%)	88 (86%)	8 (8%)	6 (6%)	1	5

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
18	R	71/88 (81%)	63 (89%)	6 (8%)	2 (3%)	5	19
19	S	79/93 (85%)	58 (73%)	11 (14%)	10 (13%)	0	0
20	T	97/106 (92%)	76 (78%)	15 (16%)	6 (6%)	1	4
21	U	23/27 (85%)	14 (61%)	8 (35%)	1 (4%)	2	10
All	All	2364/2541 (93%)	1871 (79%)	365 (15%)	128 (5%)	2	6

5 of 128 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	15	VAL
2	B	17	PHE
2	B	74	LYS
2	B	207	ALA
3	C	15	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	202/220 (92%)	183 (91%)	19 (9%)	8	26
3	C	160/188 (85%)	145 (91%)	15 (9%)	8	26
4	D	180/181 (99%)	167 (93%)	13 (7%)	14	39
5	E	115/123 (94%)	101 (88%)	14 (12%)	5	15
6	F	90/90 (100%)	86 (96%)	4 (4%)	28	61
7	G	126/127 (99%)	117 (93%)	9 (7%)	14	40
8	H	119/119 (100%)	109 (92%)	10 (8%)	11	31
9	I	98/99 (99%)	90 (92%)	8 (8%)	11	32
10	J	87/92 (95%)	83 (95%)	4 (5%)	27	60
11	K	90/99 (91%)	81 (90%)	9 (10%)	7	23
12	L	104/111 (94%)	94 (90%)	10 (10%)	8	25
13	M	100/101 (99%)	92 (92%)	8 (8%)	12	33

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
14	N	49/50 (98%)	42 (86%)	7 (14%)	3	10
15	O	79/80 (99%)	72 (91%)	7 (9%)	9	29
16	P	72/74 (97%)	67 (93%)	5 (7%)	15	41
17	Q	96/97 (99%)	90 (94%)	6 (6%)	18	46
18	R	64/77 (83%)	61 (95%)	3 (5%)	26	59
19	S	71/80 (89%)	70 (99%)	1 (1%)	67	89
20	T	76/82 (93%)	66 (87%)	10 (13%)	4	12
21	U	19/22 (86%)	19 (100%)	0	100	100
All	All	1997/2112 (95%)	1835 (92%)	162 (8%)	11	33

5 of 162 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
7	G	156	TRP
9	I	121	ARG
18	R	54	ARG
8	H	37	ARG
8	H	119	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 65 such sidechains are listed below:

Mol	Chain	Res	Type
7	G	37	ASN
9	I	23	ASN
18	R	36	ASN
7	G	56	GLN
7	G	106	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	1510/1522 (99%)	199 (13%)	58 (3%)
22	X	4/6 (66%)	1 (25%)	0
23	Y	10/17 (58%)	0	0
All	All	1524/1545 (98%)	200 (13%)	58 (3%)

5 of 200 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	8	A
1	A	9	G
1	A	31	G
1	A	32	A
1	A	39	G

5 of 58 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	560	U
1	A	812	C
1	A	1447	A
1	A	575	G
1	A	701	C

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
23	CM0	Y	34	23	16,26,27	1.74	4 (25%)	18,37,40	5.07	6 (33%)
23	6MZ	Y	37	23	18,25,26	0.85	1 (5%)	16,36,39	1.06	2 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
23	CM0	Y	34	23	-	3/8/30/31	0/2/2/2
23	6MZ	Y	37	23	-	0/5/27/28	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
23	Y	34	CM0	C4-N3	4.03	1.40	1.33
23	Y	34	CM0	C4-C5	3.46	1.48	1.40
23	Y	34	CM0	O5-C7	2.94	1.53	1.43
23	Y	34	CM0	O5-C5	-2.70	1.32	1.37
23	Y	37	6MZ	C8-N7	-2.10	1.31	1.34

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
23	Y	34	CM0	C4-N3-C2	15.80	128.48	115.14
23	Y	34	CM0	C7-O5-C5	9.65	136.75	117.76
23	Y	34	CM0	C5-C4-N3	-6.12	114.27	122.66
23	Y	34	CM0	C5-C6-N1	6.11	126.78	120.44
23	Y	34	CM0	O5-C7-C8	5.15	122.45	108.59

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
23	Y	34	CM0	C4-C5-O5-C7
23	Y	34	CM0	C8-C7-O5-C5
23	Y	34	CM0	C6-C5-O5-C7

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
23	Y	34	CM0	4	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 195 ligands modelled in this entry, 194 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
24	PAR	A	1601	-	45,45,45	1.56	7 (15%)	64,67,67	1.27	8 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
24	PAR	A	1601	-	-	2/18/94/94	0/4/4/4

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
24	A	1601	PAR	C64-C54	5.01	1.58	1.52
24	A	1601	PAR	C52-C42	3.07	1.58	1.52
24	A	1601	PAR	O54-C14	3.01	1.49	1.41
24	A	1601	PAR	C31-C21	2.70	1.56	1.53
24	A	1601	PAR	C11-C21	2.41	1.57	1.52

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
24	A	1601	PAR	O33-C14-C24	4.20	115.45	108.22
24	A	1601	PAR	O54-C54-C64	4.05	113.55	106.01
24	A	1601	PAR	C14-O54-C54	3.35	120.26	113.69
24	A	1601	PAR	O52-C13-C23	2.85	113.87	107.96
24	A	1601	PAR	O52-C13-O43	-2.51	108.71	111.43

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
24	A	1601	PAR	C23-C33-O33-C14
24	A	1601	PAR	C43-C33-O33-C14

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
24	A	1601	PAR	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	1533:C	O3'	1539:C	P	23.27

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1512/1522 (99%)	0.68	16 (1%) 80 80	30, 61, 130, 184	0
2	B	235/256 (91%)	0.74	21 (8%) 9 7	51, 85, 127, 143	0
3	C	207/239 (86%)	0.65	5 (2%) 59 56	53, 77, 110, 117	0
4	D	208/209 (99%)	0.45	6 (2%) 51 47	49, 67, 88, 96	0
5	E	151/162 (93%)	0.44	0 100 100	35, 51, 70, 88	0
6	F	101/101 (100%)	0.61	5 (4%) 28 25	60, 84, 92, 100	0
7	G	155/156 (99%)	0.46	5 (3%) 47 43	52, 72, 104, 116	0
8	H	138/138 (100%)	0.38	1 (0%) 87 87	32, 51, 66, 74	0
9	I	127/128 (99%)	0.57	9 (7%) 16 12	49, 83, 100, 106	0
10	J	99/105 (94%)	1.16	16 (16%) 1 1	50, 104, 140, 145	0
11	K	119/129 (92%)	0.58	4 (3%) 45 40	38, 59, 82, 100	0
12	L	125/135 (92%)	0.61	3 (2%) 59 56	23, 60, 79, 100	0
13	M	125/126 (99%)	0.84	16 (12%) 3 2	55, 73, 130, 159	0
14	N	60/61 (98%)	0.66	4 (6%) 17 13	56, 70, 91, 102	0
15	O	88/89 (98%)	0.42	2 (2%) 60 58	48, 66, 84, 103	0
16	P	84/88 (95%)	0.42	0 100 100	41, 51, 65, 93	0
17	Q	104/105 (99%)	0.72	5 (4%) 30 27	36, 55, 102, 132	0
18	R	73/88 (82%)	0.56	5 (6%) 17 13	54, 69, 105, 134	0
19	S	81/93 (87%)	0.69	10 (12%) 4 3	60, 87, 107, 116	0
20	T	99/106 (93%)	0.40	3 (3%) 50 45	39, 57, 81, 87	0
21	U	25/27 (92%)	0.29	0 100 100	46, 64, 85, 88	0
22	X	5/6 (83%)	0.54	0 100 100	61, 65, 98, 105	0
23	Y	11/17 (64%)	1.07	1 (9%) 9 6	69, 101, 135, 141	0
All	All	3932/4086 (96%)	0.62	137 (3%) 44 38	23, 66, 116, 184	0

The worst 5 of 137 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
17	Q	105	ALA	9.7
15	O	89	GLY	8.8
17	Q	104	LYS	8.7
11	K	128	ALA	7.6
19	S	3	ARG	7.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
23	CM0	Y	34	25/26	0.90	0.21	74,83,92,94	0
23	6MZ	Y	37	23/24	0.94	0.23	70,72,74,76	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
26	K	A	1769	1/1	0.37	0.23	113,113,113,113	0
26	K	A	1779	1/1	0.43	0.14	102,102,102,102	0
25	MG	A	1687	1/1	0.44	0.30	77,77,77,77	0
25	MG	A	1747	1/1	0.45	0.16	57,57,57,57	0
25	MG	A	1667	1/1	0.45	0.27	75,75,75,75	0
25	MG	A	1642	1/1	0.53	0.17	77,77,77,77	0
25	MG	A	1605	1/1	0.55	0.15	80,80,80,80	0
25	MG	A	1653	1/1	0.57	0.37	114,114,114,114	0
26	K	A	1771	1/1	0.59	0.19	108,108,108,108	0
25	MG	A	1603	1/1	0.59	0.26	73,73,73,73	0
26	K	A	1776	1/1	0.61	0.20	98,98,98,98	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
25	MG	A	1749	1/1	0.62	0.24	60,60,60,60	0
25	MG	A	1604	1/1	0.64	0.24	83,83,83,83	0
25	MG	A	1758	1/1	0.64	0.27	72,72,72,72	0
25	MG	A	1638	1/1	0.66	0.29	67,67,67,67	0
25	MG	A	1665	1/1	0.67	0.13	83,83,83,83	0
25	MG	A	1610	1/1	0.67	0.26	76,76,76,76	0
25	MG	A	1617	1/1	0.68	0.37	56,56,56,56	0
26	K	E	203	1/1	0.69	0.12	94,94,94,94	0
25	MG	A	1614	1/1	0.69	0.14	66,66,66,66	0
26	K	A	1773	1/1	0.69	0.21	87,87,87,87	0
25	MG	A	1732	1/1	0.71	0.24	66,66,66,66	0
25	MG	A	1750	1/1	0.71	0.18	55,55,55,55	0
25	MG	A	1741	1/1	0.72	0.17	46,46,46,46	0
25	MG	A	1782	1/1	0.72	0.25	45,45,45,45	0
25	MG	A	1650	1/1	0.74	0.26	73,73,73,73	0
26	K	A	1777	1/1	0.74	0.15	99,99,99,99	0
25	MG	A	1714	1/1	0.75	0.17	43,43,43,43	0
25	MG	A	1718	1/1	0.76	0.21	46,46,46,46	0
25	MG	A	1699	1/1	0.76	0.20	66,66,66,66	0
25	MG	A	1706	1/1	0.77	0.16	48,48,48,48	0
25	MG	A	1720	1/1	0.77	0.19	68,68,68,68	0
25	MG	A	1620	1/1	0.77	0.17	72,72,72,72	0
25	MG	A	1609	1/1	0.77	0.11	60,60,60,60	0
25	MG	G	201	1/1	0.78	0.38	77,77,77,77	0
26	K	A	1765	1/1	0.78	0.12	97,97,97,97	0
25	MG	A	1688	1/1	0.78	0.25	70,70,70,70	0
25	MG	A	1727	1/1	0.79	0.23	60,60,60,60	0
25	MG	A	1721	1/1	0.79	0.24	53,53,53,53	0
25	MG	H	202	1/1	0.80	0.45	63,63,63,63	0
25	MG	A	1679	1/1	0.80	0.32	43,43,43,43	0
25	MG	A	1615	1/1	0.80	0.20	74,74,74,74	0
25	MG	D	302	1/1	0.80	0.15	42,42,42,42	0
25	MG	A	1613	1/1	0.81	0.16	33,33,33,33	0
25	MG	A	1602	1/1	0.81	0.26	69,69,69,69	0
25	MG	A	1612	1/1	0.81	0.28	57,57,57,57	0
25	MG	F	201	1/1	0.81	0.14	68,68,68,68	0
25	MG	A	1633	1/1	0.81	0.37	42,42,42,42	0
25	MG	A	1686	1/1	0.81	0.25	85,85,85,85	0
26	K	A	1778	1/1	0.81	0.11	105,105,105,105	0
25	MG	A	1634	1/1	0.82	0.26	50,50,50,50	0
25	MG	A	1690	1/1	0.82	0.18	39,39,39,39	0
25	MG	A	1705	1/1	0.82	0.23	54,54,54,54	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
25	MG	A	1673	1/1	0.82	0.17	58,58,58,58	0
25	MG	A	1730	1/1	0.82	0.15	38,38,38,38	0
25	MG	A	1744	1/1	0.82	0.22	39,39,39,39	0
26	K	A	1772	1/1	0.82	0.09	90,90,90,90	0
25	MG	A	1626	1/1	0.82	0.24	52,52,52,52	0
25	MG	A	1702	1/1	0.83	0.10	45,45,45,45	0
25	MG	A	1740	1/1	0.83	0.11	49,49,49,49	0
25	MG	A	1674	1/1	0.83	0.39	61,61,61,61	0
25	MG	A	1616	1/1	0.84	0.14	73,73,73,73	0
25	MG	A	1717	1/1	0.84	0.13	46,46,46,46	0
25	MG	A	1651	1/1	0.84	0.37	57,57,57,57	0
26	K	A	1761	1/1	0.85	0.10	83,83,83,83	0
25	MG	A	1716	1/1	0.85	0.09	59,59,59,59	0
25	MG	A	1748	1/1	0.85	0.13	56,56,56,56	0
25	MG	A	1755	1/1	0.85	0.18	68,68,68,68	0
26	K	A	1767	1/1	0.85	0.14	102,102,102,102	0
25	MG	A	1756	1/1	0.85	0.21	79,79,79,79	0
25	MG	A	1652	1/1	0.85	0.30	57,57,57,57	0
25	MG	A	1643	1/1	0.86	0.15	28,28,28,28	0
25	MG	A	1640	1/1	0.86	0.35	46,46,46,46	0
25	MG	A	1724	1/1	0.86	0.23	54,54,54,54	0
25	MG	A	1632	1/1	0.86	0.23	34,34,34,34	0
25	MG	A	1715	1/1	0.86	0.21	51,51,51,51	0
25	MG	A	1742	1/1	0.87	0.11	55,55,55,55	0
25	MG	A	1669	1/1	0.87	0.27	55,55,55,55	0
25	MG	A	1708	1/1	0.87	0.21	33,33,33,33	0
26	K	A	1780	1/1	0.87	0.11	92,92,92,92	0
25	MG	A	1710	1/1	0.87	0.15	66,66,66,66	0
25	MG	A	1611	1/1	0.87	0.11	45,45,45,45	0
26	K	A	1763	1/1	0.87	0.08	88,88,88,88	0
25	MG	A	1641	1/1	0.88	0.26	64,64,64,64	0
25	MG	A	1654	1/1	0.88	0.33	73,73,73,73	0
25	MG	A	1704	1/1	0.88	0.19	69,69,69,69	0
25	MG	A	1735	1/1	0.89	0.18	23,23,23,23	0
25	MG	A	1703	1/1	0.89	0.15	45,45,45,45	0
25	MG	A	1757	1/1	0.89	0.21	74,74,74,74	0
25	MG	Q	201	1/1	0.89	0.17	71,71,71,71	0
26	K	A	1770	1/1	0.89	0.14	98,98,98,98	0
25	MG	A	1759	1/1	0.89	0.12	44,44,44,44	0
25	MG	A	1661	1/1	0.89	0.33	41,41,41,41	0
25	MG	A	1636	1/1	0.89	0.11	52,52,52,52	0
25	MG	A	1645	1/1	0.89	0.27	33,33,33,33	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
25	MG	A	1738	1/1	0.90	0.24	73,73,73,73	0
25	MG	H	201	1/1	0.90	0.34	62,62,62,62	0
26	K	A	1775	1/1	0.90	0.10	97,97,97,97	0
26	K	A	1764	1/1	0.90	0.21	88,88,88,88	0
26	K	A	1774	1/1	0.90	0.15	80,80,80,80	0
25	MG	A	1660	1/1	0.90	0.12	46,46,46,46	0
25	MG	A	1697	1/1	0.90	0.13	36,36,36,36	0
25	MG	A	1662	1/1	0.90	0.26	94,94,94,94	0
25	MG	A	1754	1/1	0.90	0.16	37,37,37,37	0
25	MG	A	1666	1/1	0.91	0.14	59,59,59,59	0
25	MG	A	1696	1/1	0.91	0.22	30,30,30,30	0
25	MG	A	1712	1/1	0.91	0.10	27,27,27,27	0
25	MG	A	1629	1/1	0.91	0.29	33,33,33,33	0
25	MG	A	1608	1/1	0.91	0.31	10,10,10,10	0
25	MG	A	1726	1/1	0.91	0.16	35,35,35,35	0
26	K	A	1766	1/1	0.91	0.15	88,88,88,88	0
26	K	A	1760	1/1	0.91	0.13	68,68,68,68	0
25	MG	A	1736	1/1	0.91	0.22	32,32,32,32	0
26	K	A	1768	1/1	0.91	0.08	88,88,88,88	0
25	MG	A	1606	1/1	0.91	0.38	38,38,38,38	0
25	MG	A	1621	1/1	0.91	0.34	70,70,70,70	0
25	MG	A	1625	1/1	0.91	0.31	70,70,70,70	0
25	MG	A	1646	1/1	0.91	0.27	55,55,55,55	0
25	MG	A	1623	1/1	0.91	0.37	32,32,32,32	0
25	MG	A	1713	1/1	0.92	0.22	28,28,28,28	0
25	MG	A	1656	1/1	0.92	0.34	63,63,63,63	0
25	MG	A	1635	1/1	0.92	0.23	60,60,60,60	0
25	MG	A	1745	1/1	0.92	0.19	72,72,72,72	0
25	MG	A	1725	1/1	0.92	0.14	43,43,43,43	0
25	MG	A	1685	1/1	0.92	0.25	51,51,51,51	0
25	MG	A	1670	1/1	0.92	0.12	34,34,34,34	0
25	MG	A	1678	1/1	0.92	0.17	34,34,34,34	0
25	MG	A	1694	1/1	0.92	0.12	35,35,35,35	0
25	MG	A	1619	1/1	0.92	0.15	57,57,57,57	0
25	MG	A	1701	1/1	0.92	0.13	35,35,35,35	0
25	MG	E	202	1/1	0.92	0.16	62,62,62,62	0
25	MG	A	1707	1/1	0.92	0.14	33,33,33,33	0
25	MG	A	1751	1/1	0.92	0.11	33,33,33,33	0
25	MG	A	1618	1/1	0.93	0.33	45,45,45,45	0
25	MG	A	1733	1/1	0.93	0.11	23,23,23,23	0
24	PAR	A	1601	42/42	0.93	0.32	47,52,74,78	0
25	MG	A	1689	1/1	0.93	0.28	64,64,64,64	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
25	MG	A	1719	1/1	0.93	0.15	26,26,26,26	0
25	MG	A	1753	1/1	0.93	0.15	51,51,51,51	0
25	MG	A	1691	1/1	0.93	0.35	46,46,46,46	0
25	MG	A	1743	1/1	0.93	0.13	55,55,55,55	0
25	MG	A	1683	1/1	0.93	0.35	45,45,45,45	0
25	MG	A	1698	1/1	0.93	0.22	36,36,36,36	0
25	MG	A	1728	1/1	0.93	0.16	64,64,64,64	0
25	MG	A	1695	1/1	0.93	0.27	42,42,42,42	0
25	MG	A	1723	1/1	0.93	0.41	48,48,48,48	0
25	MG	A	1746	1/1	0.94	0.17	15,15,15,15	0
25	MG	A	1677	1/1	0.94	0.20	23,23,23,23	0
25	MG	A	1647	1/1	0.94	0.24	54,54,54,54	0
25	MG	A	1644	1/1	0.94	0.36	62,62,62,62	0
25	MG	A	1682	1/1	0.94	0.22	26,26,26,26	0
25	MG	A	1627	1/1	0.94	0.32	35,35,35,35	0
25	MG	A	1722	1/1	0.94	0.12	41,41,41,41	0
25	MG	A	1676	1/1	0.94	0.22	37,37,37,37	0
25	MG	A	1752	1/1	0.94	0.25	62,62,62,62	0
25	MG	A	1711	1/1	0.95	0.42	75,75,75,75	0
25	MG	A	1668	1/1	0.95	0.20	47,47,47,47	0
25	MG	A	1658	1/1	0.95	0.15	73,73,73,73	0
25	MG	A	1700	1/1	0.95	0.14	57,57,57,57	0
25	MG	A	1692	1/1	0.95	0.17	31,31,31,31	0
25	MG	A	1681	1/1	0.95	0.21	41,41,41,41	0
25	MG	A	1639	1/1	0.95	0.32	2,2,2,2	0
25	MG	A	1648	1/1	0.95	0.25	56,56,56,56	0
25	MG	A	1693	1/1	0.96	0.28	31,31,31,31	0
25	MG	A	1684	1/1	0.96	0.17	16,16,16,16	0
25	MG	E	201	1/1	0.96	0.35	41,41,41,41	0
26	K	A	1762	1/1	0.96	0.10	74,74,74,74	0
25	MG	A	1637	1/1	0.96	0.15	35,35,35,35	0
25	MG	A	1663	1/1	0.96	0.19	38,38,38,38	0
25	MG	A	1628	1/1	0.96	0.38	32,32,32,32	0
25	MG	A	1709	1/1	0.96	0.11	44,44,44,44	0
25	MG	A	1680	1/1	0.96	0.31	35,35,35,35	0
25	MG	A	1664	1/1	0.96	0.18	42,42,42,42	0
25	MG	A	1672	1/1	0.96	0.20	37,37,37,37	0
25	MG	A	1729	1/1	0.97	0.24	33,33,33,33	0
25	MG	A	1607	1/1	0.97	0.23	36,36,36,36	0
25	MG	A	1624	1/1	0.97	0.39	37,37,37,37	0
25	MG	B	301	1/1	0.97	0.22	51,51,51,51	0
25	MG	A	1630	1/1	0.97	0.23	52,52,52,52	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
25	MG	A	1739	1/1	0.97	0.12	37,37,37,37	0
25	MG	A	1622	1/1	0.97	0.34	31,31,31,31	0
25	MG	A	1649	1/1	0.97	0.28	32,32,32,32	0
25	MG	A	1671	1/1	0.97	0.34	32,32,32,32	0
25	MG	A	1737	1/1	0.97	0.14	28,28,28,28	0
25	MG	A	1783	1/1	0.97	0.22	58,58,58,58	0
25	MG	A	1734	1/1	0.97	0.20	29,29,29,29	0
25	MG	A	1655	1/1	0.97	0.20	42,42,42,42	0
25	MG	A	1675	1/1	0.97	0.18	43,43,43,43	0
25	MG	A	1631	1/1	0.98	0.21	39,39,39,39	0
25	MG	A	1781	1/1	0.98	0.21	34,34,34,34	0
25	MG	A	1659	1/1	0.98	0.22	31,31,31,31	0
25	MG	A	1657	1/1	0.98	0.24	29,29,29,29	0
27	ZN	D	301	1/1	0.98	0.35	70,70,70,70	0
25	MG	A	1731	1/1	0.99	0.16	20,20,20,20	0
27	ZN	N	101	1/1	0.99	0.19	67,67,67,67	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around PAR A 1601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.