



wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 12:47 am BST

PDB ID : 2UUE
Title : REPLACE: A strategy for Iterative Design of Cyclin Binding Groove Inhibitors
Authors : Andrews, M.J.; Kontopidis, G.; McInnes, C.; Plater, A.; Innes, L.; Cowan, A.;
Jewsbury, P.; Fischer, P.M.
Deposited on : 2007-03-02
Resolution : 2.06 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2684 (2.08-2.04)
Clashscore	141614	2801 (2.08-2.04)
Ramachandran outliers	138981	2768 (2.08-2.04)
Sidechain outliers	138945	2768 (2.08-2.04)
RSRZ outliers	127900	2646 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	<div> <div>7%</div> <div>69%</div> <div>25%</div> <div>• •</div> </div>
1	C	298	<div> <div>7%</div> <div>69%</div> <div>23%</div> <div>6%</div> <div>•</div> </div>
2	B	259	<div> <div>3%</div> <div>75%</div> <div>20%</div> <div>•</div> </div>
2	D	259	<div> <div>5%</div> <div>75%</div> <div>20%</div> <div>•</div> </div>
3	E	5	<div> <div>60%</div> <div>20%</div> <div>20%</div> </div>
3	F	5	<div> <div>20%</div> <div>60%</div> <div>40%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MTZ	A	1297	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9332 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CELL DIVISION PROTEIN KINASE 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	296	Total	C	N	O	S	0	0	0
			2378	1547	403	420	8			
1	C	297	Total	C	N	O	S	0	0	1
			2379	1547	404	420	8			


- Molecule 2 is a protein called CYCLIN A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	258	Total	C	N	O	S	0	0	0
			2083	1350	339	383	11			
2	D	258	Total	C	N	O	S	0	0	0
			2084	1350	339	384	11			

- Molecule 3 is a protein called GVC-TETRAPEPTIDE INHIBITOR.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	5	Total	C	F	N	O	0	0	1
			40	27	1	8	4			
3	F	5	Total	C	F	N	O	0	0	1
			40	27	1	8	4			

- Molecule 4 is 4-METHYL-5-[(2E)-2-[(4-MORPHOLIN-4-YLPHENYL)IMINO]-2,5-DIHYDROPYRIMIDIN-4-YL]-1,3-THIAZOL-2-AMINE (three-letter code: MTZ) (formula: C₁₈H₂₀N₆OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			26	18	6	1	1		
4	C	1	Total	C	N	O	S	0	0
			26	18	6	1	1		

- Molecule 5 is 1-(3,5-DICHLOROPHENYL)-5-METHYL-1H-1,2,4-TRIAZOLE-3-CARBOXYLIC ACID (three-letter code: GVC) (formula: $C_{10}H_7Cl_2N_3O_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	E	1	Total	C	Cl	N	O	0	0
			16	10	2	3	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	F	1	Total	C	Cl	N	O	0	0
			16	10	2	3	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	49	Total	O	0	0
			49	49		
6	B	64	Total	O	0	0
			64	64		
6	C	65	Total	O	0	0
			65	65		
6	D	63	Total	O	0	0
			63	63		
6	E	2	Total	O	0	0
			2	2		
6	F	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CELL DIVISION PROTEIN KINASE 2




• Molecule 1: CELL DIVISION PROTEIN KINASE 2



• Molecule 2: CYCLIN A2



● Molecule 2: CYCLIN A2



● Molecule 3: GVC-TETRAPEPTIDE INHIBITOR



● Molecule 3: GVC-TETRAPEPTIDE INHIBITOR



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	74.56Å 113.80Å 155.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	91.29 – 2.06 19.90 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.8 (91.29-2.06) 99.0 (19.90-2.10)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 2.09Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.196 , 0.244 0.199 , 0.245	Depositor DCC
R_{free} test set	2299 reflections (2.99%)	wwPDB-VP
Wilson B-factor (Å ²)	36.3	Xtriage
Anisotropy	0.087	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 51.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9332	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MTZ, GVC, NH2, PFF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/2440	0.74	2/3313 (0.1%)
1	C	0.48	0/2441	0.77	1/3315 (0.0%)
2	B	0.48	0/2133	0.77	2/2896 (0.1%)
2	D	0.49	0/2134	0.76	2/2897 (0.1%)
3	E	1.54	1/26 (3.8%)	1.72	1/33 (3.0%)
3	F	1.50	1/26 (3.8%)	1.66	1/33 (3.0%)
All	All	0.49	2/9200 (0.0%)	0.77	9/12487 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	3
All	All	0	4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	1	ARG	CB-CG	-6.95	1.33	1.52
3	E	1	ARG	CB-CG	-6.94	1.33	1.52

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	250	ARG	NE-CZ-NH1	8.14	124.37	120.30
2	B	250	ARG	NE-CZ-NH2	-6.71	116.94	120.30
2	D	364	LEU	CA-CB-CG	6.27	129.72	115.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	55	LEU	CB-CG-CD1	6.14	121.44	111.00
1	C	50	ARG	NE-CZ-NH1	-5.97	117.31	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	14	THR	Peptide
1	C	162	GLU	Peptide
1	C	37	LEU	Peptide
1	C	70	ILE	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2378	0	2426	86	0
1	C	2379	0	2426	93	0
2	B	2083	0	2107	57	0
2	D	2084	0	2107	46	0
3	E	40	0	41	1	0
3	F	40	0	41	0	0
4	A	26	0	20	13	0
4	C	26	0	20	5	0
5	E	16	0	6	3	0
5	F	16	0	6	3	0
6	A	49	0	0	3	0
6	B	64	0	0	7	0
6	C	65	0	0	15	0
6	D	63	0	0	5	0
6	E	2	0	0	1	0
6	F	1	0	0	0	0
All	All	9332	0	9200	283	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 283 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:178:LYS:HA	6:A:2035:HOH:O	1.28	1.31
1:C:59:ASN:HB2	6:C:2014:HOH:O	1.49	1.11
5:E:1433:GVC:C29	5:E:1433:GVC:H361	1.73	1.11
1:A:42:GLU:OE2	2:B:275:VAL:HG23	1.50	1.11
5:E:1433:GVC:C36	5:E:1433:GVC:H29	1.80	1.09

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	294/298 (99%)	280 (95%)	12 (4%)	2 (1%)	22	11
1	C	295/298 (99%)	282 (96%)	10 (3%)	3 (1%)	15	6
2	B	256/259 (99%)	247 (96%)	8 (3%)	1 (0%)	34	25
2	D	256/259 (99%)	247 (96%)	8 (3%)	1 (0%)	34	25
3	E	2/5 (40%)	2 (100%)	0	0	100	100
3	F	2/5 (40%)	2 (100%)	0	0	100	100
All	All	1105/1124 (98%)	1060 (96%)	38 (3%)	7 (1%)	25	15

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	15	TYR
2	B	176	PRO
1	C	41	THR
2	D	431	ASN
1	A	160	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/263 (99%)	235 (90%)	26 (10%)	7	2
1	C	261/263 (99%)	225 (86%)	36 (14%)	3	1
2	B	232/233 (100%)	211 (91%)	21 (9%)	9	4
2	D	232/233 (100%)	211 (91%)	21 (9%)	9	4
3	E	3/3 (100%)	2 (67%)	1 (33%)	0	0
3	F	3/3 (100%)	2 (67%)	1 (33%)	0	0
All	All	992/998 (99%)	886 (89%)	106 (11%)	6	2

5 of 106 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	5	GLN
1	C	87	LEU
2	D	378	ARG
1	C	15	TYR
1	C	39	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	431	ASN
1	C	60	HIS
2	D	208	ASN
2	B	404	HIS
2	D	322	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	PFF	F	4	3	11,12,13	0.73	0	12,15,17	1.50	2 (16%)
3	PFF	E	4	3	11,12,13	0.82	0	12,15,17	1.38	2 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PFF	F	4	3	-	0/5/6/8	0/1/1/1
3	PFF	E	4	3	-	2/5/6/8	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	F	4	PFF	CD1-CE1-CZ	3.20	121.67	118.36
3	F	4	PFF	CE2-CZ-CE1	-2.96	118.89	122.83
3	E	4	PFF	CG-CB-CA	2.62	119.40	114.10
3	E	4	PFF	CD2-CG-CD1	2.01	121.33	118.17

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	4	PFF	O-C-CA-CB
3	E	4	PFF	C-CA-CB-CG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GVC	F	1433	3	15,17,18	1.41	2 (13%)	15,24,26	2.61	5 (33%)
5	GVC	E	1433	3	15,17,18	1.62	3 (20%)	15,24,26	2.62	7 (46%)
4	MTZ	A	1297	-	21,29,29	1.79	2 (9%)	23,40,40	1.34	3 (13%)
4	MTZ	C	1298	-	21,29,29	2.03	4 (19%)	23,40,40	1.87	7 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GVC	F	1433	3	-	4/4/6/8	0/2/2/2
5	GVC	E	1433	3	-	4/4/6/8	0/2/2/2
4	MTZ	A	1297	-	-	3/8/30/30	0/3/4/4
4	MTZ	C	1298	-	-	0/8/30/30	0/3/4/4

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	1298	MTZ	C3-C4	-6.34	1.38	1.47
4	A	1297	MTZ	C3-C4	-5.18	1.40	1.47
4	C	1298	MTZ	C8-N4	-5.03	1.34	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1297	MTZ	C8-N4	-4.94	1.34	1.42
5	E	1433	GVC	C28-CL1	3.67	1.82	1.74

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	1433	GVC	C29-C30-C31	-5.41	114.64	121.61
5	F	1433	GVC	N10-C35-N9	-4.80	110.74	114.72
5	E	1433	GVC	C29-C28-CL1	4.22	124.42	119.15
4	C	1298	MTZ	C6-N3-C5	4.13	123.46	116.03
5	E	1433	GVC	N10-C35-N9	-3.97	111.43	114.72

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	F	1433	GVC	C31-C30-N8-C34
5	F	1433	GVC	C29-C30-N8-C34
5	E	1433	GVC	C31-C30-N8-C34
5	E	1433	GVC	C29-C30-N8-C34
4	A	1297	MTZ	N2-C5-N4-C8


There are no ring outliers.

4 monomers are involved in 24 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	F	1433	GVC	3	0
5	E	1433	GVC	3	0
4	A	1297	MTZ	13	0
4	C	1298	MTZ	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient


equivalents in the CSD to analyse the geometry.




Ligand GVC E 1433



Ligand MTZ A 1297





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	296/298 (99%)	0.37	22 (7%) 14 15	36, 52, 90, 132	0
1	C	297/298 (99%)	0.26	20 (6%) 17 18	33, 48, 91, 129	0
2	B	258/259 (99%)	0.03	9 (3%) 44 46	35, 48, 71, 108	0
2	D	258/259 (99%)	0.08	12 (4%) 31 32	34, 50, 72, 104	0
3	E	3/5 (60%)	0.32	0 100 100	49, 49, 65, 66	0
3	F	3/5 (60%)	0.56	1 (33%) 0 0	46, 46, 62, 75	0
All	All	1115/1124 (99%)	0.20	64 (5%) 23 24	33, 50, 83, 132	0

The worst 5 of 64 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	15	TYR	13.1
1	C	39	THR	10.9
1	C	14	THR	10.5
1	A	13	GLY	10.3
2	B	175	VAL	8.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PFF	F	4	12/13	0.87	0.12	38,46,58,59	0
3	PFF	E	4	12/13	0.93	0.12	43,49,65,80	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	MTZ	A	1297	26/26	0.65	0.31	88,110,127,129	0
5	GVC	E	1433	16/17	0.75	0.20	52,59,69,79	0
5	GVC	F	1433	16/17	0.75	0.26	65,70,78,80	0
4	MTZ	C	1298	26/26	0.85	0.17	44,68,92,95	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around MTZ A 1297:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around GVC E 1433:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GVC F 1433:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.