



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 24, 2020 – 04:48 am BST

PDB ID : 4UUV  
Title : Structure of the DNA binding ETS domain of human ETV4 in complex with DNA  
Authors : Newman, J.A.; Cooper, C.D.O.; Kopec, J.; von Delft, F.; Arrowsmith, C.H.; Edwards, A.M.; Bountra, C.; Gileadi, O.  
Deposited on : 2014-07-31  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	100	<div> <div>2%</div> <div> <div></div> <div>89%</div> <div>7%</div> <div>•</div> </div> </div>
1	D	100	<div> <div>%</div> <div> <div></div> <div>87%</div> <div>8%</div> <div>5%</div> </div> </div>
1	G	100	<div> <div>%</div> <div> <div></div> <div>84%</div> <div>11%</div> <div>5%</div> </div> </div>
1	J	100	<div> <div>2%</div> <div> <div></div> <div>84%</div> <div>10%</div> <div>6%</div> </div> </div>
1	M	100	<div> <div></div> <div> <div></div> <div>86%</div> <div>8%</div> <div>6%</div> </div> </div>
1	P	100	<div> <div>%</div> <div> <div></div> <div>86%</div> <div>9%</div> <div>• •</div> </div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	S	100	
1	V	100	
2	B	10	
2	E	10	
2	H	10	
2	K	10	
2	N	10	
2	Q	10	
2	T	10	
2	W	10	
3	C	10	
3	F	10	
3	I	10	
3	L	10	
3	O	10	
3	R	10	
3	U	10	
4	X	10	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9352 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ETS TRANSLOCATION VARIANT 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	96	Total	C	N	O	S	0	0	0
			794	519	132	138	5			
1	D	95	Total	C	N	O	S	0	0	0
			776	509	127	135	5			
1	G	95	Total	C	N	O	S	0	0	0
			779	512	125	137	5			
1	J	94	Total	C	N	O	S	0	0	0
			785	514	130	136	5			
1	M	94	Total	C	N	O	S	0	0	0
			779	511	127	136	5			
1	P	96	Total	C	N	O	S	0	0	0
			794	519	132	138	5			
1	S	95	Total	C	N	O	S	0	0	0
			787	516	130	136	5			
1	V	94	Total	C	N	O	S	0	0	0
			755	495	123	132	5			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	336	SER	-	expression tag	UNP P43268
A	337	MET	-	expression tag	UNP P43268
D	336	SER	-	expression tag	UNP P43268
D	337	MET	-	expression tag	UNP P43268
G	336	SER	-	expression tag	UNP P43268
G	337	MET	-	expression tag	UNP P43268
J	336	SER	-	expression tag	UNP P43268
J	337	MET	-	expression tag	UNP P43268
M	336	SER	-	expression tag	UNP P43268
M	337	MET	-	expression tag	UNP P43268
P	336	SER	-	expression tag	UNP P43268
P	337	MET	-	expression tag	UNP P43268
S	336	SER	-	expression tag	UNP P43268

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
S	337	MET	-	expression tag	UNP P43268
V	336	SER	-	expression tag	UNP P43268
V	337	MET	-	expression tag	UNP P43268

- Molecule 2 is a DNA chain called 5'-D(\*AP\*CP\*CP\*GP\*GP\*AP\*AP\*GP\*TP\*GP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	10	Total	C	N	O	P	0	0	0
			206	98	43	56	9			
2	E	10	Total	C	N	O	P	0	0	0
			188	88	38	53	9			
2	H	10	Total	C	N	O	P	0	0	0
			206	98	43	56	9			
2	K	10	Total	C	N	O	P	0	0	0
			188	88	38	53	9			
2	N	10	Total	C	N	O	P	0	0	0
			188	88	38	53	9			
2	Q	10	Total	C	N	O	P	0	0	0
			206	98	43	56	9			
2	T	10	Total	C	N	O	P	0	0	0
			206	98	43	56	9			
2	W	10	Total	C	N	O	P	0	0	0
			206	98	43	56	9			

- Molecule 3 is a DNA chain called 5'-D(\*AP\*CP\*TP\*TP\*CP\*CP\*GP\*GP\*TP\*CP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	10	Total	C	N	O	P	0	0	0
			183	87	30	57	9			
3	F	10	Total	C	N	O	P	0	0	0
			198	96	33	60	9			
3	I	10	Total	C	N	O	P	0	0	0
			183	87	30	57	9			
3	L	10	Total	C	N	O	P	0	0	0
			183	87	30	57	9			
3	O	10	Total	C	N	O	P	0	0	0
			183	87	30	57	9			
3	R	10	Total	C	N	O	P	0	0	0
			198	96	33	60	9			
3	U	10	Total	C	N	O	P	0	0	0
			198	96	33	60	9			

- Molecule 4 is a DNA chain called 5'-D(\*AP\*CP\*TP\*TP\*CP\*CP\*GP\*GP\*TP\*CP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	X	10	Total	C	N	O	P	0	0	0
			183	87	30	57	9			


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ETS TRANSLOCATION VARIANT 4





- Molecule 1: ETS TRANSLOCATION VARIANT 4



- Molecule 1: ETS TRANSLOCATION VARIANT 4




- Molecule 1: ETS TRANSLOCATION VARIANT 4




- Molecule 1: ETS TRANSLOCATION VARIANT 4




- Molecule 1: ETS TRANSLOCATION VARIANT 4



- Molecule 1: ETS TRANSLOCATION VARIANT 4



- Molecule 1: ETS TRANSLOCATION VARIANT 4




- Molecule 2: 5'-D(\*AP\*CP\*CP\*GP\*GP\*AP\*AP\*GP\*TP\*GP)-3'



There are no outlier residues recorded for this chain.

- Molecule 2: 5'-D(\*AP\*CP\*CP\*GP\*GP\*AP\*AP\*GP\*TP\*GP)-3'



- Molecule 2: 5'-D(\*AP\*CP\*CP\*GP\*GP\*AP\*AP\*GP\*TP\*GP)-3'



- Molecule 2: 5'-D(\*AP\*CP\*CP\*GP\*GP\*AP\*AP\*GP\*TP\*GP)-3'



- Molecule 2: 5'-D(\*AP\*CP\*CP\*GP\*GP\*AP\*AP\*GP\*TP\*GP)-3'





There are no outlier residues recorded for this chain.


- Molecule 2: 5'-D(\*AP\*CP\*CP\*GP\*GP\*AP\*AP\*GP\*TP\*GP)-3'

Chain Q:  80% 20%



- Molecule 2: 5'-D(\*AP\*CP\*CP\*GP\*GP\*AP\*AP\*GP\*TP\*GP)-3'

Chain T:  10% 80% 20%



- Molecule 2: 5'-D(\*AP\*CP\*CP\*GP\*GP\*AP\*AP\*GP\*TP\*GP)-3'

Chain W:  90% 10%



- Molecule 3: 5'-D(\*AP\*CP\*TP\*TP\*CP\*CP\*GP\*GP\*TP\*CP)-3'

Chain C:  80% 20%



- Molecule 3: 5'-D(\*AP\*CP\*TP\*TP\*CP\*CP\*GP\*GP\*TP\*CP)-3'

Chain F:  100%

There are no outlier residues recorded for this chain.

- Molecule 3: 5'-D(\*AP\*CP\*TP\*TP\*CP\*CP\*GP\*GP\*TP\*CP)-3'

Chain I:  100%

There are no outlier residues recorded for this chain.

- Molecule 3: 5'-D(\*AP\*CP\*TP\*TP\*CP\*CP\*GP\*GP\*TP\*CP)-3'

Chain L:  90% 10%



- Molecule 3: 5'-D(\*AP\*CP\*TP\*TP\*CP\*CP\*GP\*GP\*TP\*CP)-3'

Chain O:  100%

There are no outlier residues recorded for this chain.

- Molecule 3: 5'-D(\*AP\*CP\*TP\*TP\*CP\*CP\*GP\*GP\*TP\*CP)-3'

Chain R:  100%


There are no outlier residues recorded for this chain.

- Molecule 3: 5'-D(\*AP\*CP\*TP\*TP\*CP\*CP\*GP\*GP\*TP\*CP)-3'

Chain U:  100%

There are no outlier residues recorded for this chain.

- Molecule 4: 5'-D(\*AP\*CP\*TP\*TP\*CP\*CP\*GP\*GP\*TP\*CP)-3'

Chain X:  10% 80% 20%



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	176.65Å 46.13Å 171.15Å 90.00° 96.69° 90.00°	Depositor
Resolution (Å)	40.97 – 2.80 43.72 – 2.80	Depositor EDS
% Data completeness (in resolution range)	94.6 (40.97-2.80) 94.6 (43.72-2.80)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.52 (at 2.81Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.203 , 0.248 0.205 , 0.249	Depositor DCC
$R_{free}$ test set	1534 reflections (4.69%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	80.0	Xtriage
Anisotropy	0.313	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 46.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9352	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	80.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.82% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.21	0/817	0.35	0/1104
1	D	0.21	0/799	0.34	0/1083
1	G	0.22	0/802	0.35	0/1086
1	J	0.21	0/808	0.35	0/1092
1	M	0.22	0/802	0.34	0/1085
1	P	0.21	0/817	0.34	0/1104
1	S	0.21	0/810	0.35	0/1095
1	V	0.21	0/778	0.34	0/1059
2	B	0.53	0/232	0.79	0/357
2	E	0.48	0/211	0.78	0/325
2	H	0.56	0/232	0.88	2/357 (0.6%)
2	K	0.52	0/211	0.80	0/325
2	N	0.45	0/211	0.78	0/325
2	Q	0.49	0/232	0.76	0/357
2	T	0.50	0/232	0.77	0/357
2	W	0.51	0/232	0.76	0/357
3	C	0.51	0/203	0.94	0/312
3	F	0.53	0/220	0.93	0/337
3	I	0.51	0/203	0.95	0/312
3	L	0.52	0/203	0.96	0/312
3	O	0.53	0/203	0.94	0/312
3	R	0.53	0/220	0.94	0/337
3	U	0.49	0/220	0.92	0/337
4	X	0.56	0/203	0.90	0/312
All	All	0.35	0/9901	0.60	2/14039 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	9	DT	OP1-P-O3'	5.67	117.67	105.20
2	H	9	DT	P-O3'-C3'	5.64	126.46	119.70

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	794	0	778	6	0
1	D	776	0	749	6	0
1	G	779	0	755	8	0
1	J	785	0	770	7	0
1	M	779	0	759	5	0
1	P	794	0	778	7	0
1	S	787	0	771	3	0
1	V	755	0	707	6	0
2	B	206	0	113	0	0
2	E	188	0	101	1	0
2	H	206	0	113	1	0
2	K	188	0	101	1	0
2	N	188	0	101	0	0
2	Q	206	0	113	2	0
2	T	206	0	113	1	0
2	W	206	0	113	1	0
3	C	183	0	103	2	0
3	F	198	0	115	0	0
3	I	183	0	103	0	0
3	L	183	0	103	1	0
3	O	183	0	103	0	0
3	R	198	0	115	0	0
3	U	198	0	115	0	0
4	X	183	0	103	1	0
All	All	9352	0	7795	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 45 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:Q:8:DG:H2''	2:Q:9:DT:H5'	1.73	0.69
1:G:367:MET:HG2	1:G:420:PHE:HB2	1.77	0.65
1:S:359:PHE:HB2	1:S:373:GLU:HB2	1.82	0.61
1:P:425:ASP:N	1:P:425:ASP:OD1	2.36	0.58
1:V:425:ASP:OD1	1:V:425:ASP:N	2.38	0.56

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	94/100 (94%)	88 (94%)	6 (6%)	0	100	100
1	D	93/100 (93%)	86 (92%)	7 (8%)	0	100	100
1	G	93/100 (93%)	88 (95%)	5 (5%)	0	100	100
1	J	92/100 (92%)	88 (96%)	4 (4%)	0	100	100
1	M	92/100 (92%)	88 (96%)	4 (4%)	0	100	100
1	P	94/100 (94%)	89 (95%)	5 (5%)	0	100	100
1	S	93/100 (93%)	91 (98%)	2 (2%)	0	100	100
1	V	92/100 (92%)	87 (95%)	5 (5%)	0	100	100
All	All	743/800 (93%)	705 (95%)	38 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	81/85 (95%)	81 (100%)	0	100	100
1	D	78/85 (92%)	78 (100%)	0	100	100
1	G	79/85 (93%)	79 (100%)	0	100	100
1	J	81/85 (95%)	80 (99%)	1 (1%)	71	92
1	M	80/85 (94%)	79 (99%)	1 (1%)	69	91
1	P	81/85 (95%)	80 (99%)	1 (1%)	71	92
1	S	80/85 (94%)	80 (100%)	0	100	100
1	V	74/85 (87%)	73 (99%)	1 (1%)	67	90
All	All	634/680 (93%)	630 (99%)	4 (1%)	86	96

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	425	ASP
1	M	367	MET
1	P	425	ASP
1	V	425	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	342	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	96/100 (96%)	0.23	2 (2%) 63 54	55, 66, 88, 98	0
1	D	95/100 (95%)	0.26	1 (1%) 80 75	53, 75, 100, 113	0
1	G	95/100 (95%)	0.21	1 (1%) 80 75	48, 63, 81, 92	0
1	J	94/100 (94%)	0.36	2 (2%) 63 54	56, 72, 91, 106	0
1	M	94/100 (94%)	0.08	0 100 100	53, 67, 87, 98	0
1	P	96/100 (96%)	0.43	1 (1%) 82 77	61, 76, 99, 106	0
1	S	95/100 (95%)	0.25	1 (1%) 80 75	60, 80, 96, 104	0
1	V	94/100 (94%)	0.43	1 (1%) 80 75	72, 86, 104, 121	0
2	B	10/10 (100%)	-0.16	0 100 100	63, 90, 103, 118	0
2	E	10/10 (100%)	-0.23	0 100 100	68, 82, 108, 132	0
2	H	10/10 (100%)	-0.08	0 100 100	57, 83, 106, 110	0
2	K	10/10 (100%)	0.43	0 100 100	60, 90, 112, 135	0
2	N	10/10 (100%)	-0.29	0 100 100	64, 82, 114, 120	0
2	Q	10/10 (100%)	-0.34	0 100 100	73, 87, 90, 94	0
2	T	10/10 (100%)	0.32	1 (10%) 7 4	82, 96, 116, 128	0
2	W	10/10 (100%)	-0.30	0 100 100	96, 119, 130, 131	0
3	C	10/10 (100%)	-0.34	0 100 100	65, 82, 94, 109	0
3	F	10/10 (100%)	-0.30	0 100 100	69, 83, 88, 88	0
3	I	10/10 (100%)	-0.04	0 100 100	61, 75, 89, 115	0
3	L	10/10 (100%)	-0.03	0 100 100	69, 90, 100, 110	0
3	O	10/10 (100%)	-0.08	0 100 100	66, 78, 99, 127	0
3	R	10/10 (100%)	-0.41	0 100 100	72, 93, 100, 102	0
3	U	10/10 (100%)	-0.15	0 100 100	75, 98, 102, 119	0
4	X	10/10 (100%)	0.67	1 (10%) 7 4	106, 124, 139, 153	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
All	All	919/960 (95%)	0.22	11 (1%) 79 73	48, 76, 106, 153	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	T	10	DG	3.9
4	X	10	DG	3.1
1	A	430	MET	2.6
1	V	347	LEU	2.4
1	S	365	ARG	2.4

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.