



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 17, 2020 – 11:15 AM EDT

PDB ID : 6UUM
Title : Crystal structure of antibody 438-B11 DSS mutant (Cys98A-Cys100aA)
Authors : Kumar, S.; Wilson, I.A.
Deposited on : 2019-10-30
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.4
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	215	 2% 93% 7%
1	F	215	 % 94% 6%
2	B	235	 2% 90% 7% •
2	H	235	 2% 90% 7% •

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7498 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called B11 Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	F	215	Total	C	N	O	S	0	0	0
			1649	1030	283	331	5			
1	A	215	Total	C	N	O	S	0	0	0
			1649	1030	283	331	5			

- Molecule 2 is a protein called B11 DSS Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	227	Total	C	N	O	S	0	0	0
			1717	1089	292	330	6			
2	B	227	Total	C	N	O	S	0	0	0
			1711	1086	292	327	6			

- Molecule 3 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	F	1	Total	C	O	0	0
			13	8	5		
3	F	1	Total	C	O	0	0
			5	3	2		
3	F	1	Total	C	O	0	0
			13	8	5		
3	H	1	Total	C	O	0	0
			13	8	5		
3	H	1	Total	C	O	0	0
			13	8	5		
3	H	1	Total	C	O	0	0
			9	6	3		
3	H	1	Total	C	O	0	0
			13	8	5		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			13	8	5		
3	A	1	Total	C	O	0	0
			4	3	1		
3	B	1	Total	C	O	0	0
			13	8	5		
3	B	1	Total	C	O	0	0
			13	8	5		
3	B	1	Total	C	O	0	0
			10	7	3		

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	F	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	F	1	Total	O	S	0	0
			5	4	1		
5	F	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	F	1	Total	O	S	0	0
			5	4	1		
5	F	1	Total	O	S	0	0
			5	4	1		
5	F	1	Total	O	S	0	0
			5	4	1		
5	F	1	Total	O	S	0	0
			5	4	1		
5	F	1	Total	O	S	0	0
			5	4	1		
5	H	1	Total	O	S	0	0
			5	4	1		
5	H	1	Total	O	S	0	0
			5	4	1		
5	H	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	F	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	F	133	Total	O	0	0
			133	133		
7	H	150	Total	O	0	0
			150	150		
7	A	116	Total	O	0	0
			116	116		
7	B	131	Total	O	0	0
			131	131		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: B11 Fab Light Chain





- Molecule 1: B11 Fab Light Chain



- Molecule 2: B11 DSS Fab Heavy Chain



- Molecule 2: B11 DSS Fab Heavy Chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	104.15Å 66.20Å 108.86Å 90.00° 117.96° 90.00°	Depositor
Resolution (Å)	48.08 – 2.10 48.08 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.6 (48.08-2.10) 99.6 (48.08-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.17	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.53 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.229 , 0.269 0.229 , 0.268	Depositor DCC
R_{free} test set	2008 reflections (2.61%)	wwPDB-VP
Wilson B-factor (Å ²)	28.5	Xtriage
Anisotropy	0.394	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 31.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.021 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7498	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, PG4, NAG, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/1684	0.49	0/2287
1	F	0.27	0/1684	0.48	0/2287
2	B	0.28	0/1755	0.46	0/2390
2	H	0.26	0/1761	0.46	0/2399
All	All	0.27	0/6884	0.47	0/9363

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1649	0	1595	11	0
1	F	1649	0	1597	8	0
2	B	1711	0	1692	10	0
2	H	1717	0	1695	12	0
3	A	21	0	25	1	0
3	B	36	0	46	1	0
3	F	31	0	40	0	0
3	H	48	0	65	4	0
4	A	14	0	13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	F	14	0	13	0	0
5	B	20	0	0	1	0
5	F	35	0	0	1	0
5	H	15	0	0	0	0
6	B	4	0	3	0	0
6	F	4	0	3	0	0
7	A	116	0	0	2	0
7	B	131	0	0	2	0
7	F	133	0	0	2	0
7	H	150	0	0	2	0
All	All	7498	0	6787	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 40 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:THR:CG2	7:A:606:HOH:O	2.32	0.78
1:A:10:THR:HG22	7:A:606:HOH:O	1.92	0.69
2:H:143:LYS:NZ	2:H:171:GLN:OE1	2.30	0.65
2:H:170:LEU:HB3	3:H:304:PG4:H11	1.82	0.62
2:B:68:ARG:NH2	7:B:402:HOH:O	2.34	0.60

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/215 (99%)	206 (97%)	7 (3%)	0	100	100
1	F	213/215 (99%)	205 (96%)	8 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	223/235 (95%)	218 (98%)	5 (2%)	0	100	100
2	H	223/235 (95%)	218 (98%)	5 (2%)	0	100	100
All	All	872/900 (97%)	847 (97%)	25 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/185 (100%)	184 (100%)	1 (0%)	88	92
1	F	185/185 (100%)	185 (100%)	0	100	100
2	B	192/199 (96%)	190 (99%)	2 (1%)	76	82
2	H	194/199 (98%)	193 (100%)	1 (0%)	88	92
All	All	756/768 (98%)	752 (100%)	4 (0%)	88	92

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	100(H)	TYR
1	A	10	THR
2	B	100(F)	SER
2	B	100(H)	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	F	189	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

31 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SO4	B	306	-	4,4,4	0.14	0	6,6,6	0.05	0
5	SO4	B	305	-	4,4,4	0.14	0	6,6,6	0.07	0
5	SO4	H	307	-	4,4,4	0.13	0	6,6,6	0.06	0
5	SO4	F	310	-	4,4,4	0.14	0	6,6,6	0.05	0
6	ACT	F	312	-	1,3,3	6.39	1 (100%)	0,3,3	0.00	-
6	ACT	B	308	-	1,3,3	6.29	1 (100%)	0,3,3	0.00	-
3	PG4	F	304	-	12,12,12	0.53	0	11,11,11	0.24	0
5	SO4	H	305	-	4,4,4	0.14	0	6,6,6	0.04	0
3	PG4	B	301	-	12,12,12	0.52	0	11,11,11	0.22	0
5	SO4	F	311	-	4,4,4	0.14	0	6,6,6	0.08	0
5	SO4	B	304	-	4,4,4	0.14	0	6,6,6	0.05	0
3	PG4	B	303	-	9,9,12	0.53	0	8,8,11	0.21	0
4	NAG	F	303	1	14,14,15	0.26	0	17,19,21	0.50	0
5	SO4	F	306	-	4,4,4	0.14	0	6,6,6	0.06	0
4	NAG	A	404	1	14,14,15	0.24	0	17,19,21	0.52	0
5	SO4	F	309	-	4,4,4	0.14	0	6,6,6	0.05	0
3	PG4	H	302	-	12,12,12	0.53	0	11,11,11	0.23	0
3	PG4	B	302	-	12,12,12	0.52	0	11,11,11	0.20	0
3	PG4	H	303	-	8,8,12	0.54	0	7,7,11	0.38	0
3	PG4	H	304	-	12,12,12	0.53	0	11,11,11	0.26	0
5	SO4	F	307	-	4,4,4	0.14	0	6,6,6	0.05	0
3	PG4	F	302	-	4,4,12	0.42	0	3,3,11	0.33	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PG4	A	402	-	12,12,12	0.53	0	11,11,11	0.24	0
5	SO4	F	308	-	4,4,4	0.15	0	6,6,6	0.07	0
5	SO4	F	305	-	4,4,4	0.15	0	6,6,6	0.10	0
3	PG4	H	301	-	12,12,12	0.53	0	11,11,11	0.27	0
3	PG4	A	403	-	3,3,12	0.53	0	2,2,11	0.30	0
5	SO4	H	306	-	4,4,4	0.15	0	6,6,6	0.06	0
5	SO4	B	307	-	4,4,4	0.13	0	6,6,6	0.05	0
3	PG4	F	301	-	12,12,12	0.53	0	11,11,11	0.22	0
3	PG4	A	401	-	3,3,12	0.48	0	2,2,11	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PG4	F	304	-	-	6/10/10/10	-
3	PG4	B	301	-	-	2/10/10/10	-
3	PG4	A	402	-	-	6/10/10/10	-
3	PG4	F	302	-	-	0/2/2/10	-
3	PG4	H	302	-	-	4/10/10/10	-
4	NAG	F	303	1	-	0/6/23/26	0/1/1/1
3	PG4	H	303	-	-	3/6/6/10	-
3	PG4	B	302	-	-	3/10/10/10	-
3	PG4	H	304	-	-	5/10/10/10	-
4	NAG	A	404	1	-	1/6/23/26	0/1/1/1
3	PG4	A	403	-	-	0/0/1/10	-
3	PG4	B	303	-	-	3/7/7/10	-
3	PG4	F	301	-	-	3/10/10/10	-
3	PG4	H	301	-	-	2/10/10/10	-
3	PG4	A	401	-	-	0/1/1/10	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	F	312	ACT	CH3-C	6.39	1.56	1.48
6	B	308	ACT	CH3-C	6.29	1.56	1.48

There are no bond angle outliers.

There are no chirality outliers.

5 of 38 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	H	303	PG4	C4-C3-O2-C2
3	H	304	PG4	C5-C6-O4-C7
3	H	301	PG4	O2-C3-C4-O3
3	B	302	PG4	O2-C3-C4-O3
3	H	304	PG4	O2-C3-C4-O3





There are no ring outliers.





6 monomers are involved in 8 short contacts:





Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	305	SO4	1	0
3	B	303	PG4	1	0
3	H	303	PG4	1	0
3	H	304	PG4	3	0
3	A	402	PG4	1	0
5	F	305	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	215/215 (100%)	-0.09	4 (1%) 66 71	16, 34, 59, 97	0
1	F	215/215 (100%)	-0.11	3 (1%) 75 78	18, 33, 56, 85	0
2	B	227/235 (96%)	-0.12	4 (1%) 68 72	18, 32, 62, 110	0
2	H	227/235 (96%)	-0.15	5 (2%) 62 66	17, 31, 63, 130	0
All	All	884/900 (98%)	-0.12	16 (1%) 68 72	16, 32, 61, 130	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	100(E)	THR	8.0
2	H	100(D)	ASP	5.7
1	F	212	GLY	4.7
2	B	99	GLY	4.4
2	H	100(G)	SER	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	PG4	B	302	13/13	0.59	0.23	68,77,78,79	0
4	NAG	A	404	14/15	0.60	0.27	96,100,102,104	0
3	PG4	F	301	13/13	0.61	0.21	52,62,64,65	0
3	PG4	A	403	4/13	0.67	0.32	52,53,53,54	0
3	PG4	B	301	13/13	0.67	0.29	70,74,75,76	0
4	NAG	F	303	14/15	0.70	0.22	84,85,90,91	0
3	PG4	F	304	13/13	0.73	0.20	63,64,71,71	0
3	PG4	A	402	13/13	0.73	0.23	64,66,68,69	0
6	ACT	B	308	4/4	0.74	0.26	90,90,91,91	0
3	PG4	F	302	5/13	0.81	0.33	63,63,63,64	0
5	SO4	B	306	5/5	0.81	0.25	106,106,107,107	0
3	PG4	B	303	10/13	0.82	0.21	57,57,69,70	0
3	PG4	H	303	9/13	0.82	0.20	48,58,64,65	0
3	PG4	H	304	13/13	0.84	0.23	51,62,65,65	0
3	PG4	H	302	13/13	0.87	0.14	42,48,57,58	0
5	SO4	F	311	5/5	0.87	0.15	56,57,62,67	0
5	SO4	B	307	5/5	0.88	0.15	77,79,79,80	0
5	SO4	H	305	5/5	0.88	0.18	104,105,105,105	0
6	ACT	F	312	4/4	0.90	0.07	55,58,59,59	0
5	SO4	H	307	5/5	0.90	0.16	80,82,83,83	0
5	SO4	B	305	5/5	0.91	0.16	93,93,95,96	0
5	SO4	B	304	5/5	0.91	0.20	75,76,77,79	0
3	PG4	H	301	13/13	0.92	0.17	47,51,56,56	0
5	SO4	H	306	5/5	0.92	0.16	78,80,81,83	0
3	PG4	A	401	4/13	0.92	0.10	52,52,55,55	0
5	SO4	F	306	5/5	0.94	0.14	66,68,69,69	0
5	SO4	F	307	5/5	0.95	0.15	59,60,61,62	0
5	SO4	F	310	5/5	0.96	0.09	55,62,63,64	0
5	SO4	F	309	5/5	0.98	0.09	51,53,53,56	0
5	SO4	F	308	5/5	0.98	0.07	50,51,52,56	0
5	SO4	F	305	5/5	0.98	0.11	43,47,53,55	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around PG4 B 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around PG4 F 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PG4 A 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PG4 B 301:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around PG4 F 304:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PG4 A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around PG4 F 302:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PG4 B 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around PG4 H 303:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PG4 H 304:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around PG4 H 302:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PG4 H 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around PG4 A 401:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.