



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 10, 2020 – 12:18 PM BST

PDB ID : 1UXA  
Title : ADENOVIRUS AD37 FIBRE HEAD in complex with sialyl-lactose  
Authors : Burmeister, W.P.; Guilligay, D.; Cusack, S.; Wadell, G.; Arnberg, N.  
Deposited on : 2004-02-24  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	194	<div> <div>2%</div> <div> <div></div> <div>82%</div> <div>12%</div> <div>• 5%</div> </div> </div>
1	B	194	<div> <div>7%</div> <div> <div></div> <div>87%</div> <div>7%</div> <div>• 6%</div> </div> </div>
1	C	194	<div> <div>3%</div> <div> <div></div> <div>84%</div> <div>10%</div> <div>• 6%</div> </div> </div>
2	D	2	<div> <div></div> <div> <div>50%</div> <div>50%</div> </div> </div>
2	E	2	<div> <div></div> <div> <div>50%</div> <div>50%</div> </div> </div>
3	F	2	<div> <div></div> <div> <div>50%</div> <div>50%</div> </div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4996 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FIBER PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	185	Total	C	N	O	S	0	0	0
			1467	942	238	283	4			
1	B	183	Total	C	N	O	S	0	0	0
			1447	929	236	278	4			
1	C	183	Total	C	N	O	S	0	0	0
			1447	929	236	278	4			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	173	ALA	TYR	engineered mutation	UNP Q64823
B	173	ALA	TYR	engineered mutation	UNP Q64823
C	173	ALA	TYR	engineered mutation	UNP Q64823
A	174	MET	LEU	engineered mutation	UNP Q64823
B	174	MET	LEU	engineered mutation	UNP Q64823
C	174	MET	LEU	engineered mutation	UNP Q64823
A	175	GLY	VAL	engineered mutation	UNP Q64823
B	175	GLY	VAL	engineered mutation	UNP Q64823
C	175	GLY	VAL	engineered mutation	UNP Q64823
A	176	SER	ALA	engineered mutation	UNP Q64823
B	176	SER	ALA	engineered mutation	UNP Q64823
C	176	SER	ALA	engineered mutation	UNP Q64823

- Molecule 2 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	2	Total	C	N	O	0	0	0
			32	17	1	14			
2	E	2	Total	C	N	O	0	0	0
			32	17	1	14			

- Molecule 3 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	F	2	Total	C	N	O	0	0	0
			32	17	1	14			

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		
4	A	1	Total	Zn	0	0
			1	1		
4	C	1	Total	Zn	0	0
			1	1		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total 4	C 2	O 2	0	0
5	C	1	Total 4	C 2	O 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	168	Total 168	O 168	0	0
6	B	164	Total 164	O 164	0	0
6	C	196	Total 196	O 196	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: FIBER PROTEIN



- Molecule 1: FIBER PROTEIN



- Molecule 1: FIBER PROTEIN



- Molecule 2: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose



- Molecule 2: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose



- Molecule 3: N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	60.70 Å 68.70 Å 74.70 Å 90.00° 94.90° 90.00°	Depositor
Resolution (Å)	45.18 – 1.50 49.03 – 1.50	Depositor EDS
% Data completeness (in resolution range)	92.0 (45.18-1.50) 91.9 (49.03-1.50)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.27 (at 1.50 Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.180 , 0.205 0.180 , 0.179	Depositor DCC
$R_{free}$ test set	4616 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.2	Xtriage
Anisotropy	0.109	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 50.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4996	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.16% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GAL, SIA, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.09	1/1502 (0.1%)	1.07	3/2040 (0.1%)
1	B	0.99	0/1481	1.03	3/2011 (0.1%)
1	C	1.06	0/1481	1.06	1/2011 (0.0%)
All	All	1.04	1/4464 (0.0%)	1.05	7/6062 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	304	ARG	CB-CG	-5.84	1.36	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	304	ARG	NE-CZ-NH2	-10.39	115.11	120.30
1	A	304	ARG	NE-CZ-NH2	-6.75	116.92	120.30
1	A	318	ASP	CB-CG-OD1	5.79	123.51	118.30
1	A	312	TYR	CB-CG-CD2	-5.75	117.55	121.00
1	C	188	THR	N-CA-C	5.28	125.27	111.00
1	B	304	ARG	CG-CD-NE	-5.21	100.87	111.80
1	B	191	ASP	CB-CG-OD1	5.14	122.92	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1467	0	1453	30	0
1	B	1447	0	1440	10	0
1	C	1447	0	1440	19	0
2	D	32	0	28	0	0
2	E	32	0	28	0	0
3	F	32	0	28	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
5	B	4	0	3	0	0
5	C	4	0	3	1	0
6	A	168	0	0	2	0
6	B	164	0	0	0	0
6	C	196	0	0	5	0
All	All	4996	0	4423	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:259:ASN:HD22	1:A:259:ASN:H	1.02	1.00
1:A:181:TYR:CA	1:A:184:ARG:HH11	1.84	0.90
1:A:181:TYR:O	1:A:212:LYS:HD3	1.79	0.82
1:C:268:ASN:HD22	1:C:269:PHE:H	1.28	0.81
1:C:183:THR:HB	1:C:269:PHE:CE1	2.18	0.77
1:A:259:ASN:HD22	1:A:259:ASN:N	1.83	0.76
1:A:181:TYR:CB	1:A:184:ARG:HH11	2.01	0.73
1:A:314:GLY:O	1:A:316:LYS:HD2	1.91	0.70
1:A:181:TYR:HA	1:A:184:ARG:HH11	1.56	0.69
1:A:181:TYR:HB3	1:A:286:PHE:HE1	1.57	0.69
1:A:270:ARG:HH11	1:B:215:SER:HB3	1.58	0.69
1:A:181:TYR:CB	1:A:286:PHE:HE1	2.06	0.69
1:C:276:VAL:HG21	6:C:2068:HOH:O	1.92	0.68
1:A:298:ASN:ND2	6:A:2093:HOH:O	2.25	0.68
1:A:259:ASN:H	1:A:259:ASN:ND2	1.81	0.66
1:C:187:TRP:HE1	1:C:275:ASN:ND2	1.95	0.64
1:C:295:LYS:HE2	6:C:2098:HOH:O	1.96	0.64
1:A:181:TYR:HA	1:A:184:ARG:HE	1.62	0.64

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:328:ASN:HD22	1:A:336:SER:H	1.42	0.64
1:C:328:ASN:HD22	1:C:336:SER:H	1.46	0.62
1:B:183:THR:O	1:B:183:THR:CG2	2.48	0.61
1:A:181:TYR:HB3	1:A:184:ARG:HH11	1.64	0.61
1:C:318:ASP:OD2	6:C:2131:HOH:O	2.16	0.60
1:A:181:TYR:HB3	1:A:184:ARG:NH1	2.18	0.58
1:A:181:TYR:HB2	1:A:286:PHE:CE1	2.39	0.58
1:C:268:ASN:ND2	1:C:269:PHE:H	2.00	0.57
1:C:268:ASN:HD22	1:C:269:PHE:N	1.99	0.57
1:A:181:TYR:CB	1:A:286:PHE:CE1	2.88	0.56
1:C:268:ASN:HB3	1:C:275:ASN:HB3	1.88	0.56
1:A:213:CYS:O	1:C:211:THR:HG21	2.06	0.55
1:C:281:GLU:HG2	6:C:2084:HOH:O	2.05	0.55
1:C:278:THR:HG23	6:C:2078:HOH:O	2.08	0.54
1:C:187:TRP:NE1	1:C:275:ASN:ND2	2.55	0.54
1:A:269:PHE:HB3	6:A:2062:HOH:O	2.08	0.53
1:B:328:ASN:HD22	1:B:336:SER:H	1.55	0.53
1:B:197:THR:HG23	1:B:200:GLN:C	2.31	0.51
1:C:274:SER:C	1:C:275:ASN:HD22	2.15	0.50
1:A:197:THR:HG23	1:A:200:GLN:O	2.12	0.49
1:A:211:THR:HG21	1:B:213:CYS:O	2.12	0.49
1:B:197:THR:HG23	1:B:200:GLN:O	2.13	0.49
1:A:197:THR:HG23	1:A:200:GLN:C	2.34	0.48
1:B:330:GLU:OE2	1:B:338:THR:HG21	2.13	0.48
1:A:181:TYR:HA	1:A:184:ARG:NH1	2.26	0.47
1:B:187:TRP:CH2	1:B:190:PRO:HD3	2.50	0.47
1:B:183:THR:O	1:B:183:THR:HG22	2.17	0.44
1:A:251:ASN:HB3	1:A:257:LEU:HD21	2.00	0.44
1:C:330:GLU:OE2	1:C:338:THR:HG21	2.17	0.43
1:C:232:ILE:HD11	5:C:1369:ACT:H2	2.01	0.43
1:B:259:ASN:OD1	1:B:259:ASN:N	2.52	0.43
1:A:314:GLY:C	1:A:316:LYS:HD2	2.39	0.42
1:A:328:ASN:ND2	1:A:336:SER:H	2.14	0.42
1:A:187:TRP:CH2	1:A:190:PRO:HD3	2.55	0.42
1:A:316:LYS:N	1:A:316:LYS:HD2	2.35	0.42
1:A:186:LEU:HD11	1:A:280:TYR:CE1	2.55	0.41
1:C:325:THR:HA	1:C:338:THR:O	2.21	0.41
1:C:183:THR:HB	1:C:269:PHE:CZ	2.56	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	183/194 (94%)	177 (97%)	5 (3%)	1 (0%)	29	9
1	B	181/194 (93%)	176 (97%)	5 (3%)	0	100	100
1	C	181/194 (93%)	171 (94%)	9 (5%)	1 (1%)	25	7
All	All	545/582 (94%)	524 (96%)	19 (4%)	2 (0%)	34	13

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	189	THR
1	A	182	ASP

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	165/171 (96%)	162 (98%)	3 (2%)	59	30
1	B	163/171 (95%)	160 (98%)	3 (2%)	59	30
1	C	163/171 (95%)	160 (98%)	3 (2%)	59	30
All	All	491/513 (96%)	482 (98%)	9 (2%)	59	30

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	181	TYR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	187	TRP
1	A	259	ASN
1	B	183	THR
1	B	187	TRP
1	B	277	SER
1	C	187	TRP
1	C	190	PRO
1	C	268	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	259	ASN
1	A	328	ASN
1	A	340	ASN
1	B	238	ASN
1	B	328	ASN
1	B	340	ASN
1	C	268	ASN
1	C	275	ASN
1	C	298	ASN
1	C	328	ASN
1	C	340	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	GAL	D	1	2	12,12,12	0.44	0	17,17,17	0.67	0
2	SIA	D	2	2	17,20,21	1.14	2 (11%)	21,28,31	1.15	3 (14%)
2	GAL	E	1	2	12,12,12	0.44	0	17,17,17	0.52	0
2	SIA	E	2	2	17,20,21	0.97	1 (5%)	21,28,31	0.98	1 (4%)
3	GAL	F	1	3	12,12,12	0.61	0	17,17,17	0.79	0
3	SIA	F	2	3	17,20,21	0.95	0	21,28,31	1.02	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GAL	D	1	2	-	0/2/22/22	0/1/1/1
2	SIA	D	2	2	-	0/14/34/38	0/1/1/1
2	GAL	E	1	2	-	0/2/22/22	0/1/1/1
2	SIA	E	2	2	-	0/14/34/38	0/1/1/1
3	GAL	F	1	3	-	0/2/22/22	0/1/1/1
3	SIA	F	2	3	-	0/14/34/38	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2	SIA	C11-C10	2.56	1.55	1.50
2	D	2	SIA	C4-C5	2.50	1.55	1.53
2	E	2	SIA	C7-C6	2.39	1.56	1.53

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2	SIA	O10-C10-N5	2.14	125.89	121.95
3	F	2	SIA	C3-C4-C5	-2.14	108.88	111.46
2	E	2	SIA	O10-C10-N5	2.09	125.80	121.95
2	D	2	SIA	C4-C3-C2	2.07	113.52	109.81
2	D	2	SIA	C4-C5-N5	-2.05	106.31	110.38


There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## Oligosaccharide Chain E



## Oligosaccharide Chain F





## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	ACT	C	1369	4	1,3,3	2.12	1 (100%)	0,3,3	0.00	-
5	ACT	B	1369	4	1,3,3	4.13	1 (100%)	0,3,3	0.00	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1369	ACT	CH3-C	4.13	1.54	1.48
5	C	1369	ACT	CH3-C	2.12	1.51	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	1369	ACT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	185/194 (95%)	-0.06	3 (1%) 72 77	12, 19, 34, 51	0
1	B	183/194 (94%)	0.11	13 (7%) 16 17	11, 22, 44, 57	0
1	C	183/194 (94%)	0.03	6 (3%) 46 51	10, 17, 34, 50	0
All	All	551/582 (94%)	0.03	22 (3%) 38 42	10, 19, 38, 57	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	181	TYR	11.0
1	A	182	ASP	5.7
1	C	183	THR	4.4
1	B	276	VAL	4.2
1	C	276	VAL	3.9
1	B	266	TYR	3.9
1	B	277	SER	3.8
1	B	183	THR	3.5
1	B	240	LYS	3.3
1	B	281	GLU	2.9
1	B	258	ASP	2.5
1	B	331	THR	2.5
1	B	199	ALA	2.5
1	B	279	ALA	2.4
1	C	273	ASN	2.4
1	B	278	THR	2.4
1	C	275	ASN	2.3
1	C	271	SER	2.3
1	B	241	ILE	2.3
1	C	272	GLY	2.3
1	B	239	PRO	2.2
1	A	266	TYR	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GAL	D	1	12/12	0.84	0.11	29,42,45,52	12
3	GAL	F	1	12/12	0.91	0.21	11,17,26,31	12
2	GAL	E	1	12/12	0.91	0.12	22,36,41,47	12
2	SIA	D	2	20/21	0.92	0.20	8,20,31,42	20
2	SIA	E	2	20/21	0.92	0.12	10,16,22,22	20
3	SIA	F	2	20/21	0.94	0.14	8,13,19,29	20

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around Chain D:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain F:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	ACT	B	1369	4/4	0.90	0.18	21,25,30,33	0
5	ACT	C	1369	4/4	0.98	0.09	9,27,31,36	0
4	ZN	B	1367	1/1	0.98	0.05	20,20,20,20	0
4	ZN	A	1367	1/1	0.99	0.06	27,27,27,27	0
4	ZN	C	1367	1/1	0.99	0.04	23,23,23,23	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.