



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 06:05 am BST

PDB ID : 4UYQ
Title : High resolution structure of the third cohesin ScaC in complex with the ScaB dockerin with a mutation in the C-terminal helix (IN to SI) from *Acetivibrio cellulolyticus* displaying a type I interaction.
Authors : Cameron, K.; Fontes, C.M.G.A.; Najmudin, S.
Deposited on : 2014-09-02
Resolution : 1.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	7484 (1.84-1.80)
Clashscore	141614	8401 (1.84-1.80)
Ramachandran outliers	138981	8290 (1.84-1.80)
Sidechain outliers	138945	8290 (1.84-1.80)
RSRZ outliers	127900	7371 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	151	
2	B	75	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2003 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cellulosomal scaffoldin anchoring protein C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	144	Total	C	N	O	S	0	1	1
			1092	691	173	226	2			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP Q7WYN2
A	144	LEU	-	expression tag	UNP Q7WYN2
A	145	GLU	-	expression tag	UNP Q7WYN2
A	146	HIS	-	expression tag	UNP Q7WYN2
A	147	HIS	-	expression tag	UNP Q7WYN2
A	148	HIS	-	expression tag	UNP Q7WYN2
A	149	HIS	-	expression tag	UNP Q7WYN2
A	150	HIS	-	expression tag	UNP Q7WYN2
A	151	HIS	-	expression tag	UNP Q7WYN2

- Molecule 2 is a protein called Cellulosomal scaffoldin adaptor protein B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	73	Total	C	N	O	S	0	3	0
			589	385	92	111	1			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	51	SER	ILE	engineered mutation	UNP Q7WYN3
B	52	ILE	ASN	engineered mutation	UNP Q7WYN3

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total 2	Ca 2	0	0
3	A	1	Total 1	Ca 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	232	Total 232	O 232	0	0
4	B	87	Total 87	O 87	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cellulosomal scaffoldin anchoring protein C

Chain A:  87% 8% 5%



- Molecule 2: Cellulosomal scaffoldin adaptor protein B

Chain B:  4% 76% 20%



4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	68.17Å 68.17Å 57.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	68.17 – 1.81 30.49 – 1.81	Depositor EDS
% Data completeness (in resolution range)	96.8 (68.17-1.81) 96.8 (30.49-1.81)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.91 (at 1.82Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.150 , 0.195 0.162 , 0.207	Depositor DCC
R_{free} test set	1174 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	17.9	Xtriage
Anisotropy	0.601	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 45.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.048 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2003	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.20	1/1115 (0.1%)	1.10	7/1517 (0.5%)
2	B	1.09	2/608 (0.3%)	1.20	2/820 (0.2%)
All	All	1.16	3/1723 (0.2%)	1.13	9/2337 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	1	0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	73	ALA	C-N	21.26	1.82	1.34
2	B	72[A]	GLU	CD-OE1	6.54	1.32	1.25
2	B	72[B]	GLU	CD-OE1	6.54	1.32	1.25

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	83	ARG	NE-CZ-NH2	-12.37	114.11	120.30
2	B	58	ARG	NE-CZ-NH1	8.37	124.49	120.30
1	A	83	ARG	NE-CZ-NH1	7.87	124.23	120.30
1	A	73	ALA	CA-C-O	7.41	135.66	120.10
1	A	73	ALA	O-C-N	-6.26	112.68	122.70
1	A	109	LEU	CB-CG-CD1	6.11	121.39	111.00
1	A	109	LEU	CA-CB-CG	-5.54	102.55	115.30
2	B	8	ASP	CB-CG-OD1	5.20	122.98	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	37	ARG	NE-CZ-NH1	-5.06	117.77	120.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	1[A]	LYS	CA

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1092	0	1056	15	0
2	B	589	0	603	22	0
3	A	1	0	0	0	0
3	B	2	0	0	0	0
4	A	232	0	0	16	3
4	B	87	0	0	13	2
All	All	2003	0	1659	37	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73:ALA:C	1:A:74:SER:N	1.82	1.31
1:A:141:VAL:HG12	4:A:2025:HOH:O	1.28	1.28
2:B:1[A]:LYS:NZ	4:B:2004:HOH:O	1.66	1.26
4:A:2056:HOH:O	2:B:19[B]:VAL:HG21	1.49	1.10
1:A:141:VAL:CG1	4:A:2025:HOH:O	1.89	1.07
1:A:118:TYR:HD2	4:A:2189:HOH:O	1.40	1.03
1:A:115:ASN:OD1	4:A:2189:HOH:O	1.78	1.01
4:A:2056:HOH:O	2:B:19[B]:VAL:CG2	2.10	0.92
2:B:61:VAL:HG11	4:B:2032:HOH:O	1.74	0.87

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:50:LYS:NZ	4:B:2064:HOH:O	2.07	0.85
1:A:115:ASN:CG	4:A:2189:HOH:O	2.17	0.82
1:A:118:TYR:CD2	4:A:2189:HOH:O	2.19	0.81
1:A:127:ASP:OD1	4:A:2213:HOH:O	2.00	0.79
1:A:118:TYR:HB2	4:A:2189:HOH:O	1.88	0.73
2:B:61:VAL:HG21	4:B:2032:HOH:O	1.89	0.72
2:B:71:GLU:OE2	4:B:2086:HOH:O	2.11	0.68
2:B:15:ILE:HA	4:B:2032:HOH:O	1.94	0.66
2:B:15:ILE:CA	4:B:2032:HOH:O	2.47	0.61
4:A:2056:HOH:O	2:B:22:ARG:NH2	2.33	0.61
1:A:101:SER:HB3	4:A:2086:HOH:O	2.03	0.57
2:B:38:MET:SD	2:B:49:ILE:CD1	2.92	0.56
2:B:72[A]:GLU:HG3	2:B:73:LYS:HD3	1.88	0.56
1:A:131:ASN:OD1	1:A:131:ASN:N	2.29	0.54
4:A:2056:HOH:O	2:B:19[A]:VAL:CG1	2.57	0.53
2:B:72[A]:GLU:OE2	4:B:2087:HOH:O	2.19	0.50
4:A:2056:HOH:O	2:B:19[A]:VAL:HG11	2.12	0.49
2:B:72[A]:GLU:HG3	2:B:73:LYS:CD	2.44	0.48
2:B:61:VAL:CB	4:B:2032:HOH:O	2.62	0.48
2:B:38:MET:SD	2:B:49:ILE:HD12	2.54	0.47
1:A:73:ALA:CA	1:A:74:SER:N	2.72	0.45
1:A:118:TYR:CB	4:A:2189:HOH:O	2.56	0.45
2:B:25:VAL:HB	4:B:2036:HOH:O	2.18	0.43
2:B:26:LEU:HG	4:B:2036:HOH:O	2.18	0.43
1:A:32:TYR:CE2	2:B:19[A]:VAL:HG11	2.54	0.42
1:A:12:LYS:HE3	4:A:2023:HOH:O	2.20	0.40
2:B:34:TYR:HE2	4:B:2009:HOH:O	2.03	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:2153:HOH:O	4:B:2072:HOH:O[4_575]	2.05	0.15
4:A:2058:HOH:O	4:B:2072:HOH:O[4_575]	2.12	0.08
4:A:2028:HOH:O	4:A:2179:HOH:O[4_675]	2.15	0.05

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	143/151 (95%)	143 (100%)	0	0	100	100
2	B	73/75 (97%)	72 (99%)	1 (1%)	0	100	100
All	All	216/226 (96%)	215 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	122/129 (95%)	122 (100%)	0	100	100
2	B	64/63 (102%)	64 (100%)	0	100	100
All	All	186/192 (97%)	186 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	73:ALA	C	74:SER	N	1.82

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	144/151 (95%)	-0.35	0	100 100	12, 20, 38, 49	0
2	B	73/75 (97%)	-0.00	3 (4%)	37 31	13, 20, 34, 62	0
All	All	217/226 (96%)	-0.23	3 (1%)	75 72	12, 20, 38, 62	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	73	LYS	3.1
2	B	61	VAL	2.1
2	B	30	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CA	B	202	1/1	0.99	0.15	34,34,34,34	0
3	CA	B	201	1/1	1.00	0.07	15,15,15,15	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	CA	A	200	1/1	1.00	0.06	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.