



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 11:08 pm BST

PDB ID : 3V07
Title : Crystal structure of R-6'-Me-3'-fluoro hexitol nucleic acid
Authors : Pallan, P.S.; Egli, M.
Deposited on : 2011-12-07
Resolution : 1.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2024 (1.28-1.20)
Clashscore	141614	1007 (1.26-1.22)
RSRZ outliers	127900	1987 (1.28-1.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	10	<div> <div></div> <div>10%</div> <div>90%</div> </div>
1	B	10	<div> <div></div> <div>30%</div> <div>60%</div> <div>10%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 490 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*GP*CP*GP*TP*AP*(F6H)P*AP*CP*GP*C)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	10	Total	C	F	N	O	P	0	0	0
			205	99	1	38	58	9			
1	B	10	Total	C	F	N	O	P	0	0	0
			205	99	1	38	58	9			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	37	Total	O	0	0
			37	37		
2	B	43	Total	O	0	0
			43	43		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D(*GP*CP*GP*TP*AP*(F6H)P*AP*CP*GP*C)-3')

Chain A: 

G101	G102	G103	T104	A105	T106	A107	G108	G109	C110
------	------	------	------	------	------	------	------	------	------

- Molecule 1: DNA (5'-D(*GP*CP*GP*TP*AP*(F6H)P*AP*CP*GP*C)-3')

Chain B: 

G201	C202	G203	T204	A205	T206	A207	C208	G209	C210
------	------	------	------	------	------	------	------	------	------

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	27.30Å 43.48Å 45.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.45 – 1.24 31.45 – 1.24	Depositor EDS
% Data completeness (in resolution range)	98.2 (31.45-1.24) 98.2 (31.45-1.24)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.62 (at 1.24Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.159 , 0.207 0.157 , 0.206	Depositor DCC
R_{free} test set	776 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	14.7	Xtriage
Anisotropy	0.774	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.023 for -h,l,k	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	490	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: F6H

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	2.28	6/203 (3.0%)	2.81	29/309 (9.4%)
1	B	2.41	10/203 (4.9%)	2.44	14/309 (4.5%)
All	All	2.35	16/406 (3.9%)	2.63	43/618 (7.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	109	DG	N7-C5	7.96	1.44	1.39
1	A	101	DG	N7-C5	7.92	1.44	1.39
1	B	201	DG	C8-N7	7.85	1.35	1.30
1	B	203	DG	N9-C4	7.23	1.43	1.38
1	B	210	DC	C4'-O4'	6.29	1.51	1.45
1	B	203	DG	N7-C5	6.02	1.42	1.39
1	A	107	DA	P-O5'	5.76	1.65	1.59
1	A	110	DC	C4'-C3'	-5.75	1.46	1.52
1	B	205	DA	N3-C4	5.58	1.38	1.34
1	B	208	DC	C3'-O3'	5.53	1.51	1.44
1	A	103	DG	C6-N1	5.28	1.43	1.39
1	B	201	DG	N3-C4	5.27	1.39	1.35
1	B	210	DC	P-OP2	-5.25	1.40	1.49
1	B	209	DG	C6-N1	-5.24	1.35	1.39
1	B	208	DC	C2'-C1'	5.12	1.57	1.52
1	A	108	DC	N3-C4	-5.08	1.30	1.33

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	210	DC	N3-C4-C5	11.74	126.60	121.90
1	A	108	DC	C2-N3-C4	10.83	125.31	119.90
1	B	210	DC	C4-C5-C6	-10.34	112.23	117.40
1	A	109	DG	O4'-C4'-C3'	-10.10	99.94	106.00
1	A	102	DC	N1-C2-O2	-9.73	113.06	118.90
1	A	107	DA	N1-C2-N3	-9.64	124.48	129.30
1	A	105	DA	O4'-C4'-C3'	-9.42	100.35	106.00
1	A	108	DC	N3-C4-C5	-8.92	118.33	121.90
1	A	109	DG	N9-C4-C5	8.67	108.87	105.40
1	A	108	DC	O4'-C4'-C3'	-7.85	101.29	106.00
1	A	109	DG	C4-C5-N7	-7.59	107.76	110.80
1	A	110	DC	C5-C6-N1	-7.30	117.35	121.00
1	B	209	DG	C2-N3-C4	7.18	115.49	111.90
1	A	105	DA	C2-N3-C4	7.04	114.12	110.60
1	B	207	DA	N7-C8-N9	-7.04	110.28	113.80
1	A	102	DC	C6-N1-C2	-6.87	117.55	120.30
1	A	107	DA	C4-C5-N7	6.82	114.11	110.70
1	B	207	DA	C4'-C3'-C2'	-6.56	97.20	103.10
1	B	207	DA	N1-C2-N3	-6.53	126.04	129.30
1	A	110	DC	C6-N1-C2	6.25	122.80	120.30
1	A	105	DA	N1-C2-N3	-6.20	126.20	129.30
1	A	108	DC	C5-C6-N1	6.15	124.08	121.00
1	A	103	DG	C4-C5-N7	-6.11	108.36	110.80
1	A	102	DC	C5-C4-N4	6.08	124.45	120.20
1	B	209	DG	O4'-C4'-C3'	-6.05	102.08	104.50
1	A	104	DT	C2-N3-C4	-5.99	123.61	127.20
1	B	205	DA	C2-N3-C4	-5.97	107.62	110.60
1	B	203	DG	O4'-C1'-C2'	5.95	110.66	105.90
1	A	107	DA	C6-N1-C2	5.74	122.04	118.60
1	B	209	DG	N1-C2-N3	-5.70	120.48	123.90
1	A	102	DC	O4'-C4'-C3'	-5.69	102.23	104.50
1	A	104	DT	N3-C4-C5	5.58	118.55	115.20
1	B	209	DG	C5-C6-N1	5.54	114.27	111.50
1	A	102	DC	N3-C4-N4	-5.51	114.15	118.00
1	A	107	DA	O4'-C4'-C3'	-5.45	102.32	104.50
1	A	109	DG	N3-C4-N9	-5.40	122.76	126.00
1	B	201	DG	O4'-C1'-C2'	5.33	110.16	105.90
1	A	108	DC	N3-C4-N4	5.22	121.65	118.00
1	B	205	DA	N3-C4-N9	-5.20	123.24	127.40
1	B	205	DA	N9-C4-C5	5.20	107.88	105.80
1	A	107	DA	N3-C4-C5	5.14	130.40	126.80
1	A	110	DC	C4-C5-C6	5.14	119.97	117.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	102	DC	C4'-C3'-C2'	-5.01	98.59	103.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	209	DG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	205	0	117	0	0
1	B	205	0	117	0	0
2	A	37	0	0	0	0
2	B	43	0	0	0	0
All	All	490	0	234	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	F6H	A	106	1	16,24,25	1.58	4 (25%)	12,35,38	2.68	3 (25%)
1	F6H	B	206	1	16,24,25	1.19	0	12,35,38	2.95	4 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	F6H	A	106	1	-	0/6/29/30	0/2/2/2
1	F6H	B	206	1	-	0/6/29/30	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	106	F6H	O4'-C4'	2.85	1.49	1.43
1	A	106	F6H	C1'-C2'	2.69	1.55	1.52
1	A	106	F6H	C3'-C4'	2.40	1.54	1.52
1	A	106	F6H	O4-C4	2.25	1.30	1.24

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	206	F6H	C4-N3-C2	8.52	122.34	115.14
1	A	106	F6H	C4-N3-C2	7.04	121.09	115.14
1	A	106	F6H	C1'-O5'-C5'	4.00	118.03	111.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	206	F6H	C1'-O5'-C5'	3.57	117.34	111.48
1	A	106	F6H	C7'-C6'-C5'	3.23	120.44	113.34
1	B	206	F6H	C5-C6-N1	2.96	125.39	122.19
1	B	206	F6H	C7-C5-C6	2.12	123.15	118.68

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	9/10 (90%)	-0.30	0 100 100	15, 16, 18, 18	0
1	B	9/10 (90%)	-0.40	0 100 100	16, 16, 18, 20	0
All	All	18/20 (90%)	-0.35	0 100 100	15, 16, 18, 20	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	F6H	A	106	23/24	0.98	0.06	13,16,19,22	0
1	F6H	B	206	23/24	0.98	0.07	13,16,19,20	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.