



Full wwPDB NMR Structure Validation Report ⓘ

Jun 20, 2022 – 10:19 AM EDT

PDB ID : 7V06
Title : Encoded Conformational Dynamics of the HIV Splice Site A3 Regulatory Locus: Implications for differential binding of hnRNP Splicing Auxiliary Factors
Authors : Tolbert, B.S.; Chiu, L.Y.; Sugarman, A.L.
Deposited on : 2022-05-09

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	2.29
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.29


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR, SOLUTION SCATTERING

The overall completeness of chemical shifts assignment is 17%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
RNA backbone	4643	676

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	43	 35% 65%

2 Ensemble composition and analysis ⓘ

This entry contains 10 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

3 Entry composition [i](#)


There is only 1 type of molecule in this entry. The entry contains 1369 atoms, of which 461 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called RNA (43-MER).

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
1	A	43	1369	407	461	155	304	42	0


4.2.3 Score per residue for model 3

- Molecule 1: RNA (43-MER)




4.2.4 Score per residue for model 4

- Molecule 1: RNA (43-MER)




4.2.5 Score per residue for model 5

- Molecule 1: RNA (43-MER)




4.2.6 Score per residue for model 6

- Molecule 1: RNA (43-MER)




4.2.7 Score per residue for model 7

- Molecule 1: RNA (43-MER)




4.2.8 Score per residue for model 8

- Molecule 1: RNA (43-MER)




4.2.9 Score per residue for model 9

- Molecule 1: RNA (43-MER)



4.2.10 Score per residue for model 10

- Molecule 1: RNA (43-MER)



5 Refinement protocol and experimental data overview

The models were refined using the following method: *molecular dynamics*.

Of the 1024 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Amber	refinement	
X-PLOR NIH	structure calculation	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	137
Number of shifts mapped to atoms	137
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	17%

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.37±0.01	0±0/1013 (0.0± 0.0%)	2.24±0.04	58±4/1576 (3.7± 0.3%)
All	All	1.37	0/10130 (0.0%)	2.24	585/15760 (3.7%)

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	20	A	N1-C6-N6	-11.89	111.47	118.60	7	10
1	A	11	A	C5-C6-N1	11.37	123.38	117.70	10	10
1	A	11	A	N1-C6-N6	-10.36	112.38	118.60	10	10
1	A	22	A	N1-C6-N6	-9.99	112.61	118.60	2	10
1	A	36	A	N1-C6-N6	-9.70	112.78	118.60	7	10
1	A	29	U	N3-C2-O2	-9.44	115.59	122.20	10	3
1	A	34	A	N1-C6-N6	-9.15	113.11	118.60	10	10
1	A	23	A	N1-C6-N6	-9.08	113.15	118.60	9	10
1	A	15	A	N1-C6-N6	-8.88	113.28	118.60	9	10
1	A	11	A	C4-C5-C6	-8.52	112.74	117.00	9	10
1	A	24	U	N3-C2-O2	-8.42	116.31	122.20	9	9
1	A	25	U	N3-C2-O2	-8.42	116.31	122.20	6	1
1	A	20	A	C5-C6-N1	7.89	121.65	117.70	4	10
1	A	20	A	C4-C5-C6	-7.89	113.06	117.00	7	10
1	A	38	A	N1-C6-N6	-7.86	113.88	118.60	7	10
1	A	22	A	C5-C6-N1	7.70	121.55	117.70	9	10
1	A	13	C	N3-C2-O2	-7.64	116.55	121.90	9	10
1	A	14	C	N3-C2-O2	-7.63	116.56	121.90	3	10
1	A	43	C	N3-C2-O2	-7.60	116.58	121.90	4	10
1	A	15	A	C5-C6-N1	7.57	121.49	117.70	6	10
1	A	41	A	N1-C6-N6	-7.52	114.09	118.60	1	10
1	A	5	C	N3-C2-O2	-7.50	116.65	121.90	8	10
1	A	34	A	C5-C6-N1	7.44	121.42	117.70	9	10
1	A	42	C	N3-C2-O2	-7.36	116.75	121.90	9	10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	22	A	C4-C5-C6	-7.33	113.33	117.00	1	10
1	A	23	A	C5-C6-N1	7.31	121.35	117.70	9	10
1	A	12	U	N3-C2-O2	-7.30	117.09	122.20	10	3
1	A	36	A	C5-C6-N1	7.26	121.33	117.70	10	10
1	A	40	C	N3-C2-O2	-7.25	116.83	121.90	5	10
1	A	15	A	C4-C5-C6	-7.19	113.40	117.00	8	10
1	A	28	G	C4-N9-C1'	7.11	135.74	126.50	10	3
1	A	32	C	N3-C2-O2	-7.01	116.99	121.90	1	10
1	A	19	C	N3-C2-O2	-7.01	116.99	121.90	6	10
1	A	38	A	C5-C6-N1	6.79	121.09	117.70	10	10
1	A	41	A	C5-C6-N1	6.77	121.09	117.70	9	10
1	A	23	A	C4-C5-C6	-6.72	113.64	117.00	9	10
1	A	35	C	N3-C2-O2	-6.62	117.26	121.90	7	10
1	A	41	A	C4-C5-C6	-6.48	113.76	117.00	9	10
1	A	38	A	C4-C5-C6	-6.47	113.76	117.00	1	10
1	A	9	U	O4'-C1'-N1	6.47	113.38	108.20	8	8
1	A	36	A	C4-C5-C6	-6.38	113.81	117.00	7	10
1	A	13	C	N1-C2-O2	6.37	122.72	118.90	10	10
1	A	11	A	C6-C5-N7	6.33	136.73	132.30	9	3
1	A	28	G	C8-N9-C1'	-6.29	118.82	127.00	10	3
1	A	30	G	N1-C6-O6	-6.26	116.14	119.90	7	5
1	A	34	A	C4-C5-C6	-6.24	113.88	117.00	3	10
1	A	32	C	O4'-C1'-N1	6.11	113.09	108.20	5	4
1	A	19	C	O4'-C1'-N1	6.09	113.07	108.20	4	3
1	A	5	C	N1-C2-O2	6.08	122.55	118.90	8	10
1	A	7	G	N1-C6-O6	-6.08	116.25	119.90	6	10
1	A	29	U	N1-C2-O2	6.04	127.03	122.80	8	3
1	A	30	G	N3-C4-C5	-6.01	125.60	128.60	9	1
1	A	28	G	N1-C6-O6	-5.93	116.34	119.90	8	10
1	A	24	U	C2-N1-C1'	5.92	124.80	117.70	9	1
1	A	25	U	N1-C2-O2	5.87	126.91	122.80	6	1
1	A	8	U	O4'-C1'-N1	5.86	112.88	108.20	1	3
1	A	24	U	N1-C2-O2	5.75	126.83	122.80	9	1
1	A	25	U	C2-N1-C1'	5.72	124.57	117.70	6	1
1	A	8	U	N3-C2-O2	-5.72	118.20	122.20	2	7
1	A	19	C	N1-C2-O2	5.71	122.32	118.90	6	6
1	A	42	C	N1-C2-O2	5.70	122.32	118.90	4	10
1	A	28	G	N7-C8-N9	5.69	115.94	113.10	8	3
1	A	14	C	N1-C2-O2	5.68	122.31	118.90	3	9
1	A	20	A	C6-C5-N7	5.67	136.27	132.30	7	3
1	A	43	C	N1-C2-O2	5.67	122.30	118.90	1	10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	18	U	O4'-C1'-N1	5.66	112.73	108.20	4	5
1	A	40	C	N1-C2-O2	5.64	122.28	118.90	9	10
1	A	17	U	O4'-C1'-N1	5.58	112.66	108.20	10	5
1	A	33	G	N1-C6-O6	-5.54	116.58	119.90	10	8
1	A	24	U	O4'-C1'-N1	5.54	112.63	108.20	1	8
1	A	32	C	N1-C2-O2	5.50	122.20	118.90	1	3
1	A	28	G	N3-C4-C5	-5.50	125.85	128.60	10	3
1	A	30	G	C5-C6-N1	5.48	114.24	111.50	5	5
1	A	35	C	O4'-C1'-N1	5.47	112.58	108.20	7	1
1	A	15	A	C6-C5-N7	5.26	135.98	132.30	8	7
1	A	11	A	C6-N1-C2	-5.17	115.50	118.60	10	2
1	A	11	A	O4'-C1'-N9	5.15	112.32	108.20	9	2
1	A	31	U	O4'-C1'-N1	5.15	112.32	108.20	3	3
1	A	18	U	N3-C2-O2	-5.13	118.61	122.20	5	5
1	A	10	U	N3-C2-O2	-5.12	118.62	122.20	1	1
1	A	6	U	N3-C2-O2	-5.12	118.61	122.20	8	6
1	A	9	U	N1-C2-N3	5.11	117.96	114.90	8	3
1	A	35	C	N1-C2-O2	5.06	121.94	118.90	7	2
1	A	18	U	N1-C2-N3	5.06	117.94	114.90	4	1
1	A	25	U	C1'-O4'-C4'	-5.00	105.90	109.90	9	1

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	908	461	461	0±0
All	All	9080	4610	4610	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:11:A:H62	1:A:28:G:N2	0.41	2.14	9	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

6.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	A	42/43 (98%)	2±0 (4±1%)	0±0 (0±0%)	0.58±0.02
All	All	420/430 (98%)	17 (4%)	0 (0%)	0.58

The overall RNA backbone suiteness is 0.58.

All unique RNA backbone outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	A	21	G	6
1	A	19	C	4
1	A	32	C	3
1	A	28	G	3
1	A	20	A	1

There are no RNA pucker outliers to report.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry

There are no ligands in this entry.

6.7 Other polymers

There are no such molecules in this entry.

6.8 Polymer linkage issues

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 17% for the well-defined parts and 17% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chemical_shifts_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	137
Number of shifts mapped to atoms	137
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 17%, i.e. 137 atoms were assigned a chemical shift out of a possible 800. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Sidechain	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Aromatic	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Overall	137/800 (17%)	137/456 (30%)	0/290 (0%)	0/54 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 17%, i.e. 137 atoms were assigned a chemical shift out of a possible 800. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Sidechain	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Aromatic	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Overall	137/800 (17%)	137/456 (30%)	0/290 (0%)	0/54 (0%)

7.1.4 Statistically unusual chemical shifts ⓘ

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots ⓘ

No *random coil index* (RCI) plot could be generated from the current chemical shift list (assigned_chemical_shifts_1). RCI is only applicable to proteins.