



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 06:31 AM BST

PDB ID : 5V2K
Title : Crystal structure of UDP-glucosyltransferase, UGT74F2 (T15A), with UDP and 2-bromobenzoic acid
Authors : George Thompson, A.M.; Iancu, C.V.; Dean, J.V.; Choe, J.
Deposited on : 2017-03-04
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	449	<div> <div>2%</div> <div> <div></div> <div>88%</div> <div>10%</div> <div>..</div> </div> </div>
1	B	449	<div> <div>5%</div> <div> <div></div> <div>89%</div> <div>9%</div> <div>.</div> </div> </div>
2	C	2	<div> <div></div> <div>100%</div> </div>
2	D	2	<div> <div></div> <div>50%</div> <div>50%</div> </div>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7322 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UDP-glycosyltransferase 74F2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	444	Total	C	N	O	S	0	0	0
			3535	2276	572	671	16			
1	B	444	Total	C	N	O	S	0	0	0
			3535	2276	572	671	16			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	15	ALA	THR	engineered mutation	UNP O22822
B	15	ALA	THR	engineered mutation	UNP O22822

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-3)-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	2	Total	C	O	0	0	0
			22	12	10			
2	D	2	Total	C	O	0	0	0
			22	12	10			

- Molecule 3 is beta-D-glucopyranose (three-letter code: BGC) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		
3	B	1	Total	C	O	0	0
			11	6	5		

- Molecule 4 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula: C₉H₁₄N₂O₁₂P₂).



- Molecule 5 is 2-bromobenzoic acid (three-letter code: 7WV) (formula: $C_7H_5BrO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	Br	C	O	0	0
			10	1	7	2		
5	B	1	Total	Br	C	O	0	0
			10	1	7	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	76	Total	O	0	0
			76	76		
6	B	40	Total	O	0	0
			40	40		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.12Å 87.27Å 162.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.93 – 2.00 43.93 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.6 (43.93-2.00) 99.6 (43.93-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.72 (at 2.00Å)	Xtriage
Refinement program	PHENIX (dev_2666: ???)	Depositor
R, R_{free}	0.193 , 0.244 0.193 , 0.244	Depositor DCC
R_{free} test set	3060 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å ²)	43.3	Xtriage
Anisotropy	0.207	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 41.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7322	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, UDP, 7WV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.46	0/3624	0.57	0/4920
1	B	0.39	0/3624	0.53	0/4920
All	All	0.43	0/7248	0.55	0/9840

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	3
All	All	0	4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	251	ASP	Peptide
1	B	316	LYS	Peptide
1	B	50	PRO	Peptide
1	B	51	ASP	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3535	0	3470	27	0
1	B	3535	0	3470	23	0
2	C	22	0	19	0	0
2	D	22	0	19	0	0
3	A	11	0	9	0	0
3	B	11	0	10	0	0
4	A	25	0	11	0	0
4	B	25	0	11	0	0
5	A	10	0	0	0	0
5	B	10	0	0	0	0
6	A	76	0	0	1	0
6	B	40	0	0	0	0
All	All	7322	0	7019	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:363:GLN:H	1:B:367:GLN:HE21	1.41	0.67
1:B:52:LEU:HD21	1:B:56:ILE:HG21	1.77	0.66
1:A:23:ARG:NH1	1:A:242:ASP:OD1	2.28	0.65
1:B:95:ILE:HA	1:B:105:ILE:HD12	1.79	0.65
1:B:312:GLU:N	1:B:312:GLU:OE1	2.30	0.64
1:A:361:MET:HE2	1:A:384:VAL:HG21	1.80	0.62
1:B:363:GLN:H	1:B:367:GLN:NE2	2.00	0.59
1:A:279:ASN:O	1:A:283:GLU:HG2	2.02	0.59
1:A:375:GLN:NE2	1:A:383:ARG:H	2.03	0.57
1:B:52:LEU:HD21	1:B:56:ILE:CG2	2.34	0.57
1:A:375:GLN:HG2	1:A:376:ASP:OD1	2.05	0.56
1:A:412:LYS:HD3	1:A:415:LYS:HE2	1.88	0.55
1:A:415:LYS:HG3	1:A:416:LYS:N	2.21	0.55
1:A:23:ARG:HE	1:A:56:ILE:HD12	1.72	0.54
1:B:85:THR:O	1:B:89:LYS:HG2	2.09	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:389:GLU:HG2	1:B:390:SER:OG	2.10	0.52
1:B:5:ARG:NH2	1:B:447:GLN:O	2.35	0.52
1:A:153:GLN:OE1	1:A:153:GLN:N	2.43	0.52
1:A:386:THR:HG22	1:A:393:ALA:HB2	1.92	0.51
1:B:269:VAL:HG13	1:B:296:TRP:HA	1.91	0.51
1:B:88:SER:HB2	1:B:89:LYS:HE3	1.94	0.49
1:B:231:LEU:HD21	1:B:353:THR:CG2	2.43	0.49
1:A:415:LYS:O	1:A:419:LYS:HG2	2.13	0.49
1:A:15:ALA:HB1	1:A:274:MET:HE1	1.93	0.49
1:B:52:LEU:HD11	1:B:56:ILE:HD12	1.95	0.48
1:B:283:GLU:HG2	1:B:307:PRO:HG3	1.95	0.47
1:A:251:ASP:HB3	1:A:254:CYS:HB3	1.97	0.47
1:B:130:PRO:HD2	1:B:196:PHE:O	2.15	0.46
1:A:93:ASP:OD1	1:A:97:LYS:NZ	2.49	0.46
1:A:152:LEU:HG	1:A:156:ILE:HD11	1.97	0.45
1:A:412:LYS:HD3	1:A:415:LYS:CE	2.46	0.44
1:A:353:THR:O	1:A:422:ARG:HB2	2.18	0.43
1:B:271:PHE:CZ	1:B:285:LEU:HD11	2.53	0.43
1:B:415:LYS:O	1:B:419:LYS:HG3	2.18	0.43
1:A:40:LEU:O	1:A:60:THR:HA	2.19	0.42
1:A:412:LYS:HA	1:A:415:LYS:HG2	2.01	0.42
1:A:286:ALA:HB1	1:A:310:PHE:CE1	2.54	0.42
1:A:44:VAL:O	1:A:48:ILE:HG13	2.20	0.42
1:A:274:MET:HE3	6:A:614:HOH:O	2.19	0.42
1:A:443:VAL:O	1:A:447:GLN:HG2	2.20	0.41
1:B:315:ASN:O	1:B:317:GLU:N	2.52	0.41
1:A:248:SER:C	1:A:250:ASP:H	2.24	0.41
1:A:382:VAL:HG12	1:A:405:VAL:CG2	2.51	0.41
1:A:135:PRO:HD3	1:A:345:TRP:CG	2.56	0.41
1:A:332:SER:HB3	1:A:354:PHE:CE1	2.55	0.40
1:B:34:LEU:HA	1:B:34:LEU:HD12	1.85	0.40
1:B:40:LEU:O	1:B:60:THR:HA	2.21	0.40
1:B:360:ALA:O	1:B:384:VAL:HG22	2.20	0.40
1:B:192:GLU:HG2	1:B:193:LYS:HG3	2.03	0.40
1:B:316:LYS:HG3	1:B:316:LYS:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	442/449 (98%)	429 (97%)	13 (3%)	0	100	100
1	B	442/449 (98%)	424 (96%)	18 (4%)	0	100	100
All	All	884/898 (98%)	853 (96%)	31 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	395/400 (99%)	388 (98%)	7 (2%)	59	63
1	B	395/400 (99%)	387 (98%)	8 (2%)	55	58
All	All	790/800 (99%)	775 (98%)	15 (2%)	57	61

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	101	SER
1	A	191	PHE
1	A	211	GLU
1	A	254	CYS
1	A	308	SER
1	A	422	ARG
1	A	427	LYS
1	B	89	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	103	ASN
1	B	174	SER
1	B	249	LYS
1	B	285	LEU
1	B	316	LYS
1	B	388	LYS
1	B	428	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	375	GLN
1	B	103	ASN
1	B	367	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BGC	C	1	1,2	11,11,12	0.98	0	15,15,17	1.34	2 (13%)
2	BGC	C	2	2	11,11,12	0.46	0	15,15,17	1.18	2 (13%)
2	BGC	D	1	1,2	11,11,12	0.88	0	15,15,17	1.36	3 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	D	2	2	11,11,12	0.44	0	15,15,17	1.05	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	C	1	1,2	-	0/2/19/22	0/1/1/1
2	BGC	C	2	2	-	0/2/19/22	0/1/1/1
2	BGC	D	1	1,2	-	0/2/19/22	0/1/1/1
2	BGC	D	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	BGC	C1-O5-C5	3.04	116.31	112.19
2	D	1	BGC	O3-C3-C2	-2.81	104.61	109.99
2	C	1	BGC	C3-C4-C5	2.52	114.74	110.24
2	C	1	BGC	O2-C2-C3	-2.50	105.12	110.14
2	D	1	BGC	O2-C2-C1	2.42	114.11	109.15
2	C	2	BGC	C1-C2-C3	-2.16	107.02	109.67
2	D	1	BGC	O5-C5-C6	2.14	110.56	107.20



There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	UDP	B	503	-	20,26,26	3.57	9 (45%)	25,40,40	1.17	3 (12%)
5	7WV	A	504	-	8,10,10	4.68	2 (25%)	9,13,13	1.08	1 (11%)
4	UDP	A	503	-	20,26,26	3.65	8 (40%)	25,40,40	1.21	3 (12%)
5	7WV	B	504	-	8,10,10	4.31	2 (25%)	9,13,13	1.01	0
3	BGC	A	501	1	11,11,12	1.17	1 (9%)	15,15,17	2.13	3 (20%)
3	BGC	B	501	1	11,11,12	1.19	1 (9%)	15,15,17	1.52	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	UDP	B	503	-	-	3/14/32/32	0/2/2/2
5	7WV	A	504	-	-	0/0/4/4	0/1/1/1
4	UDP	A	503	-	-	3/14/32/32	0/2/2/2
5	7WV	B	504	-	-	0/0/4/4	0/1/1/1
3	BGC	A	501	1	-	1/2/19/22	0/1/1/1
3	BGC	B	501	1	-	0/2/19/22	0/1/1/1

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	504	7WV	C05-C07	10.70	1.57	1.47
5	A	504	7WV	C05-C07	10.06	1.57	1.47
5	A	504	7WV	BR1-C04	8.24	2.08	1.89
4	A	503	UDP	C6-N1	7.88	1.45	1.35
4	B	503	UDP	C6-N1	6.88	1.44	1.35
4	B	503	UDP	C4-N3	6.86	1.45	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	503	UDP	C4-N3	6.65	1.44	1.33
4	B	503	UDP	O4'-C4'	6.16	1.58	1.45
4	A	503	UDP	O4'-C4'	5.98	1.58	1.45
4	B	503	UDP	C3'-C4'	-5.69	1.38	1.53
4	A	503	UDP	C3'-C4'	-5.63	1.38	1.53
5	B	504	7WV	BR1-C04	5.53	2.02	1.89
4	A	503	UDP	O4'-C1'	-5.51	1.33	1.41
4	B	503	UDP	O4'-C1'	-5.34	1.33	1.41
4	A	503	UDP	C6-C5	5.15	1.49	1.38
4	B	503	UDP	C6-C5	5.10	1.49	1.38
4	B	503	UDP	C2-N3	4.06	1.46	1.38
4	A	503	UDP	C2-N3	4.02	1.46	1.38
4	B	503	UDP	O3'-C3'	2.62	1.49	1.43
4	A	503	UDP	O3'-C3'	2.57	1.49	1.43
3	A	501	BGC	C6-C5	2.25	1.59	1.51
4	B	503	UDP	O2'-C2'	-2.09	1.38	1.43
3	B	501	BGC	C6-C5	2.06	1.58	1.51

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	BGC	C1-O5-C5	5.45	119.57	112.19
3	B	501	BGC	C1-C2-C3	4.67	115.40	109.67
3	A	501	BGC	C1-C2-C3	4.02	114.61	109.67
4	B	503	UDP	C2'-C3'-C4'	3.58	109.60	102.64
4	A	503	UDP	C2'-C3'-C4'	3.55	109.54	102.64
5	A	504	7WV	C06-C05-C07	-2.37	116.67	120.20
4	A	503	UDP	C5-C4-N3	-2.30	118.24	123.31
4	A	503	UDP	O3B-PB-O2B	2.14	115.83	107.64
4	B	503	UDP	C5-C4-N3	-2.05	118.80	123.31
3	A	501	BGC	C6-C5-C4	-2.02	108.27	113.00
4	B	503	UDP	O2'-C2'-C3'	2.00	118.30	111.82

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	503	UDP	PB-O3A-PA-O5'
4	B	503	UDP	PB-O3A-PA-O5'
4	B	503	UDP	O4'-C4'-C5'-O5'
4	A	503	UDP	O4'-C4'-C5'-O5'
4	B	503	UDP	C3'-C4'-C5'-O5'

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	503	UDP	C3'-C4'-C5'-O5'
3	A	501	BGC	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	444/449 (98%)	-0.01	8 (1%) 68 66	30, 50, 88, 118	0
1	B	444/449 (98%)	0.21	24 (5%) 25 24	36, 55, 98, 129	0
All	All	888/898 (98%)	0.10	32 (3%) 42 42	30, 53, 93, 129	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	52	LEU	5.3
1	B	253	PHE	3.9
1	A	253	PHE	3.7
1	B	250	ASP	3.6
1	B	310	PHE	3.6
1	B	5	ARG	3.4
1	A	311	LEU	3.3
1	B	4	LYS	3.3
1	B	313	THR	3.2
1	B	309	GLY	3.1
1	B	249	LYS	3.0
1	A	392	ILE	2.9
1	B	308	SER	2.9
1	B	314	VAL	2.7
1	A	314	VAL	2.7
1	B	49	ASN	2.6
1	A	316	LYS	2.5
1	B	191	PHE	2.4
1	B	317	GLU	2.4
1	B	318	LYS	2.3
1	B	294	PHE	2.3
1	B	311	LEU	2.3
1	B	307	PRO	2.3
1	B	193	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	309	GLY	2.2
1	B	257	TRP	2.1
1	A	317	GLU	2.1
1	B	189	ILE	2.1
1	B	51	ASP	2.1
1	B	312	GLU	2.1
1	A	315	ASN	2.1
1	B	389	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	BGC	D	2	11/12	0.93	0.10	50,57,60,61	0
2	BGC	C	2	11/12	0.96	0.10	36,40,46,50	0
2	BGC	D	1	11/12	0.97	0.09	44,52,61,64	0
2	BGC	C	1	11/12	0.97	0.11	35,39,44,57	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BGC	B	501	11/12	0.92	0.09	56,63,70,71	0
3	BGC	A	501	11/12	0.95	0.10	42,45,51,68	0
4	UDP	A	503	25/25	0.98	0.12	33,38,45,45	0
4	UDP	B	503	25/25	0.98	0.12	36,41,46,48	0
5	7WV	A	504	10/10	0.98	0.08	42,45,53,56	0
5	7WV	B	504	10/10	0.99	0.08	43,47,59,62	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers ⓘ

There are no such residues in this entry.