



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 6, 2021 – 12:11 PM JST

PDB ID : 7V25
Title : Crystal Structure of phthalate dioxygenase in complex with phthalate
Authors : Mahto, J.K.; Kumar, P.
Deposited on : 2021-08-07
Resolution : 2.74 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.24
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.24

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1271 (2.76-2.72)
Clashscore	141614	1322 (2.76-2.72)
Ramachandran outliers	138981	1297 (2.76-2.72)
Sidechain outliers	138945	1298 (2.76-2.72)
RSRZ outliers	127900	1243 (2.76-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	439	<div><div>2%</div><div><div></div><div>74%</div><div>17%</div><div>•</div><div>7%</div></div></div>
1	B	439	<div><div>5%</div><div><div></div><div>72%</div><div>22%</div><div>•</div><div>•</div></div></div>
1	C	439	<div><div>5%</div><div><div></div><div>73%</div><div>19%</div><div>•</div><div>7%</div></div></div>
1	D	439	<div><div>3%</div><div><div></div><div>68%</div><div>22%</div><div>•</div><div>8%</div></div></div>
1	E	439	<div><div>10%</div><div><div></div><div>69%</div><div>19%</div><div>•</div><div>10%</div></div></div>
1	F	439	<div><div>9%</div><div><div></div><div>65%</div><div>26%</div><div>•</div><div>7%</div></div></div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 20542 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Rieske (2Fe-2S) domain protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	423	Total	C	N	O	S	0	0	0
			3323	2094	588	617	24			
1	A	407	Total	C	N	O	S	0	0	0
			3213	2029	567	594	23			
1	C	407	Total	C	N	O	S	0	0	0
			3212	2029	567	593	23			
1	D	405	Total	C	N	O	S	0	0	0
			3203	2023	565	592	23			
1	E	395	Total	C	N	O	S	0	0	0
			3095	1952	545	574	24			
1	F	409	Total	C	N	O	S	0	0	0
			3228	2038	569	597	24			

- Molecule 2 is FE (II) ION (three-letter code: FE2) (formula: Fe) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Fe	0	0
			1	1		
2	A	1	Total	Fe	0	0
			1	1		
2	C	1	Total	Fe	0	0
			1	1		
2	D	1	Total	Fe	0	0
			1	1		
2	E	1	Total	Fe	0	0
			1	1		
2	F	1	Total	Fe	0	0
			1	1		

- Molecule 3 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe₂S₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	Fe	S	0	0
			4	2	2		
3	A	1	Total	Fe	S	0	0
			4	2	2		
3	C	1	Total	Fe	S	0	0
			4	2	2		
3	D	1	Total	Fe	S	0	0
			4	2	2		
3	E	1	Total	Fe	S	0	0
			4	2	2		
3	F	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 4 is PHTHALIC ACID (three-letter code: PHT) (formula: C₈H₆O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			12	8	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	208	Total	O	0	0
			208	208		
5	A	204	Total	O	0	0
			204	204		
5	C	208	Total	O	0	0
			208	208		
5	D	198	Total	O	0	0
			198	198		
5	E	200	Total	O	0	0
			200	200		
5	F	208	Total	O	0	0
			208	208		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Rieske (2Fe-2S) domain protein




• Molecule 1: Rieske (2Fe-2S) domain protein




• Molecule 1: Rieske (2Fe-2S) domain protein







• Molecule 1: Rieske (2Fe-2S) domain protein



• Molecule 1: Rieske (2Fe-2S) domain protein



• Molecule 1: Rieske (2Fe-2S) domain protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	179.03Å 122.18Å 162.94Å 90.00° 101.13° 90.00°	Depositor
Resolution (Å)	159.88 – 2.74 159.88 – 2.74	Depositor EDS
% Data completeness (in resolution range)	98.0 (159.88-2.74) 98.0 (159.88-2.74)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 2.73Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.207 , 0.283 0.207 , 0.283	Depositor DCC
R_{free} test set	4388 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	45.9	Xtriage
Anisotropy	0.101	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 45.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	20542	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PHT, FES, FE2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/3301	0.74	1/4497 (0.0%)
1	B	0.33	0/3414	0.75	1/4652 (0.0%)
1	C	0.33	0/3301	0.72	1/4498 (0.0%)
1	D	0.34	0/3292	0.72	0/4485
1	E	0.34	0/3176	0.73	0/4320
1	F	0.32	0/3317	0.72	0/4518
All	All	0.34	0/19801	0.73	3/26970 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	339	PHE	CB-CA-C	5.72	121.84	110.40
1	B	329	LYS	CB-CA-C	5.50	121.40	110.40
1	C	300	ARG	CG-CD-NE	-5.03	101.23	111.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3213	0	3096	47	0
1	B	3323	0	3210	75	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3212	0	3097	61	0
1	D	3203	0	3084	75	0
1	E	3095	0	2993	64	0
1	F	3228	0	3110	105	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	4	0	0	1	0
3	B	4	0	0	0	0
3	C	4	0	0	1	0
3	D	4	0	0	0	0
3	E	4	0	0	1	0
3	F	4	0	0	0	0
4	B	12	0	4	0	0
5	A	204	0	0	9	0
5	B	208	0	0	6	0
5	C	208	0	0	8	0
5	D	198	0	0	9	0
5	E	200	0	0	8	0
5	F	208	0	0	16	0
All	All	20542	0	18594	404	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 404 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2:MET:HE1	1:B:372:ARG:HD3	1.22	1.16
1:F:397:THR:HB	1:F:398:PRO:HD3	1.31	1.08
1:B:66:MET:CE	1:B:120:HIS:HD2	1.70	1.03
1:B:66:MET:HE3	1:B:120:HIS:HD2	1.27	0.99
1:E:54:VAL:HG22	1:E:68:GLU:HA	1.45	0.98

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	401/439 (91%)	368 (92%)	29 (7%)	4 (1%)	15	28
1	B	421/439 (96%)	386 (92%)	33 (8%)	2 (0%)	29	48
1	C	401/439 (91%)	369 (92%)	29 (7%)	3 (1%)	22	40
1	D	399/439 (91%)	367 (92%)	29 (7%)	3 (1%)	19	36
1	E	387/439 (88%)	350 (90%)	30 (8%)	7 (2%)	8	15
1	F	403/439 (92%)	368 (91%)	31 (8%)	4 (1%)	15	28
All	All	2412/2634 (92%)	2208 (92%)	181 (8%)	23 (1%)	15	28

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	236	ASN
1	F	15	GLY
1	A	74	ARG
1	E	409	LYS
1	B	198	ALA

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	337/363 (93%)	315 (94%)	22 (6%)	17	30
1	B	348/363 (96%)	329 (94%)	19 (6%)	21	37
1	C	337/363 (93%)	319 (95%)	18 (5%)	22	39

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	337/363 (93%)	314 (93%)	23 (7%)	16	28
1	E	326/363 (90%)	309 (95%)	17 (5%)	23	39
1	F	340/363 (94%)	322 (95%)	18 (5%)	22	39
All	All	2025/2178 (93%)	1908 (94%)	117 (6%)	20	35

5 of 117 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	402	SER
1	F	240	ASN
1	D	235	SER
1	F	227	TYR
1	F	28	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	263	ASN
1	F	260	ASN
1	D	288	GLN
1	F	358	HIS
1	E	342	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 13 ligands modelled in this entry, 6 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	FES	C	502	1	0,4,4	-	-	-		
4	PHT	B	503	-	8,12,12	2.58	2 (25%)	8,16,16	1.95	2 (25%)
3	FES	B	502	1	0,4,4	-	-	-		
3	FES	F	502	1	0,4,4	-	-	-		
3	FES	E	502	1	0,4,4	-	-	-		
3	FES	A	502	1	0,4,4	-	-	-		
3	FES	D	502	1	0,4,4	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PHT	B	503	-	-	0/0/8/8	0/1/1/1
3	FES	C	502	1	-	-	0/1/1/1
3	FES	B	502	1	-	-	0/1/1/1
3	FES	F	502	1	-	-	0/1/1/1
3	FES	E	502	1	-	-	0/1/1/1
3	FES	A	502	1	-	-	0/1/1/1
3	FES	D	502	1	-	-	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	503	PHT	C2-C10	5.76	1.53	1.47
4	B	503	PHT	C3-C7	4.47	1.51	1.47

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	503	PHT	C4-C3-C7	-4.05	114.17	120.20
4	B	503	PHT	C1-C2-C10	-3.45	115.07	120.20

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.


3 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	502	FES	1	0
3	E	502	FES	1	0
3	A	502	FES	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	407/439 (92%)	0.24	9 (2%) 62 69	22, 40, 75, 116	0
1	B	423/439 (96%)	0.45	21 (4%) 28 32	21, 40, 104, 159	0
1	C	407/439 (92%)	0.39	22 (5%) 25 29	23, 40, 98, 136	0
1	D	405/439 (92%)	0.37	14 (3%) 44 49	25, 44, 91, 128	0
1	E	395/439 (89%)	0.59	44 (11%) 5 5	24, 43, 132, 187	0
1	F	409/439 (93%)	0.68	40 (9%) 7 7	28, 49, 114, 132	0
All	All	2446/2634 (92%)	0.45	150 (6%) 21 24	21, 43, 107, 187	0

The worst 5 of 150 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	399	THR	9.5
1	C	111	ALA	8.4
1	E	400	VAL	7.4
1	B	201	THR	6.4
1	E	234	LEU	6.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	PHT	B	503	12/12	0.84	0.26	62,75,84,88	0
2	FE2	F	501	1/1	0.96	0.12	45,45,45,45	0
2	FE2	E	501	1/1	0.96	0.16	40,40,40,40	0
3	FES	C	502	4/4	0.98	0.11	39,41,41,50	0
2	FE2	D	501	1/1	0.98	0.12	34,34,34,34	0
2	FE2	C	501	1/1	0.99	0.14	30,30,30,30	0
3	FES	A	502	4/4	0.99	0.12	30,33,36,40	0
2	FE2	B	501	1/1	0.99	0.12	36,36,36,36	0
3	FES	D	502	4/4	0.99	0.14	35,38,40,44	0
3	FES	E	502	4/4	0.99	0.14	25,27,30,32	0
3	FES	F	502	4/4	0.99	0.12	33,35,36,45	0
2	FE2	A	501	1/1	0.99	0.17	27,27,27,27	0
3	FES	B	502	4/4	1.00	0.12	29,29,33,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around PHT B 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE2 F 501:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE2 E 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around FES C 502:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FE2 D 501:**




$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)








Electron density around FES A 502:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FE2 B 501:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FES D 502:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FES E 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FES F 502:


$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FE2 A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.