



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 02:29 am BST

PDB ID : 5V3D
Title : Crystal structure of fosfomycin resistance protein from *Klebsiella pneumoniae* with bound fosfomycin
Authors : Klontz, E.; Guenther, S.; Silverstein, Z.; Sundberg, E.
Deposited on : 2017-03-07
Resolution : 1.54 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2556 (1.56-1.52)
Clashscore	141614	2634 (1.56-1.52)
Ramachandran outliers	138981	2580 (1.56-1.52)
Sidechain outliers	138945	2577 (1.56-1.52)
RSRZ outliers	127900	2524 (1.56-1.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	145	<div> <div>5%</div> <div> <div></div> <div>87%</div> <div>8%</div> <div>5%</div> </div> </div>
1	B	145	<div> <div>10%</div> <div> <div></div> <div>83%</div> <div>10%</div> <div>6%</div> </div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 2480 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fosfomycin resistance protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	138	Total	C	N	O	S	0	2	0
			1094	698	190	199	7			
1	B	137	Total	C	N	O	S	0	1	0
			1078	689	185	197	7			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	140	HIS	-	expression tag	UNP W8UNW6
A	141	HIS	-	expression tag	UNP W8UNW6
A	142	HIS	-	expression tag	UNP W8UNW6
A	143	HIS	-	expression tag	UNP W8UNW6
A	144	HIS	-	expression tag	UNP W8UNW6
A	145	HIS	-	expression tag	UNP W8UNW6
B	140	HIS	-	expression tag	UNP W8UNW6
B	141	HIS	-	expression tag	UNP W8UNW6
B	142	HIS	-	expression tag	UNP W8UNW6
B	143	HIS	-	expression tag	UNP W8UNW6
B	144	HIS	-	expression tag	UNP W8UNW6
B	145	HIS	-	expression tag	UNP W8UNW6

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mn	0	0
			2	2		

- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	K	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total K 1 1	0	0

- Molecule 4 is FOSFOMYCIN (three-letter code: FCN) (formula: $C_3H_7O_4P$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O P 8 3 4 1	0	0
4	A	1	Total C O P 8 3 4 1	0	0

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	152	Total 152	O 152	0	0
7	B	125	Total 125	O 125	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fosfomycin resistance protein



- Molecule 1: Fosfomycin resistance protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	40.07Å 47.16Å 149.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.29 – 1.54 29.29 – 1.54	Depositor EDS
% Data completeness (in resolution range)	93.5 (29.29-1.54) 96.7 (29.29-1.54)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 1.54Å)	Xtriage
Refinement program	PHENIX 1.10.1 _2155	Depositor
R, R_{free}	0.168 , 0.206 0.168 , 0.204	Depositor DCC
R_{free} test set	2069 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	17.6	Xtriage
Anisotropy	1.027	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2480	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, K, MN, EDO, FCN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/1124	0.54	0/1526
1	B	0.49	2/1108 (0.2%)	0.55	0/1505
All	All	0.43	2/2232 (0.1%)	0.55	0/3031

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	74	GLU	CD-OE1	-7.76	1.17	1.25
1	B	74	GLU	CD-OE2	-6.99	1.18	1.25

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1094	0	1050	8	0
1	B	1078	0	1036	11	0
2	A	2	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	16	0	9	4	0
5	B	7	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	4	0	6	1	0
7	A	152	0	0	1	3
7	B	125	0	0	0	1
All	All	2480	0	2109	17	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:124:ALA:O	1:B:128:GLU:OE1	2.05	0.74
4:A:203:FCN:H2	1:B:48[B]:CYS:SG	2.34	0.66
1:B:16:ALA:HB2	1:B:33:ARG:HH22	1.74	0.52
1:A:5:LEU:HD22	1:B:5:LEU:HD22	1.92	0.52
1:A:35:ASP:OD2	1:B:132:LYS:NZ	2.37	0.51
1:A:48[B]:CYS:SG	4:A:205:FCN:H2	2.52	0.49
1:A:96:ARG:NH1	7:A:303:HOH:O	2.40	0.48
1:A:101:SER:HA	1:A:114:LEU:O	2.14	0.48
1:B:101:SER:HA	1:B:114:LEU:O	2.16	0.46
1:A:16:ALA:HB2	1:A:33:ARG:HH22	1.81	0.45
4:A:205:FCN:H32	1:B:122:ARG:HH12	1.81	0.44
1:A:131:TYR:HB2	1:A:134:MET:HE3	2.02	0.42
1:B:55:ARG:HA	1:B:55:ARG:HD2	1.91	0.42
1:B:16:ALA:HB3	1:B:17:PRO:HD3	2.02	0.42
1:B:98:GLU:OE2	1:B:121:GLN:NE2	2.26	0.41
1:B:5:LEU:HB3	6:B:203:EDO:H12	2.02	0.41
1:A:122:ARG:HH12	4:A:203:FCN:H32	1.86	0.41

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:449:HOH:O	7:A:451:HOH:O[4_545]	2.05	0.15
7:A:348:HOH:O	7:B:380:HOH:O[3_554]	2.08	0.12
7:A:314:HOH:O	7:A:371:HOH:O[4_545]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	138/145 (95%)	137 (99%)	1 (1%)	0	100	100
1	B	136/145 (94%)	136 (100%)	0	0	100	100
All	All	274/290 (94%)	273 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/119 (95%)	112 (99%)	1 (1%)	78	60
1	B	112/119 (94%)	109 (97%)	3 (3%)	44	15
All	All	225/238 (94%)	221 (98%)	4 (2%)	59	29

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	106	ASP
1	B	74	GLU
1	B	106	ASP
1	B	132	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	FCN	A	203	2	5,8,8	1.54	1 (20%)	5,13,13	1.94	2 (40%)
6	EDO	B	203	-	3,3,3	0.52	0	2,2,2	0.08	0
4	FCN	A	205	2	5,8,8	1.55	2 (40%)	5,13,13	1.40	1 (20%)
5	PEG	B	201	-	6,6,6	0.53	0	5,5,5	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FCN	A	203	2	-	0/0/11/11	0/1/1/1
6	EDO	B	203	-	-	0/1/1/1	-
4	FCN	A	205	2	-	0/0/11/11	0/1/1/1
5	PEG	B	201	-	-	1/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	205	FCN	P-O2P	-2.09	1.51	1.54
4	A	203	FCN	P-O3P	-2.04	1.51	1.54
4	A	205	FCN	P-O1P	2.01	1.53	1.49

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	A	203	FCN	O1P-P-C1	-3.34	105.87	113.37
4	A	205	FCN	O3P-P-O1P	-2.58	106.96	113.45
4	A	203	FCN	O-C2-C3	2.44	121.13	116.61

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	201	PEG	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	203	FCN	2	0
6	B	203	EDO	1	0
4	A	205	FCN	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	138/145 (95%)	0.10	7 (5%) 28 31	17, 26, 43, 57	0
1	B	137/145 (94%)	0.39	15 (10%) 5 5	18, 29, 50, 63	0
All	All	275/290 (94%)	0.24	22 (8%) 12 14	17, 28, 47, 63	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	75	ALA	5.1
1	A	1	MET	3.5
1	B	10	LEU	3.3
1	B	136	PHE	2.9
1	B	8	LEU	2.8
1	B	128	GLU	2.7
1	B	125	ALA	2.7
1	A	138	GLU	2.7
1	B	74	GLU	2.6
1	B	1	MET	2.6
1	A	137	PHE	2.6
1	A	68	TYR	2.5
1	B	120	ALA	2.4
1	A	112	LEU	2.4
1	B	48[A]	CYS	2.4
1	A	69	ALA	2.4
1	B	47	LEU	2.3
1	B	124	ALA	2.2
1	B	132	LYS	2.2
1	A	8	LEU	2.2
1	B	54	GLN	2.1
1	B	137	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	PEG	B	201	7/7	0.84	0.22	29,31,34,35	0
6	EDO	B	203	4/4	0.91	0.13	26,27,28,31	0
4	FCN	A	205	8/8	0.94	0.13	36,39,43,45	0
4	FCN	A	203	8/8	0.98	0.09	24,28,38,42	0
3	K	B	202	1/1	0.98	0.07	41,41,41,41	0
2	MN	A	204	1/1	0.99	0.14	36,36,36,36	0
3	K	A	202	1/1	1.00	0.04	22,22,22,22	0
2	MN	A	201	1/1	1.00	0.10	17,17,17,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.