



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 09:25 pm BST

PDB ID : 1V4A
Title : Structure of the N-terminal Domain of Escherichia coli Glutamine Synthetase adenylyltransferase
Authors : Xu, Y.; Zhang, R.; Joachimiak, A.; Carr, P.D.; Ollis, D.L.; Vasudevan, S.G.
Deposited on : 2003-11-12
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	440	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3541 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate-ammonia-ligase adenylyltransferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	429	Total	C	N	O	S	Se	0	0	0
			3370	2146	593	615	3	13			

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	UNP P30870
A	87	MSE	MET	MODIFIED RESIDUE	UNP P30870
A	98	MSE	MET	MODIFIED RESIDUE	UNP P30870
A	160	MSE	MET	MODIFIED RESIDUE	UNP P30870
A	202	MSE	MET	MODIFIED RESIDUE	UNP P30870
A	224	MSE	MET	MODIFIED RESIDUE	UNP P30870
A	258	MSE	MET	MODIFIED RESIDUE	UNP P30870
A	264	MSE	MET	MODIFIED RESIDUE	UNP P30870
A	278	MSE	MET	MODIFIED RESIDUE	UNP P30870
A	299	MSE	MET	MODIFIED RESIDUE	UNP P30870
A	302	MSE	MET	MODIFIED RESIDUE	UNP P30870
A	409	MSE	MET	MODIFIED RESIDUE	UNP P30870
A	425	MSE	MET	MODIFIED RESIDUE	UNP P30870


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	171	Total	O	0	0
			171	171		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutamate-ammonia-ligase adenylyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	116.84Å 116.84Å 67.74Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.00 44.24 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-2.00) 99.9 (44.24-2.00)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.10 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.228 , 0.269 0.239 , 0.277	Depositor DCC
R_{free} test set	1823 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	27.9	Xtriage
Anisotropy	0.228	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3541	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	1/3429 (0.0%)	0.69	4/4636 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	258	MSE	SE-CE	-6.56	1.56	1.95

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	266	ASP	CB-CG-OD2	5.71	123.44	118.30
1	A	211	ASP	CB-CG-OD2	5.25	123.03	118.30
1	A	40	ASP	CB-CG-OD2	5.17	122.96	118.30
1	A	223	ASP	CB-CG-OD2	5.08	122.87	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3370	0	3275	27	0
2	A	171	0	0	0	0
All	All	3541	0	3275	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:295:SER:C	1:A:299:MSE:HE3	2.07	0.75
1:A:420:ALA:O	1:A:424:HIS:HD2	1.76	0.68
1:A:160:MSE:HE1	1:A:258:MSE:SE	2.45	0.66
1:A:147:ASN:O	1:A:191:ARG:N	2.30	0.65
1:A:124:GLU:HG3	1:A:264:MSE:HG3	1.82	0.60
1:A:372:VAL:HG12	1:A:424:HIS:HE1	1.67	0.59
1:A:316:ILE:HG23	1:A:377:LEU:HD22	1.89	0.54
1:A:295:SER:OG	1:A:299:MSE:HE2	2.08	0.53
1:A:295:SER:OG	1:A:299:MSE:CE	2.56	0.53
1:A:243:GLU:HG3	1:A:278:MSE:HE2	1.92	0.52
1:A:280:ARG:NH2	1:A:345:GLN:HE22	2.08	0.52
1:A:124:GLU:HB2	1:A:264:MSE:CE	2.41	0.51
1:A:372:VAL:HG12	1:A:424:HIS:CE1	2.45	0.50
1:A:46:VAL:HG21	1:A:53:LEU:CB	2.44	0.47
1:A:223:ASP:OD1	1:A:225:ARG:HD3	2.15	0.46
1:A:131:ARG:HG3	1:A:157:ILE:HD11	1.98	0.45
1:A:280:ARG:CZ	1:A:345:GLN:HE22	2.29	0.45
1:A:139:CYS:SG	1:A:145:PRO:HA	2.56	0.45
1:A:2:LYS:O	1:A:407:TRP:NE1	2.49	0.45
1:A:384:LEU:HD13	1:A:392:THR:HG23	1.99	0.44
1:A:295:SER:O	1:A:299:MSE:HE3	2.17	0.44
1:A:280:ARG:HB3	1:A:281:PRO:HD3	2.00	0.42
1:A:131:ARG:NH1	1:A:132:ASP:OD1	2.53	0.42
1:A:1:MSE:HE1	1:A:31:GLN:CG	2.50	0.42
1:A:142:TRP:O	1:A:144:THR:N	2.53	0.41
1:A:146:CYS:HB2	1:A:195:ASN:OD1	2.21	0.41
1:A:124:GLU:HB2	1:A:264:MSE:HE3	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	425/440 (97%)	411 (97%)	9 (2%)	5 (1%)	13 7

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	18	ARG
1	A	143	GLY
1	A	144	THR
1	A	3	PRO
1	A	181	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	341/362 (94%)	315 (92%)	26 (8%)	13 8

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	SER
1	A	33	LYS
1	A	47	ILE
1	A	51	GLU
1	A	73	LEU
1	A	86	LEU
1	A	140	ARG
1	A	147	ASN
1	A	156	LEU
1	A	167	GLU
1	A	168	LEU
1	A	191	ARG
1	A	211	ASP
1	A	225	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	232	SER
1	A	235	LEU
1	A	242	LEU
1	A	264	MSE
1	A	275	LEU
1	A	347	ARG
1	A	353	LEU
1	A	360	HIS
1	A	364	GLU
1	A	370	LEU
1	A	384	LEU
1	A	400	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	345	GLN
1	A	424	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	416/440 (94%)	0.10	14 (3%) 45 44	14, 31, 57, 71	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	312	LEU	4.8
1	A	144	THR	4.8
1	A	191	ARG	4.4
1	A	193	LEU	4.2
1	A	143	GLY	3.7
1	A	415	PRO	3.4
1	A	27	SER	3.4
1	A	412	ALA	3.3
1	A	287	TYR	2.9
1	A	58	SER	2.9
1	A	142	TRP	2.8
1	A	79	ASN	2.4
1	A	78	CYS	2.1
1	A	56	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.