



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 7, 2020 – 01:04 AM BST

PDB ID : 6V44
Title : The crystal structure of hemagglutinin from swine influenza virus A/swine/Missouri/A01727926/2015
Authors : Yang, H.; Stevens, J.
Deposited on : 2019-11-27
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	332	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> % </div> <div style="width: 100%; height: 10px; background-color: green; position: relative;"> 84% 12% • </div> </div>
1	C	332	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> % </div> <div style="width: 100%; height: 10px; background-color: green; position: relative;"> 81% 15% • • </div> </div>
1	E	332	<div> <div style="width: 100%; height: 10px; background-color: green; position: relative;"> 84% 12% • </div> </div>
2	B	186	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> 3% </div> <div style="width: 100%; height: 10px; background-color: green; position: relative;"> 75% 16% • 7% </div> </div>
2	D	186	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> 3% </div> <div style="width: 100%; height: 10px; background-color: green; position: relative;"> 80% 12% • 7% </div> </div>
2	F	186	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> 5% </div> <div style="width: 100%; height: 10px; background-color: green; position: relative;"> 83% 9% • 7% </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	6	 17%83%
3	H	6	 17%83%
3	I	6	 83%17%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 12422 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin HA1 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	320	Total	C	N	O	S	0	0	0
			2487	1562	440	474	11			
1	C	320	Total	C	N	O	S	0	0	0
			2487	1562	440	474	11			
1	E	320	Total	C	N	O	S	0	0	0
			2487	1562	440	474	11			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	ALA	-	expression tag	UNP A0A140D8S6
A	-3	ASP	-	expression tag	UNP A0A140D8S6
A	-2	LEU	-	expression tag	UNP A0A140D8S6
A	-1	GLY	-	expression tag	UNP A0A140D8S6
C	-4	ALA	-	expression tag	UNP A0A140D8S6
C	-3	ASP	-	expression tag	UNP A0A140D8S6
C	-2	LEU	-	expression tag	UNP A0A140D8S6
C	-1	GLY	-	expression tag	UNP A0A140D8S6
E	-4	ALA	-	expression tag	UNP A0A140D8S6
E	-3	ASP	-	expression tag	UNP A0A140D8S6
E	-2	LEU	-	expression tag	UNP A0A140D8S6
E	-1	GLY	-	expression tag	UNP A0A140D8S6


- Molecule 2 is a protein called Hemagglutinin HA2 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	173	Total	C	N	O	S	0	0	0
			1415	878	251	282	4			
2	D	173	Total	C	N	O	S	0	0	0
			1415	878	251	282	4			
2	F	173	Total	C	N	O	S	0	0	0
			1415	878	251	282	4			

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	175	SER	-	expression tag	UNP A0A140D8S6
B	176	GLY	-	expression tag	UNP A0A140D8S6
B	177	ARG	-	expression tag	UNP A0A140D8S6
B	178	SER	-	expression tag	UNP A0A140D8S6
B	179	GLY	-	expression tag	UNP A0A140D8S6
B	180	ARG	-	expression tag	UNP A0A140D8S6
B	181	LEU	-	expression tag	UNP A0A140D8S6
B	182	VAL	-	expression tag	UNP A0A140D8S6
B	183	PRO	-	expression tag	UNP A0A140D8S6
B	184	ARG	-	expression tag	UNP A0A140D8S6
B	185	GLY	-	expression tag	UNP A0A140D8S6
B	186	SER	-	expression tag	UNP A0A140D8S6
D	175	SER	-	expression tag	UNP A0A140D8S6
D	176	GLY	-	expression tag	UNP A0A140D8S6
D	177	ARG	-	expression tag	UNP A0A140D8S6
D	178	SER	-	expression tag	UNP A0A140D8S6
D	179	GLY	-	expression tag	UNP A0A140D8S6
D	180	ARG	-	expression tag	UNP A0A140D8S6
D	181	LEU	-	expression tag	UNP A0A140D8S6
D	182	VAL	-	expression tag	UNP A0A140D8S6
D	183	PRO	-	expression tag	UNP A0A140D8S6
D	184	ARG	-	expression tag	UNP A0A140D8S6
D	185	GLY	-	expression tag	UNP A0A140D8S6
D	186	SER	-	expression tag	UNP A0A140D8S6
F	175	SER	-	expression tag	UNP A0A140D8S6
F	176	GLY	-	expression tag	UNP A0A140D8S6
F	177	ARG	-	expression tag	UNP A0A140D8S6
F	178	SER	-	expression tag	UNP A0A140D8S6
F	179	GLY	-	expression tag	UNP A0A140D8S6
F	180	ARG	-	expression tag	UNP A0A140D8S6
F	181	LEU	-	expression tag	UNP A0A140D8S6
F	182	VAL	-	expression tag	UNP A0A140D8S6
F	183	PRO	-	expression tag	UNP A0A140D8S6
F	184	ARG	-	expression tag	UNP A0A140D8S6
F	185	GLY	-	expression tag	UNP A0A140D8S6
F	186	SER	-	expression tag	UNP A0A140D8S6

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	G	6	Total	C	N	O	0	0	0
			72	40	2	30			
3	H	6	Total	C	N	O	0	0	0
			72	40	2	30			
3	I	6	Total	C	N	O	0	0	0
			72	40	2	30			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	E	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	83	Total	O	0	0
			83	83		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	61	Total 61	O 61	0	0
5	C	99	Total 99	O 99	0	0
5	D	55	Total 55	O 55	0	0
5	E	94	Total 94	O 94	0	0
5	F	66	Total 66	O 66	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hemagglutinin HA1 chain




- Molecule 1: Hemagglutinin HA1 chain




- Molecule 1: Hemagglutinin HA1 chain





- Molecule 2: Hemagglutinin HA2 chain








- Molecule 2: Hemagglutinin HA2 chain




- Molecule 2: Hemagglutinin HA2 chain




- Molecule 3: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	68.97Å 240.00Å 68.95Å 90.00° 119.86° 90.00°	Depositor
Resolution (Å)	33.21 – 2.20 33.21 – 2.20	Depositor EDS
% Data completeness (in resolution range)	97.2 (33.21-2.20) 96.4 (33.21-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.94 (at 2.20Å)	Xtriage
Refinement program	PHENIX 1.14 _3260	Depositor
R, R_{free}	0.203 , 0.244 0.203 , 0.244	Depositor DCC
R_{free} test set	4762 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	45.7	Xtriage
Anisotropy	0.369	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 24.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.357 for l,k,-h-l 0.357 for -h-l,k,h 0.028 for -h-l,-k,l 0.021 for h,-k,-h-l 0.019 for l,-k,h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12422	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.44	0/2546	0.67	1/3472 (0.0%)
1	C	0.46	0/2546	0.71	1/3472 (0.0%)
1	E	0.44	0/2546	0.66	1/3472 (0.0%)
2	B	0.54	1/1439 (0.1%)	0.71	3/1938 (0.2%)
2	D	0.56	1/1439 (0.1%)	0.68	2/1938 (0.1%)
2	F	0.46	0/1439	0.68	2/1938 (0.1%)
All	All	0.48	2/11955 (0.0%)	0.69	10/16230 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	30	GLU	CG-CD	9.12	1.65	1.51
2	B	69	GLU	CG-CD	5.63	1.60	1.51

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	260	GLN	CA-CB-CG	-9.74	91.96	113.40
2	F	17	LEU	CB-CG-CD1	6.51	122.07	111.00
2	F	71	GLU	CA-CB-CG	-6.17	99.83	113.40
2	B	69	GLU	CB-CA-C	6.11	122.62	110.40
2	D	86	ASP	CB-CG-OD1	5.41	123.17	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2487	0	2433	34	0
1	C	2487	0	2433	53	0
1	E	2487	0	2433	40	0
2	B	1415	0	1333	35	0
2	D	1415	0	1333	30	0
2	F	1415	0	1333	15	0
3	G	72	0	61	1	0
3	H	72	0	61	1	0
3	I	72	0	61	3	0
4	A	14	0	13	0	0
4	C	14	0	13	0	0
4	E	14	0	13	0	0
5	A	83	0	0	2	0
5	B	61	0	0	6	0
5	C	99	0	0	10	0
5	D	55	0	0	4	0
5	E	94	0	0	9	0
5	F	66	0	0	1	0
All	All	12422	0	11520	156	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 156 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:25:ARG:HG3	2:F:34:THR:HG22	1.46	0.95
1:C:152:LYS:NZ	5:C:503:HOH:O	2.00	0.94
2:B:128:GLU:OE1	5:B:202:HOH:O	1.95	0.85
2:B:11:GLU:OE1	5:B:201:HOH:O	1.93	0.84
2:D:153:ARG:NH1	5:D:201:HOH:O	2.11	0.82

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	318/332 (96%)	301 (95%)	16 (5%)	1 (0%)	41	46
1	C	318/332 (96%)	302 (95%)	15 (5%)	1 (0%)	41	46
1	E	318/332 (96%)	303 (95%)	14 (4%)	1 (0%)	41	46
2	B	171/186 (92%)	159 (93%)	10 (6%)	2 (1%)	13	10
2	D	171/186 (92%)	160 (94%)	9 (5%)	2 (1%)	13	10
2	F	171/186 (92%)	159 (93%)	10 (6%)	2 (1%)	13	10
All	All	1467/1554 (94%)	1384 (94%)	74 (5%)	9 (1%)	25	26

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	61	ASP
1	C	61	ASP
1	E	61	ASP
2	B	60	ASN
2	D	60	ASN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	281/290 (97%)	279 (99%)	2 (1%)	84	91
1	C	281/290 (97%)	277 (99%)	4 (1%)	67	80
1	E	281/290 (97%)	276 (98%)	5 (2%)	59	72

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	149/159 (94%)	146 (98%)	3 (2%)	55	69
2	D	149/159 (94%)	148 (99%)	1 (1%)	84	91
2	F	149/159 (94%)	147 (99%)	2 (1%)	69	81
All	All	1290/1347 (96%)	1273 (99%)	17 (1%)	69	81

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	262	LYS
1	C	266	LEU
1	E	239	LEU
1	C	239	LEU
1	E	260	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	53	ASN
2	B	72	GLN
2	D	72	GLN
1	E	14	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

18 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	G	1	1,3	14,14,15	0.35	0	17,19,21	0.49	0
3	NAG	G	2	3	14,14,15	0.63	0	17,19,21	0.60	0
3	BMA	G	3	3	11,11,12	1.00	1 (9%)	15,15,17	1.13	2 (13%)
3	MAN	G	4	3	11,11,12	1.19	1 (9%)	15,15,17	1.79	3 (20%)
3	MAN	G	5	3	11,11,12	1.72	3 (27%)	15,15,17	2.80	6 (40%)
3	MAN	G	6	3	11,11,12	1.99	2 (18%)	15,15,17	1.95	2 (13%)
3	NAG	H	1	1,3	14,14,15	0.36	0	17,19,21	0.52	0
3	NAG	H	2	3	14,14,15	0.56	0	17,19,21	0.59	0
3	BMA	H	3	3	11,11,12	0.95	1 (9%)	15,15,17	1.14	2 (13%)
3	MAN	H	4	3	11,11,12	1.10	1 (9%)	15,15,17	1.63	4 (26%)
3	MAN	H	5	3	11,11,12	1.77	2 (18%)	15,15,17	2.68	5 (33%)
3	MAN	H	6	3	11,11,12	1.89	2 (18%)	15,15,17	1.96	3 (20%)
3	NAG	I	1	1,3	14,14,15	0.30	0	17,19,21	0.74	0
3	NAG	I	2	3	14,14,15	0.51	0	17,19,21	0.63	0
3	BMA	I	3	3	11,11,12	1.46	2 (18%)	15,15,17	1.11	2 (13%)
3	MAN	I	4	3	11,11,12	0.94	0	15,15,17	2.10	3 (20%)
3	MAN	I	5	3	11,11,12	1.68	3 (27%)	15,15,17	2.78	5 (33%)
3	MAN	I	6	3	11,11,12	2.17	3 (27%)	15,15,17	1.70	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	G	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	G	2	3	-	0/6/23/26	0/1/1/1
3	BMA	G	3	3	-	2/2/19/22	0/1/1/1
3	MAN	G	4	3	-	0/2/19/22	0/1/1/1
3	MAN	G	5	3	-	2/2/19/22	0/1/1/1
3	MAN	G	6	3	-	0/2/19/22	0/1/1/1
3	NAG	H	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	H	2	3	-	0/6/23/26	0/1/1/1
3	BMA	H	3	3	-	2/2/19/22	0/1/1/1
3	MAN	H	4	3	-	0/2/19/22	0/1/1/1
3	MAN	H	5	3	-	2/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAN	H	6	3	-	0/2/19/22	0/1/1/1
3	NAG	I	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	I	2	3	-	2/6/23/26	0/1/1/1
3	BMA	I	3	3	-	2/2/19/22	0/1/1/1
3	MAN	I	4	3	-	0/2/19/22	0/1/1/1
3	MAN	I	5	3	-	2/2/19/22	0/1/1/1
3	MAN	I	6	3	-	0/2/19/22	0/1/1/1

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	I	6	MAN	C1-C2	4.61	1.62	1.52
3	H	6	MAN	C2-C3	4.49	1.59	1.52
3	G	6	MAN	C2-C3	4.32	1.58	1.52
3	G	6	MAN	C1-C2	4.18	1.61	1.52
3	I	6	MAN	O5-C1	3.98	1.50	1.43

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	I	5	MAN	C1-O5-C5	6.57	121.10	112.19
3	I	4	MAN	O2-C2-C3	-6.44	97.24	110.14
3	G	5	MAN	C1-C2-C3	6.12	117.19	109.67
3	I	5	MAN	C1-C2-C3	6.10	117.17	109.67
3	H	5	MAN	C1-C2-C3	5.88	116.89	109.67

There are no chirality outliers.

5 of 14 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	I	5	MAN	O5-C5-C6-O6
3	H	5	MAN	O5-C5-C6-O6
3	G	5	MAN	O5-C5-C6-O6
3	H	5	MAN	C4-C5-C6-O6
3	I	2	NAG	O5-C5-C6-O6





There are no ring outliers.


5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	2	NAG	1	0
3	H	2	NAG	1	0
3	G	2	NAG	1	0
3	I	3	BMA	1	0
3	I	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



Oligosaccharide Chain H**Bond lengths****Bond angles****Torsions****Rings**



5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	E	407	1	14,14,15	0.21	0	17,19,21	0.66	1 (5%)
4	NAG	A	407	1	14,14,15	0.36	0	17,19,21	0.67	1 (5%)
4	NAG	C	407	1	14,14,15	0.58	0	17,19,21	0.68	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	E	407	1	-	1/6/23/26	0/1/1/1
4	NAG	A	407	1	-	0/6/23/26	0/1/1/1
4	NAG	C	407	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	407	NAG	C1-O5-C5	2.48	115.56	112.19
4	A	407	NAG	C1-O5-C5	2.40	115.44	112.19
4	E	407	NAG	C1-O5-C5	2.32	115.33	112.19

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	E	407	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	320/332 (96%)	-0.03	3 (0%) 84 83	31, 47, 66, 81	0
1	C	320/332 (96%)	-0.04	2 (0%) 89 88	32, 47, 67, 82	0
1	E	320/332 (96%)	-0.03	0 100 100	33, 47, 65, 75	0
2	B	173/186 (93%)	-0.00	6 (3%) 44 42	35, 49, 85, 141	0
2	D	173/186 (93%)	0.16	6 (3%) 44 42	34, 50, 83, 144	0
2	F	173/186 (93%)	0.04	9 (5%) 27 26	36, 50, 86, 155	0
All	All	1479/1554 (95%)	0.00	26 (1%) 68 66	31, 48, 70, 155	0

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	58	LYS	6.8
2	D	56	ILE	6.2
2	F	173	ILE	4.1
2	B	58	LYS	3.8
2	B	60	ASN	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MAN	I	5	11/12	0.72	0.23	79,97,104,108	0
3	MAN	H	5	11/12	0.72	0.20	81,93,100,106	0
3	MAN	G	6	11/12	0.76	0.15	78,88,93,94	0
3	MAN	H	6	11/12	0.79	0.16	82,91,98,105	0
3	MAN	G	5	11/12	0.81	0.16	68,86,96,101	0
3	MAN	I	6	11/12	0.82	0.17	85,93,105,106	0
3	MAN	I	4	11/12	0.87	0.14	53,59,66,85	0
3	NAG	I	1	14/15	0.89	0.14	45,56,70,72	0
3	BMA	I	3	11/12	0.90	0.14	42,51,62,68	0
3	NAG	G	1	14/15	0.91	0.12	47,56,72,73	0
3	BMA	G	3	11/12	0.91	0.11	48,59,68,73	0
3	NAG	I	2	14/15	0.91	0.17	43,51,56,58	0
3	NAG	G	2	14/15	0.92	0.14	41,51,58,63	0
3	NAG	H	1	14/15	0.92	0.19	46,54,70,72	0
3	MAN	H	4	11/12	0.92	0.11	49,55,61,75	0
3	BMA	H	3	11/12	0.94	0.12	48,53,63,73	0
3	NAG	H	2	14/15	0.94	0.18	43,49,56,63	0
3	MAN	G	4	11/12	0.94	0.12	50,55,63,70	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain G:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	C	407	14/15	0.86	0.16	55,70,79,80	0
4	NAG	A	407	14/15	0.89	0.12	55,65,70,76	0
4	NAG	E	407	14/15	0.92	0.13	49,66,78,78	0

6.5 Other polymers [i](#)

There are no such residues in this entry.