



# Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 09:23 am BST

PDB ID : 1V6S  
Title : Crystal Structure of Phosphoglycerate Kinase from *Thermus thermophilus* HB8  
Authors : Mizutani, H.; Kunishima, N.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2003-12-04  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	390	<div> <div>6%</div> <div> <div></div> <div>92%</div> <div>8%</div> </div> </div>
1	B	390	<div> <div>7%</div> <div> <div></div> <div>91%</div> <div>9%</div> </div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6659 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphoglycerate kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	390	Total	C	N	O	S	0	0	0
			2949	1877	525	542	5			
1	B	390	Total	C	N	O	S	0	0	0
			2949	1877	525	542	5			

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Na	0	0
			2	2		
2	A	1	Total	Na	0	0
			1	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 6	C 3	O 3	0	0
3	A	1	Total 6	C 3	O 3	0	0
3	B	1	Total 6	C 3	O 3	0	0
3	B	1	Total 6	C 3	O 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	376	Total 376	O 376	0	0
4	B	358	Total 358	O 358	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphoglycerate kinase



- Molecule 1: Phosphoglycerate kinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	143.18 Å   143.18 Å   75.64 Å 90.00°   90.00°   120.00°	Depositor
Resolution (Å)	30.00 – 1.50 29.43 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.7 (30.00-1.50) 99.9 (29.43-1.50)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.27 (at 1.50 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.201 , 0.215 0.195 , 0.193	Depositor DCC
$R_{free}$ test set	7064 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.3	Xtriage
Anisotropy	0.038	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 44.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.022 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6659	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.26% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.30	0/3006	0.63	1/4071 (0.0%)
1	B	0.29	0/3006	0.59	1/4071 (0.0%)
All	All	0.29	0/6012	0.61	2/8142 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	51	SER	N-CA-C	-5.50	96.16	111.00
1	A	173	ARG	NE-CZ-NH2	-5.07	117.77	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2949	0	3016	22	0
1	B	2949	0	3016	24	0
2	A	1	0	0	0	0
2	B	2	0	0	0	0
3	A	12	0	16	1	0
3	B	12	0	16	1	0
4	A	376	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	358	0	0	7	0
All	All	6659	0	6064	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:9:PRO:HB2	1:B:50:ALA:HB2	1.51	0.91
1:A:172:VAL:HG21	1:A:385:LEU:HD21	1.69	0.72
1:A:180:LYS:HD2	1:A:180:LYS:O	1.92	0.69
1:B:177:ARG:O	1:B:182:PRO:HD3	1.99	0.61
1:B:12:LYS:HE3	4:B:1618:HOH:O	2.04	0.57
1:A:2:ARG:HH12	1:A:390:GLY:C	2.09	0.55
1:A:176:SER:O	1:A:180:LYS:HG3	2.07	0.55
1:B:147:HIS:H	1:B:147:HIS:CD2	2.24	0.54
1:B:360:LYS:HE2	4:B:1620:HOH:O	2.07	0.54
1:B:285:VAL:HG13	1:B:286:PRO:HA	1.90	0.54
1:B:2:ARG:HH12	1:B:390:GLY:N	2.06	0.54
1:A:2:ARG:NH1	1:A:390:GLY:OXT	2.44	0.51
1:B:242:LEU:HD13	1:B:242:LEU:C	2.32	0.50
1:B:193:ALA:HA	1:B:217:ALA:HB3	1.94	0.49
1:B:33:THR:O	1:B:37:GLU:HG3	2.12	0.49
1:A:184:ARG:O	1:A:308:ARG:HD2	2.14	0.47
1:A:183:GLU:HG2	1:A:308:ARG:CB	2.45	0.47
1:A:71:ALA:HB3	1:A:72:PRO:HD3	1.96	0.47
1:A:154:VAL:HG13	1:A:387:VAL:HG11	1.96	0.47
1:B:132:ALA:O	4:B:1427:HOH:O	2.20	0.47
1:B:97:ARG:O	1:B:101:GLU:HG3	2.15	0.46
1:A:164:ALA:HB2	1:A:388:LEU:HD13	1.98	0.45
1:B:71:ALA:HB3	1:B:72:PRO:HD3	1.97	0.45
1:A:183:GLU:HG2	1:A:308:ARG:HB3	1.97	0.45
1:A:242:LEU:C	1:A:242:LEU:HD13	2.37	0.45
1:B:336:ALA:O	4:B:1428:HOH:O	2.21	0.44
1:A:302:ARG:HD3	4:A:1666:HOH:O	2.17	0.44
1:A:172:VAL:CG2	1:A:385:LEU:HD21	2.45	0.44
1:B:232:ARG:HE	1:B:285:VAL:HG13	1.83	0.43
1:A:97:ARG:O	1:A:101:GLU:HG3	2.19	0.43
1:A:7:LEU:O	1:A:9:PRO:HD3	2.18	0.43
1:A:2:ARG:HH12	1:A:390:GLY:N	2.16	0.43

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:126:GLU:HG3	3:B:1403:GOL:H11	2.00	0.43
1:B:147:HIS:H	1:B:147:HIS:HD2	1.64	0.42
1:A:193:ALA:HA	1:A:217:ALA:HB3	2.02	0.41
1:B:354:VAL:HG13	1:B:359:LEU:HB2	2.02	0.41
1:A:270:GLU:HB3	1:A:273:VAL:HB	2.02	0.41
1:A:95:GLU:O	1:A:99:GLU:HG2	2.21	0.41
1:A:126:GLU:HG2	3:A:1401:GOL:H11	2.01	0.41
1:B:82:PRO:HB2	4:B:1695:HOH:O	2.21	0.41
1:B:164:ALA:HB2	1:B:388:LEU:HD13	2.03	0.41
1:B:192:GLY:HA3	4:B:1666:HOH:O	2.21	0.40
1:B:285:VAL:CG1	1:B:286:PRO:HA	2.51	0.40
1:A:97:ARG:HD3	1:A:133:ARG:CZ	2.51	0.40
1:B:320:GLU:HG3	4:B:1757:HOH:O	2.21	0.40
1:B:169:GLU:O	1:B:173:ARG:HG3	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	388/390 (100%)	381 (98%)	6 (2%)	1 (0%)	41	18
1	B	388/390 (100%)	375 (97%)	13 (3%)	0	100	100
All	All	776/780 (100%)	756 (97%)	19 (2%)	1 (0%)	51	25

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	369	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/300 (100%)	297 (99%)	3 (1%)	76	57
1	B	300/300 (100%)	297 (99%)	3 (1%)	76	57
All	All	600/600 (100%)	594 (99%)	6 (1%)	76	57

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	180	LYS
1	A	186	TYR
1	A	350	SER
1	B	147	HIS
1	B	186	TYR
1	B	350	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	25	GLN
1	B	147	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	B	1404	-	5,5,5	0.29	0	5,5,5	0.23	0
3	GOL	A	1402	-	5,5,5	1.11	0	5,5,5	1.68	2 (40%)
3	GOL	B	1403	-	5,5,5	0.31	0	5,5,5	0.31	0
3	GOL	A	1401	-	5,5,5	0.29	0	5,5,5	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	1404	-	-	0/4/4/4	-
3	GOL	A	1402	-	-	0/4/4/4	-
3	GOL	B	1403	-	-	0/4/4/4	-
3	GOL	A	1401	-	-	0/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1402	GOL	C3-C2-C1	2.84	122.73	111.70
3	A	1402	GOL	O2-C2-C3	-2.01	100.28	109.12

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1403	GOL	1	0
3	A	1401	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	390/390 (100%)	0.61	23 (5%)	22 24	8, 16, 28, 37	0
1	B	390/390 (100%)	0.60	28 (7%)	15 16	9, 16, 29, 39	0
All	All	780/780 (100%)	0.60	51 (6%)	18 20	8, 16, 29, 39	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	390	GLY	11.2
1	A	369	GLY	10.1
1	B	390	GLY	8.6
1	B	371	GLY	7.4
1	B	181	ASP	4.9
1	A	368	THR	4.7
1	B	372	ALA	4.6
1	B	370	GLY	4.6
1	A	104	ARG	3.9
1	B	271	ALA	3.7
1	B	26	ASP	3.6
1	B	369	GLY	3.5
1	B	25	GLN	3.4
1	A	268	ARG	3.4
1	A	25	GLN	3.3
1	B	340	GLY	3.3
1	B	238	ASP	3.2
1	A	26	ASP	3.1
1	A	361	GLU	3.1
1	B	5	LEU	3.0
1	B	320	GLU	2.9
1	A	389	GLU	2.8
1	B	104	ARG	2.7
1	A	287	TYR	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	180	LYS	2.6
1	A	370	GLY	2.6
1	B	4	LEU	2.6
1	B	270	GLU	2.6
1	B	292	ILE	2.5
1	B	60	ARG	2.5
1	B	232	ARG	2.5
1	A	276	ARG	2.4
1	A	284	PRO	2.4
1	B	212	LEU	2.4
1	B	285	VAL	2.3
1	A	214	ILE	2.3
1	A	270	GLU	2.3
1	B	214	ILE	2.3
1	A	238	ASP	2.3
1	A	85	ARG	2.2
1	B	268	ARG	2.2
1	B	247	LEU	2.2
1	A	286	PRO	2.2
1	B	368	THR	2.1
1	B	242	LEU	2.1
1	A	215	GLY	2.1
1	B	213	LEU	2.1
1	A	285	VAL	2.0
1	A	28	LYS	2.0
1	A	181	ASP	2.0
1	A	190	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	A	1402	6/6	0.82	0.22	25,26,28,30	0
3	GOL	B	1404	6/6	0.89	0.19	27,28,29,30	0
2	NA	B	1407	1/1	0.93	0.21	43,43,43,43	0
2	NA	B	1406	1/1	0.94	0.30	27,27,27,27	0
3	GOL	B	1403	6/6	0.95	0.11	13,15,17,17	0
2	NA	A	1405	1/1	0.95	0.21	24,24,24,24	0
3	GOL	A	1401	6/6	0.96	0.09	14,15,15,15	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.