



wwPDB X-ray Structure Validation Summary Report ⓘ

May 21, 2020 – 03:46 am BST

PDB ID : 2V6E
Title : protelomerase TelK complexed with substrate DNA
Authors : Aihara, H.; Huang, W.M.; Ellenberger, T.
Deposited on : 2007-07-17
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	558	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 41%, yellow 47%, orange 7%, grey 5%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 41% 47% 7% 5% </div> </div>
1	B	558	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, green 40%, yellow 48%, orange 7%, grey 5%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 2% 40% 48% 7% 5% </div> </div>
2	C	25	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 28%, yellow 64%, orange 8%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 28% 64% 8% </div> </div>
2	E	25	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 28%, yellow 64%, orange 8%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 28% 64% 8% </div> </div>
3	D	19	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 11%, yellow 89%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 11% 89% </div> </div>
3	F	19	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 11%, yellow 89%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 11% 89% </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10082 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTELEMORASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	532	Total	C	N	O	S	0	0	1
			4142	2629	732	767	14			
1	B	532	Total	C	N	O	S	0	0	1
			4142	2629	732	767	14			


- Molecule 2 is a DNA chain called TELRL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	25	Total	C	N	O	P	0	0	0
			516	246	96	150	24			
2	E	25	Total	C	N	O	P	0	0	0
			516	246	96	150	24			

- Molecule 3 is a DNA chain called TELRL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	19	Total	C	N	O	P	0	0	0
			380	184	68	110	18			
3	F	19	Total	C	N	O	P	0	0	0
			380	184	68	110	18			

- Molecule 4 is VANADATE ION (three-letter code: VO4) (formula: O₄V).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	V	0	0
			3	2	1		
4	B	1	Total	O	V	0	0
			3	2	1		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PROTELEMORASE








- Molecule 2: TELRL




- Molecule 2: TELRL



- Molecule 3: TELRL



- Molecule 3: TELRL



4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	157.97Å 157.97Å 90.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.20 47.58 – 3.00	Depositor EDS
% Data completeness (in resolution range)	97.4 (50.00-3.20) 96.1 (47.58-3.00)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 3.01Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.255 , 0.286 0.253 , 0.281	Depositor DCC
R_{free} test set	1680 reflections (3.88%)	wwPDB-VP
Wilson B-factor (Å ²)	55.4	Xtriage
Anisotropy	0.297	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 41.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.409 for h,-k,-l	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	10082	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: VO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/4227	0.69	2/5727 (0.0%)
1	B	0.45	0/4227	0.69	1/5727 (0.0%)
2	C	0.47	0/579	0.87	1/894 (0.1%)
2	E	0.49	0/579	0.86	1/894 (0.1%)
3	D	0.50	0/425	0.82	0/652
3	F	0.51	0/425	0.83	0/652
All	All	0.46	0/10462	0.73	5/14546 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1
2	E	0	1
All	All	0	2

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	24	DT	N1-C1'-C2'	6.22	124.41	112.60
2	E	24	DT	N1-C1'-C2'	6.14	124.26	112.60
1	A	304	GLU	N-CA-C	-5.26	96.78	111.00
1	B	304	GLU	N-CA-C	-5.24	96.86	111.00
1	A	281	LEU	N-CA-C	5.02	124.55	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	9	DT	Sidechain
2	E	9	DT	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4142	0	3975	339	0
1	B	4142	0	3975	341	0
2	C	516	0	283	34	0
2	E	516	0	283	35	0
3	D	380	0	215	27	0
3	F	380	0	215	29	0
4	A	3	0	0	1	0
4	B	3	0	0	0	0
All	All	10082	0	8946	744	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 39.

The worst 5 of 744 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:34:DT:H2''	3:D:35:DG:H5'	1.14	1.13
1:B:378:VAL:HG11	2:E:11:DA:H3'	1.31	1.10
3:F:34:DT:H2''	3:F:35:DG:H5'	1.14	1.08
1:A:165:ASP:HB3	1:A:174:LYS:HD3	1.38	1.05
1:A:378:VAL:HG11	2:C:11:DA:H3'	1.38	1.04

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	530/558 (95%)	404 (76%)	93 (18%)	33 (6%)	1	11
1	B	530/558 (95%)	405 (76%)	92 (17%)	33 (6%)	1	11
All	All	1060/1116 (95%)	809 (76%)	185 (18%)	66 (6%)	1	11

5 of 66 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	11	ASN
1	A	32	LYS
1	A	107	LEU
1	A	157	PRO
1	A	303	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	416/486 (86%)	378 (91%)	38 (9%)	9	34
1	B	416/486 (86%)	379 (91%)	37 (9%)	9	35
All	All	832/972 (86%)	757 (91%)	75 (9%)	9	34

5 of 75 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	406	GLU
1	B	121	LEU
1	B	404	VAL
1	A	457	MET
1	B	47	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 36 such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	521	GLN
1	B	144	ASN
1	B	514	GLN
1	B	67	ASN
1	B	160	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	VO4	B	1536	1,3,2	0,2,4	0.00	-	-		
4	VO4	A	1536	1,3,2	0,2,4	0.00	-	-		

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1536	VO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	532/558 (95%)	-0.09	7 (1%) 77 65	24, 55, 94, 108	0
1	B	532/558 (95%)	-0.05	13 (2%) 59 44	25, 55, 94, 108	0
2	C	25/25 (100%)	0.03	0 100 100	29, 56, 76, 86	0
2	E	25/25 (100%)	0.04	0 100 100	29, 55, 76, 86	0
3	D	19/19 (100%)	-0.04	0 100 100	29, 41, 80, 85	0
3	F	19/19 (100%)	-0.12	0 100 100	28, 40, 80, 86	0
All	All	1152/1204 (95%)	-0.06	20 (1%) 70 57	24, 55, 93, 108	0

The worst 5 of 20 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	144	ASN	3.5
1	A	139	LYS	3.2
1	B	139	LYS	3.1
1	B	145	SER	2.8
1	B	56	GLY	2.7

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	VO4	B	1536	3/5	0.98	0.17	36,36,37,38	0
4	VO4	A	1536	3/5	0.99	0.19	37,37,38,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.