



# Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 06:53 pm BST

PDB ID : 6V7J  
Title : The C2221 crystal form of canavalin at 173 K  
Authors : McPherson, A.  
Deposited on : 2019-12-08  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	445	<div> <div>2%</div> <div> <div></div> <div>68%</div> <div>11%</div> <div>21%</div> </div> </div>
1	B	445	<div> <div></div> <div> <div>69%</div> <div>10%</div> <div>21%</div> </div> </div>
1	C	445	<div> <div>2%</div> <div> <div></div> <div>69%</div> <div>11%</div> <div>21%</div> </div> </div>

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 9121 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Canavalin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	351	Total	C	N	O	S	0	4	0
			2845	1797	490	551	7			
1	B	351	Total	C	N	O	S	0	8	0
			2872	1816	491	557	8			
1	C	353	Total	C	N	O	S	0	7	0
			2878	1820	494	557	7			

- Molecule 2 is BENZOIC ACID (three-letter code: BEZ) (formula:  $C_7H_6O_2$ ) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			9	7	2		
2	B	1	Total	C	O	0	0
			9	7	2		
2	C	1	Total	C	O	0	0
			9	7	2		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Ca	0	0
			2	2		
4	C	1	Total	Ca	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	165	Total	O	0	0
			165	165		
5	B	161	Total	O	0	1
			162	162		
5	C	151	Total	O	0	0
			151	151		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Canavalin




#### • Molecule 1: Canavalin



#### • Molecule 1: Canavalin





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	136.35Å 149.25Å 128.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.01 – 2.00 79.14 – 2.00	Depositor EDS
% Data completeness (in resolution range)	81.2 (50.01-2.00) 81.3 (79.14-2.00)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	0.16	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.14 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0253	Depositor
R, $R_{free}$	0.196 , 0.247 0.203 , 0.248	Depositor DCC
$R_{free}$ test set	3636 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	38.3	Xtriage
Anisotropy	0.546	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 42.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9121	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BEZ, GOL, CA, OCS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.68	1/2899 (0.0%)	0.81	2/3922 (0.1%)
1	B	0.65	0/2935	0.75	0/3970
1	C	0.64	0/2941	0.74	0/3977
All	All	0.66	1/8775 (0.0%)	0.77	2/11869 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	245	GLN	C-N	12.00	1.61	1.34

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	245	GLN	O-C-N	12.56	142.79	122.70
1	A	245	GLN	CA-C-N	-9.81	95.61	117.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	245	GLN	Mainchain



## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2845	0	2818	36	0
1	B	2872	0	2852	32	0
1	C	2878	0	2863	40	0
2	A	9	0	5	0	0
2	B	9	0	5	0	0
2	C	9	0	5	0	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
3	C	6	0	8	0	0
4	A	2	0	0	0	0
4	C	1	0	0	0	0
5	A	165	0	0	7	0
5	B	162	0	0	5	0
5	C	151	0	0	2	0
All	All	9121	0	8572	100	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (100) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:321:GLU:OE2	1:C:331:MET:CE	2.05	1.04
1:C:224:LYS:HA	5:C:701:HOH:O	1.59	1.00
1:C:321:GLU:OE2	1:C:331:MET:HE3	1.69	0.92
1:A:84:ARG:NH2	1:B:81[B]:GLU:OE2	2.09	0.85
1:C:321:GLU:OE2	1:C:331:MET:HE1	1.76	0.84
1:A:245:GLN:CB	5:A:604:HOH:O	2.29	0.80
1:B:84:ARG:NH2	1:C:81:GLU:OE2	2.16	0.79
1:C:407:GLU:O	1:C:411:GLU:HG2	1.91	0.71
1:C:335:ARG:HD3	1:C:337:ALA:HB2	1.72	0.71
1:B:407:GLU:O	1:B:411[A]:GLU:HG2	1.90	0.70
1:A:407:GLU:O	1:A:411:GLU:HG2	1.91	0.69
1:A:245:GLN:HB3	5:A:604:HOH:O	1.92	0.69
1:C:275:GLN:O	1:C:278[B]:ASP:OD1	2.17	0.63

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:309:ASN:O	1:C:342:GLU:HG2	1.99	0.62
1:B:173:GLU:OE2	1:C:321:GLU:HG3	2.00	0.62
1:A:321:GLU:HA	5:A:721:HOH:O	2.00	0.60
1:A:245:GLN:C	5:A:604:HOH:O	2.39	0.60
1:C:392[B]:ARG:HH11	1:C:392[B]:ARG:HB3	1.66	0.60
1:C:396:ASP:OD1	1:C:402:SER:HA	2.01	0.59
1:B:114:GLU:OE1	1:B:158:ARG:HD3	2.03	0.59
1:A:322:GLN:NE2	1:A:330:SER:O	2.35	0.59
1:A:335:ARG:HD3	1:C:189:ALA:O	2.02	0.58
1:A:49:TYR:OH	1:A:321:GLU:OE2	2.21	0.58
1:B:134:GLN:HG2	5:B:725:HOH:O	2.03	0.58
1:A:247:LYS:HB2	1:A:248:PRO:HD2	1.86	0.57
1:A:114:GLU:OE1	1:A:158:ARG:HD3	2.04	0.57
1:C:224:LYS:HD2	1:C:227:ILE:HD11	1.86	0.57
1:C:362:LEU:HD11	1:C:364:MET:SD	2.45	0.56
1:C:114:GLU:OE1	1:C:158:ARG:HD3	2.06	0.56
1:B:55:LYS:HE3	5:B:607:HOH:O	2.04	0.56
1:B:404:GLU:N	1:B:404:GLU:OE1	2.39	0.55
1:B:89:LEU:HB2	1:B:161:LYS:HB2	1.88	0.55
1:A:189:ALA:O	1:B:335:ARG:HD2	2.07	0.54
1:A:89:LEU:HB2	1:A:161:LYS:HB2	1.89	0.54
1:A:207:GLU:HA	1:A:211:LEU:HB2	1.91	0.53
1:A:415:GLU:HB3	1:A:419:VAL:HG12	1.90	0.53
1:A:245:GLN:HB2	5:A:604:HOH:O	2.01	0.53
1:C:362:LEU:HD11	1:C:364:MET:CG	2.39	0.52
1:A:72:ARG:NH1	1:A:170:GLY:O	2.43	0.51
1:B:51:PHE:HA	1:B:55:LYS:HE2	1.92	0.51
1:B:179:SER:HB3	1:B:213:GLU:HA	1.93	0.51
1:C:89:LEU:HB2	1:C:161:LYS:HB2	1.93	0.51
1:B:141:GLN:NE2	5:B:606:HOH:O	2.43	0.50
1:B:252:ARG:CZ	1:B:287:GLN:OE1	2.60	0.50
1:A:191:SER:HB3	1:B:335:ARG:NH2	2.26	0.50
1:B:362:LEU:HD13	1:B:364[A]:MET:HG3	1.94	0.49
1:B:362:LEU:HD12	1:B:362:LEU:C	2.33	0.49
1:C:362:LEU:HD11	1:C:364:MET:HG3	1.95	0.49
1:B:105:ASP:HA	1:C:351:SER:OG	2.13	0.48
1:B:393:GLN:OE1	5:B:601:HOH:O	2.20	0.48
1:A:116:GLN:HB2	1:A:131:LYS:HE2	1.94	0.47
1:C:282:LEU:C	1:C:282:LEU:HD12	2.34	0.47
1:A:282:LEU:HD12	1:A:282:LEU:C	2.35	0.47
1:B:282:LEU:C	1:B:282:LEU:HD12	2.35	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:216:GLU:HA	1:C:216:GLU:OE2	2.15	0.47
1:B:110:VAL:O	1:B:137:ALA:HA	2.15	0.47
1:A:152:ASP:OD2	1:A:155:GLN:N	2.32	0.47
1:A:387:ILE:HG22	1:A:410:LEU:HD11	1.96	0.46
1:A:362:LEU:HD12	1:A:362:LEU:C	2.36	0.46
1:B:127:ARG:CZ	1:C:391:PRO:HG3	2.44	0.46
1:A:110:VAL:O	1:A:137:ALA:HA	2.15	0.46
1:C:72:ARG:NH1	1:C:170:GLY:O	2.48	0.46
1:B:116:GLN:HB2	1:B:131:LYS:HE2	1.98	0.46
1:A:247:LYS:HB2	1:A:248:PRO:CD	2.46	0.45
1:C:387:ILE:HG22	1:C:410:LEU:HD11	1.98	0.45
1:B:224:LYS:HD2	5:C:621:HOH:O	2.16	0.45
1:A:168:ARG:NH1	5:A:613:HOH:O	2.49	0.45
1:C:110:VAL:O	1:C:137:ALA:HA	2.17	0.45
1:C:207:GLU:HA	1:C:211:LEU:HB2	1.99	0.45
1:C:392[B]:ARG:HH22	1:C:404:GLU:HG3	1.82	0.45
1:A:309:ASN:O	1:A:342:GLU:HG3	2.17	0.44
1:B:362:LEU:O	1:B:362:LEU:HD12	2.16	0.44
1:A:362:LEU:O	1:A:362:LEU:HD12	2.18	0.44
1:A:268:ILE:HG22	1:A:276:LEU:HD12	2.00	0.44
1:B:415:GLU:HB2	1:B:419:VAL:HG12	2.00	0.44
1:B:109:LEU:HD12	1:B:109:LEU:HA	1.83	0.44
1:C:320:LEU:O	1:C:321:GLU:HG2	2.18	0.44
1:A:111:LEU:CD2	1:A:365:VAL:HG11	2.48	0.44
1:B:148:LEU:HD12	1:B:148:LEU:HA	1.85	0.43
1:C:377:ASN:OD1	1:C:386:VAL:HG11	2.17	0.43
1:A:320:LEU:HA	1:A:320:LEU:HD23	1.90	0.43
1:C:94:LYS:HB3	1:C:95:PRO:HD2	2.01	0.43
1:B:268:ILE:HG22	1:B:276:LEU:HD12	2.00	0.43
1:C:252:ARG:CZ	1:C:287:GLN:OE1	2.67	0.43
1:B:299:ASN:HB2	5:B:658:HOH:O	2.19	0.42
1:C:392[B]:ARG:NE	1:C:407:GLU:OE1	2.52	0.42
1:C:148:LEU:HD12	1:C:148:LEU:HA	1.85	0.42
1:C:320:LEU:O	1:C:321:GLU:CG	2.68	0.42
1:B:68:ARG:NH1	1:B:90:GLU:OE1	2.51	0.42
1:B:358:ALA:HA	1:B:362:LEU:HD23	2.00	0.42
1:C:392[B]:ARG:HH22	1:C:404:GLU:HA	1.85	0.42
1:A:247:LYS:HE3	5:A:750:HOH:O	2.19	0.41
1:A:254:ARG:NH2	1:A:272:LYS:HB3	2.35	0.41
1:A:164:ILE:HD12	1:A:164:ILE:N	2.35	0.41
1:A:49:TYR:HH	1:A:321:GLU:CD	2.24	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:111:LEU:CD2	1:C:365:VAL:HG11	2.50	0.41
1:C:392[B]:ARG:CZ	1:C:407:GLU:OE1	2.68	0.41
1:A:286:LEU:O	1:A:363:ASN:HA	2.22	0.40
1:C:393[B]:GLN:H	1:C:393[B]:GLN:CD	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	348/445 (78%)	333 (96%)	14 (4%)	1 (0%)	41	37
1	B	352/445 (79%)	339 (96%)	12 (3%)	1 (0%)	41	37
1	C	353/445 (79%)	338 (96%)	14 (4%)	1 (0%)	41	37
All	All	1053/1335 (79%)	1010 (96%)	40 (4%)	3 (0%)	41	37

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	246	ASP
1	B	223	PRO
1	C	223	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/391 (82%)	317 (99%)	2 (1%)	86	90
1	B	323/391 (83%)	321 (99%)	2 (1%)	86	90
1	C	324/391 (83%)	324 (100%)	0	100	100
All	All	966/1173 (82%)	962 (100%)	4 (0%)	91	93

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	148	LEU
1	A	294	PHE
1	B	294	PHE
1	B	341	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	OCS	B	285	1	7,8,9	1.06	1 (14%)	6,11,13	1.55	1 (16%)
1	OCS	A	285	1	7,8,9	1.59	1 (14%)	6,11,13	2.28	2 (33%)
1	OCS	C	285	1	7,8,9	1.73	1 (14%)	6,11,13	2.59	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	OCS	B	285	1	-	1/4/7/9	-
1	OCS	A	285	1	-	1/4/7/9	-
1	OCS	C	285	1	-	1/4/7/9	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	285	OCS	OD1-SG	4.03	1.56	1.45
1	A	285	OCS	OD1-SG	3.78	1.56	1.45
1	B	285	OCS	OD2-SG	2.27	1.55	1.47

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	285	OCS	OD1-SG-CB	-5.06	100.92	106.94
1	A	285	OCS	OD2-SG-OD3	3.67	120.24	111.27
1	A	285	OCS	OD1-SG-CB	-3.65	102.60	106.94
1	B	285	OCS	OD2-SG-CB	-3.15	100.72	105.74
1	C	285	OCS	OD2-SG-OD3	3.11	118.87	111.27

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	285	OCS	N-CA-CB-SG
1	A	285	OCS	N-CA-CB-SG
1	C	285	OCS	N-CA-CB-SG

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 3 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	BEZ	B	501	-	7,9,9	0.97	1 (14%)	8,11,11	0.35	0
3	GOL	C	502	-	5,5,5	0.09	0	5,5,5	0.26	0
3	GOL	A	502	-	5,5,5	0.11	0	5,5,5	0.32	0
2	BEZ	C	501	-	7,9,9	1.39	1 (14%)	8,11,11	0.30	0
2	BEZ	A	501	-	7,9,9	0.97	1 (14%)	8,11,11	0.40	0
3	GOL	B	502	-	5,5,5	0.11	0	5,5,5	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BEZ	B	501	-	-	0/0/4/4	0/1/1/1
3	GOL	C	502	-	-	0/4/4/4	-
3	GOL	A	502	-	-	4/4/4/4	-
2	BEZ	C	501	-	-	0/0/4/4	0/1/1/1
2	BEZ	A	501	-	-	0/0/4/4	0/1/1/1
3	GOL	B	502	-	-	4/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	501	BEZ	C1-C	3.67	1.51	1.47
2	B	501	BEZ	C1-C	2.56	1.49	1.47
2	A	501	BEZ	C1-C	2.55	1.49	1.47

There are no bond angle outliers.

There are no chirality outliers.


All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	GOL	O1-C1-C2-C3
3	A	502	GOL	C1-C2-C3-O3
3	B	502	GOL	O1-C1-C2-O2
3	B	502	GOL	O1-C1-C2-C3
3	B	502	GOL	C1-C2-C3-O3
3	A	502	GOL	O2-C2-C3-O3
3	B	502	GOL	O2-C2-C3-O3
3	A	502	GOL	O1-C1-C2-O2


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	245:GLN	C	246:ASP	N	1.61

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	350/445 (78%)	0.04	7 (2%) 65 63	33, 47, 78, 109	0
1	B	350/445 (78%)	-0.02	1 (0%) 94 93	34, 49, 73, 110	0
1	C	352/445 (79%)	0.11	7 (1%) 65 63	34, 47, 79, 135	0
All	All	1052/1335 (78%)	0.04	15 (1%) 75 74	33, 47, 78, 135	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	226	GLN	8.6
1	C	227	ILE	6.1
1	A	245	GLN	4.7
1	C	223	PRO	4.4
1	C	225	ASP	4.4
1	B	215	GLN	3.7
1	A	46	ASN	3.6
1	A	320	LEU	3.5
1	A	321	GLU	3.3
1	A	330	SER	2.9
1	A	57	LEU	2.6
1	C	321	GLU	2.5
1	C	331	MET	2.4
1	C	330	SER	2.2
1	A	334	ARG	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	OCS	B	285	9/10	0.94	0.13	39,42,61,68	0
1	OCS	A	285	9/10	0.94	0.16	36,40,65,65	0
1	OCS	C	285	9/10	0.94	0.15	38,42,67,74	0

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

### 6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	A	502	6/6	0.69	0.17	66,71,72,76	0
3	GOL	B	502	6/6	0.80	0.23	67,71,75,78	0
4	CA	A	504	1/1	0.82	1.01	131,131,131,131	0
4	CA	C	503	1/1	0.87	0.26	112,112,112,112	0
3	GOL	C	502	6/6	0.88	0.16	77,79,81,82	0
2	BEZ	A	501	9/9	0.93	0.12	41,42,48,49	0
4	CA	A	503	1/1	0.93	0.05	74,74,74,74	0
2	BEZ	B	501	9/9	0.95	0.23	46,56,62,62	0
2	BEZ	C	501	9/9	0.95	0.10	40,45,56,56	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around BEZ A 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around BEZ B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around BEZ C 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.