



Full wwPDB X-ray Structure Validation Report ⓘ

May 27, 2020 – 10:18 pm BST

PDB ID : 2V8W
Title : Crystallographic and mass spectrometric characterisation of eIF4E with N7-cap derivatives
Authors : Brown, C.J.; Mcnae, I.; Fischer, P.M.; Walkinshaw, M.D.
Deposited on : 2007-08-16
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	217	<div> <div>66%</div> <div>14%</div> <div>•</div> <div>17%</div> </div>
1	E	217	<div> <div>5%</div> <div>57%</div> <div>23%</div> <div>•</div> <div>18%</div> </div>
2	B	14	<div> <div>64%</div> <div>29%</div> <div>7%</div> </div>
2	F	14	<div> <div>79%</div> <div>21%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3614 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called EUKARYOTIC TRANSLATION INITIATION FACTOR 4E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	181	Total	C	N	O	S	0	0	0
			1504	960	265	273	6			
1	E	179	Total	C	N	O	S	0	0	0
			1484	948	261	269	6			

- Molecule 2 is a protein called EUKARYOTIC TRANSLATION INITIATION FACTOR 4E-BINDING PROTEIN 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	14	Total	C	N	O	S	0	0	0
			129	81	25	21	2			
2	F	14	Total	C	N	O	S	0	0	0
			129	81	25	21	2			

- Molecule 3 is [[(2R,3S,4R,5R)-5-(6-AMINO-3-METHYL-4-OXO-5H-IMIDAZO[4,5-C]PYRIDIN-1-YL)-3,4-DIHYDROXY-OXOLAN-2-YL]METHOXY-HYDROXY-PHOSPHORYL] PHOSPHONO HYDROGEN PHOSPHATE (three-letter code: MGO) (formula: C₁₂H₂₀N₄O₁₄P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			33	12	4	14	3		
3	E	1	Total	C	N	O	P	0	0
			33	12	4	14	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	136	Total	O	0	0
			136	136		
4	B	13	Total	O	0	0
			13	13		
4	E	138	Total	O	0	0
			138	138		
4	F	15	Total	O	0	0
			15	15		



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	37.99Å 99.61Å 135.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.34 – 2.30 34.34 – 2.30	Depositor EDS
% Data completeness (in resolution range)	95.5 (34.34-2.30) 95.5 (34.34-2.30)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.62 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.184 , 0.243 0.180 , 0.236	Depositor DCC
R_{free} test set	1156 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	20.4	Xtriage
Anisotropy	0.233	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 52.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3614	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MGO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	1/1542 (0.1%)	0.84	4/2084 (0.2%)
1	E	0.84	0/1521	0.84	1/2055 (0.0%)
2	B	0.82	0/130	0.95	0/169
2	F	0.87	0/130	0.70	0/169
All	All	0.83	1/3323 (0.0%)	0.84	5/4477 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	136	CYS	CB-SG	-5.41	1.73	1.81

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	186	ARG	NE-CZ-NH2	-6.04	117.28	120.30
1	A	186	ARG	NE-CZ-NH1	5.88	123.24	120.30
1	A	112	ARG	NE-CZ-NH1	5.68	123.14	120.30
1	E	60	LEU	CA-CB-CG	5.46	127.87	115.30
1	A	114	LEU	CA-CB-CG	5.05	126.92	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1504	0	1468	33	0
1	E	1484	0	1451	43	0
2	B	129	0	132	3	0
2	F	129	0	132	3	0
3	A	33	0	16	0	0
3	E	33	0	16	0	0
4	A	136	0	0	20	1
4	B	13	0	0	1	0
4	E	138	0	0	18	0
4	F	15	0	0	2	0
All	All	3614	0	3215	81	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:E:2081:HOH:O	2:F:56:ARG:HD2	1.53	1.05
1:E:121:GLN:NE2	4:E:2072:HOH:O	1.98	0.96
1:A:61:ARG:HG3	4:A:2023:HOH:O	1.66	0.95
1:E:37:HIS:HD2	1:E:70:GLU:OE1	1.49	0.94
1:E:209:SER:CA	4:E:2132:HOH:O	2.17	0.92
1:E:61:ARG:HD3	4:E:2011:HOH:O	1.79	0.83
1:E:120:GLN:HE22	1:E:125:ASP:H	1.26	0.80
1:E:37:HIS:CD2	1:E:70:GLU:OE1	2.37	0.77
1:A:177:THR:CB	4:A:2108:HOH:O	2.33	0.76
1:E:169:GLU:OE1	1:E:171:GLU:HG2	1.85	0.76
1:E:184:LYS:HE2	4:E:2125:HOH:O	1.87	0.75
1:E:186:ARG:NH2	4:E:2118:HOH:O	2.19	0.74
1:A:105:GLU:OE1	1:A:108:LYS:NZ	2.20	0.74
1:E:132:GLU:OE2	1:E:186:ARG:NH1	2.23	0.71
1:E:181:ARG:NE	4:E:2116:HOH:O	1.91	0.71
1:A:173:ARG:O	1:A:177:THR:HB	1.91	0.71
1:E:34:TYR:CE1	4:E:2021:HOH:O	2.44	0.70
1:E:181:ARG:NH2	4:E:2116:HOH:O	2.27	0.68
1:A:177:THR:HB	4:A:2108:HOH:O	1.91	0.68
1:A:116:THR:HG23	4:A:2071:HOH:O	1.94	0.67
1:A:125:ASP:OD2	1:A:128:ARG:NH2	2.28	0.67
1:A:182:VAL:HG23	4:A:2109:HOH:O	1.94	0.66
1:E:137:LEU:HD22	1:E:149:VAL:HG11	1.78	0.65
1:A:120:GLN:HG2	4:A:2075:HOH:O	1.96	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:200:HIS:HE1	4:E:2062:HOH:O	1.81	0.63
1:A:61:ARG:NH1	4:A:2025:HOH:O	2.31	0.62
1:A:181:ARG:O	1:A:185:GLU:HG3	1.99	0.61
1:E:159:LYS:NZ	4:E:2101:HOH:O	2.35	0.58
1:A:194:VAL:HG23	4:A:2116:HOH:O	2.04	0.57
1:E:193:ILE:O	1:E:193:ILE:HG13	2.05	0.55
1:E:42:ARG:HG2	1:E:96:ASP:HB2	1.88	0.55
2:F:51:ARG:NH1	4:F:2001:HOH:O	2.38	0.55
1:E:57:GLN:HE21	1:E:101:MET:CE	2.20	0.54
1:E:202:ASP:OD1	1:E:203:THR:N	2.40	0.54
1:E:159:LYS:HE2	4:E:2102:HOH:O	2.06	0.54
2:F:51:ARG:HD3	4:F:2004:HOH:O	2.07	0.54
2:B:55:ASP:HB2	4:B:2006:HOH:O	2.06	0.54
1:A:112:ARG:O	1:A:112:ARG:HG2	2.09	0.53
1:A:192:LYS:HB2	4:A:2117:HOH:O	2.08	0.53
1:A:34:TYR:CE2	1:A:35:ILE:HG13	2.44	0.53
1:E:112:ARG:HG2	1:E:112:ARG:O	2.10	0.52
1:A:61:ARG:HD2	4:A:2018:HOH:O	2.11	0.50
1:A:116:THR:CG2	4:A:2071:HOH:O	2.58	0.49
1:A:164:ALA:HA	4:A:2068:HOH:O	2.11	0.49
1:E:193:ILE:HD13	4:E:2120:HOH:O	2.11	0.49
4:A:2090:HOH:O	2:B:51:ARG:HD3	2.11	0.49
1:E:173:ARG:HG2	1:E:215:PHE:HE2	1.77	0.48
1:E:76:TYR:OH	4:E:2027:HOH:O	2.10	0.48
1:E:48:PHE:CE1	1:E:59:ASN:HB2	2.50	0.47
1:E:113:TRP:O	1:E:164:ALA:HA	2.14	0.47
1:E:34:TYR:CD1	4:E:2021:HOH:O	2.64	0.47
1:E:209:SER:N	4:E:2132:HOH:O	2.42	0.46
1:A:112:ARG:HD2	4:A:2119:HOH:O	2.14	0.46
1:E:178:HIS:HD2	1:E:181:ARG:HH22	1.63	0.46
1:A:118:ASN:H	1:A:121:GLN:HE21	1.64	0.46
1:A:118:ASN:ND2	1:A:121:GLN:HG3	2.32	0.45
1:A:184:LYS:NZ	1:E:174:GLU:OE1	2.41	0.45
1:A:137:LEU:HD22	1:A:149:VAL:HG11	1.98	0.45
1:A:32:GLU:N	4:A:2005:HOH:O	2.50	0.45
1:A:159:LYS:HE3	4:A:2095:HOH:O	2.17	0.44
1:E:203:THR:HA	4:E:2129:HOH:O	2.18	0.44
1:A:61:ARG:NH1	1:A:61:ARG:HB3	2.33	0.44
1:A:78:HIS:HB3	4:A:2036:HOH:O	2.18	0.44
1:E:56:TRP:HZ3	1:E:101:MET:HE2	1.83	0.44
1:E:44:ALA:HB1	1:E:62:LEU:CD1	2.48	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:200:HIS:C	1:E:202:ASP:H	2.22	0.43
1:A:85:LEU:HD11	1:A:91:TYR:CE1	2.54	0.43
1:E:112:ARG:HD2	4:E:2066:HOH:O	2.19	0.43
1:E:190:PRO:HG2	1:E:193:ILE:HB	1.99	0.43
2:B:60:MET:O	2:B:63:ARG:NH1	2.43	0.42
1:A:32:GLU:N	4:A:2003:HOH:O	2.53	0.42
1:E:184:LYS:HE2	1:E:217:VAL:O	2.19	0.42
1:A:118:ASN:HD21	1:A:121:GLN:HG3	1.85	0.41
1:A:113:TRP:O	1:A:164:ALA:HA	2.21	0.41
1:E:176:VAL:HG11	1:E:215:PHE:CD1	2.56	0.41
1:E:106:LYS:HA	1:E:106:LYS:HD2	1.78	0.41
1:E:130:TRP:CZ2	1:E:134:LEU:HD21	2.55	0.41
1:A:32:GLU:HB2	4:A:2006:HOH:O	2.20	0.41
1:E:173:ARG:HG2	1:E:215:PHE:CE2	2.55	0.41
1:A:181:ARG:HB3	4:A:2109:HOH:O	2.21	0.40
1:E:120:GLN:HE22	1:E:125:ASP:N	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:2055:HOH:O	4:A:2075:HOH:O[1_455]	1.90	0.30

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	177/217 (82%)	173 (98%)	4 (2%)	0	100	100
1	E	175/217 (81%)	167 (95%)	6 (3%)	2 (1%)	14	15
2	B	12/14 (86%)	12 (100%)	0	0	100	100
2	F	12/14 (86%)	12 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	376/462 (81%)	364 (97%)	10 (3%)	2 (0%)	29	35

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	210	THR
1	E	194	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	162/195 (83%)	151 (93%)	11 (7%)	16	21
1	E	160/195 (82%)	145 (91%)	15 (9%)	8	10
2	B	14/14 (100%)	12 (86%)	2 (14%)	3	3
2	F	14/14 (100%)	13 (93%)	1 (7%)	14	19
All	All	350/418 (84%)	321 (92%)	29 (8%)	11	14

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	78	HIS
1	A	84	ASN
1	A	99	GLU
1	A	112	ARG
1	A	118	ASN
1	A	148	ASP
1	A	153	VAL
1	A	177	THR
1	A	192	LYS
1	A	194	VAL
1	A	212	LYS
2	B	55	ASP
2	B	57	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	42	ARG
1	E	54	LYS
1	E	62	LEU
1	E	78	HIS
1	E	112	ARG
1	E	114	LEU
1	E	122	ARG
1	E	148	ASP
1	E	156	VAL
1	E	159	LYS
1	E	185	GLU
1	E	186	ARG
1	E	198	GLN
1	E	210	THR
1	E	216	VAL
2	F	57	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	80	GLN
1	A	118	ASN
1	A	121	GLN
1	A	198	GLN
1	E	37	HIS
1	E	57	GLN
1	E	120	GLN
1	E	178	HIS
1	E	198	GLN
1	E	200	HIS
1	E	213	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MGO	E	1218	-	28,35,35	1.85	5 (17%)	34,56,56	1.51	7 (20%)
3	MGO	A	1218	-	28,35,35	1.60	6 (21%)	34,56,56	1.68	10 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MGO	E	1218	-	-	2/18/38/38	0/3/3/3
3	MGO	A	1218	-	-	2/18/38/38	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	1218	MGO	O4'-C1'	5.56	1.48	1.41
3	A	1218	MGO	CAV-NAP	5.02	1.41	1.33
3	E	1218	MGO	CAV-NAP	4.22	1.40	1.33
3	E	1218	MGO	CAU-NAP	3.74	1.41	1.35
3	A	1218	MGO	CAU-NAP	2.98	1.40	1.35
3	E	1218	MGO	PBE-OAD	2.89	1.66	1.54
3	A	1218	MGO	PBE-OAJ	2.86	1.65	1.54
3	E	1218	MGO	PBE-OAJ	2.57	1.64	1.54
3	A	1218	MGO	PBG-OAL	2.07	1.65	1.55
3	A	1218	MGO	CAX-CAW	-2.07	1.39	1.44
3	A	1218	MGO	PBE-OAD	2.02	1.62	1.54

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	1218	MGO	CAU-CAM-CAW	-3.87	119.16	122.31
3	A	1218	MGO	CAU-CAM-CAW	-3.74	119.27	122.31
3	A	1218	MGO	CAX-CAV-NAP	-3.50	118.65	123.43
3	E	1218	MGO	CAX-CAV-NAP	-3.22	119.03	123.43
3	A	1218	MGO	NAB-CAU-NAP	3.14	121.46	116.49
3	A	1218	MGO	CAV-NAP-CAU	3.09	120.94	116.95
3	A	1218	MGO	O4'-C4'-C5'	-2.91	99.81	109.37
3	E	1218	MGO	PBG-OAS-PBE	-2.66	123.71	132.83
3	A	1218	MGO	OAD-PBE-OAS	2.65	113.51	104.64
3	E	1218	MGO	CAV-NAP-CAU	2.59	120.29	116.95
3	E	1218	MGO	NAB-CAU-NAP	2.58	120.56	116.49
3	E	1218	MGO	PBF-OAT-PBG	-2.49	124.27	132.83
3	A	1218	MGO	PBG-OAS-PBE	-2.46	124.37	132.83
3	A	1218	MGO	O4'-C4'-C3'	-2.44	100.28	105.11
3	A	1218	MGO	PBF-OAT-PBG	-2.34	124.80	132.83
3	A	1218	MGO	CAX-CAW-NBC	2.07	108.29	105.72
3	E	1218	MGO	CAX-CAW-NBC	2.01	108.22	105.72

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	1218	MGO	PBG-OAS-PBE-OAJ
3	A	1218	MGO	PBG-OAS-PBE-OAD
3	E	1218	MGO	PBG-OAS-PBE-OAI
3	A	1218	MGO	PBG-OAS-PBE-OAI

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	181/217 (83%)	-0.47	1 (0%) 89 92	4, 15, 29, 38	0
1	E	179/217 (82%)	-0.17	11 (6%) 21 27	3, 15, 45, 58	0
2	B	14/14 (100%)	0.38	0 100 100	16, 27, 44, 45	0
2	F	14/14 (100%)	-0.17	0 100 100	13, 19, 27, 33	0
All	All	388/462 (83%)	-0.29	12 (3%) 49 56	3, 15, 39, 58	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	52	LYS	3.4
1	E	191	PRO	3.2
1	E	209	SER	3.2
1	E	53	SER	3.0
1	E	203	THR	2.8
1	E	55	THR	2.6
1	E	193	ILE	2.5
1	E	202	ASP	2.4
1	E	190	PRO	2.4
1	E	214	ARG	2.2
1	E	171	GLU	2.1
1	A	32	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no carbohydrates in this entry.


6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MGO	E	1218	33/33	0.94	0.13	20,30,47,49	0
3	MGO	A	1218	33/33	0.97	0.09	11,14,36,38	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.