



# Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 02:10 am BST

PDB ID : 5V8U  
Title : Small Molecule Inhibitor ABS-143 Bound to the Botulinum Neurotoxin Serotype A Light Chain  
Authors : Allen, K.N.; Silvaggi, N.R.  
Deposited on : 2017-03-22  
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	444	<div> <div>6%</div> <div> <div></div> <div>79%</div> <div>9%</div> <div>12%</div> </div> </div>
1	B	444	<div> <div>7%</div> <div> <div></div> <div>81%</div> <div>6%</div> <div>13%</div> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12881 atoms, of which 6159 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Botulinum neurotoxin type A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	389	Total	C	H	N	O	S	0	0	0
			6249	2034	3094	521	592	8			
1	B	385	Total	C	H	N	O	S	0	0	0
			6186	2011	3065	517	585	8			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP P10845
A	-18	GLY	-	expression tag	UNP P10845
A	-17	SER	-	expression tag	UNP P10845
A	-16	SER	-	expression tag	UNP P10845
A	-15	HIS	-	expression tag	UNP P10845
A	-14	HIS	-	expression tag	UNP P10845
A	-13	HIS	-	expression tag	UNP P10845
A	-12	HIS	-	expression tag	UNP P10845
A	-11	HIS	-	expression tag	UNP P10845
A	-10	HIS	-	expression tag	UNP P10845
A	-9	SER	-	expression tag	UNP P10845
A	-8	SER	-	expression tag	UNP P10845
A	-7	GLY	-	expression tag	UNP P10845
A	-6	LEU	-	expression tag	UNP P10845
A	-5	VAL	-	expression tag	UNP P10845
A	-4	PRO	-	expression tag	UNP P10845
A	-3	ARG	-	expression tag	UNP P10845
A	-2	GLY	-	expression tag	UNP P10845
A	-1	SER	-	expression tag	UNP P10845
A	0	HIS	-	expression tag	UNP P10845
A	2	GLN	PRO	engineered mutation	UNP P10845
B	-19	MET	-	initiating methionine	UNP P10845
B	-18	GLY	-	expression tag	UNP P10845
B	-17	SER	-	expression tag	UNP P10845
B	-16	SER	-	expression tag	UNP P10845

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-15	HIS	-	expression tag	UNP P10845
B	-14	HIS	-	expression tag	UNP P10845
B	-13	HIS	-	expression tag	UNP P10845
B	-12	HIS	-	expression tag	UNP P10845
B	-11	HIS	-	expression tag	UNP P10845
B	-10	HIS	-	expression tag	UNP P10845
B	-9	SER	-	expression tag	UNP P10845
B	-8	SER	-	expression tag	UNP P10845
B	-7	GLY	-	expression tag	UNP P10845
B	-6	LEU	-	expression tag	UNP P10845
B	-5	VAL	-	expression tag	UNP P10845
B	-4	PRO	-	expression tag	UNP P10845
B	-3	ARG	-	expression tag	UNP P10845
B	-2	GLY	-	expression tag	UNP P10845
B	-1	SER	-	expression tag	UNP P10845
B	0	HIS	-	expression tag	UNP P10845
B	2	GLN	PRO	engineered mutation	UNP P10845

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is N-[3-(4-fluorophenyl)-4-methyl-1H-pyrazol-5-yl]-2-sulfanylacetamide (three-letter code: 90M) (formula: C<sub>12</sub>H<sub>12</sub>FN<sub>3</sub>OS).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	S	0	0
			18	12	1	3	1	1		
3	B	1	Total	C	F	N	O	S	0	0
			18	12	1	3	1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	216	Total	O	0	0
			216	216		
4	B	192	Total	O	0	0
			192	192		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Botulinum neurotoxin type A



#### • Molecule 1: Botulinum neurotoxin type A



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.09Å 67.65Å 98.09Å 90.00° 105.22° 90.00°	Depositor
Resolution (Å)	37.56 – 2.05 37.56 – 2.05	Depositor EDS
% Data completeness (in resolution range)	96.0 (37.56-2.05) 91.6 (37.56-2.05)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.80 (at 2.05Å)	Xtriage
Refinement program	PHENIX (1.11_2567)	Depositor
R, $R_{free}$	0.229 , 0.260 0.239 , 0.234	Depositor DCC
$R_{free}$ test set	2665 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.5	Xtriage
Anisotropy	0.193	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 61.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	12881	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.57% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 90M

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.25	0/3224	0.42	0/4351
1	B	0.26	0/3189	0.43	0/4304
All	All	0.25	0/6413	0.43	0/8655

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3155	3094	3104	36	0
1	B	3121	3065	3070	25	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	18	0	0	1	0
3	B	18	0	0	1	0
4	A	216	0	0	26	0
4	B	192	0	0	16	0
All	All	6722	6159	6174	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.



All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:320:LYS:N	4:A:602:HOH:O	1.84	1.10
1:A:402:ASN:OD1	4:A:601:HOH:O	1.80	0.98
1:B:284:LEU:O	4:B:601:HOH:O	1.81	0.96
1:B:416:LEU:O	4:B:602:HOH:O	1.89	0.90
1:A:413:PHE:O	4:A:603:HOH:O	1.94	0.86
1:B:211:GLY:N	4:B:608:HOH:O	2.10	0.85
1:A:292:ASP:OD2	4:A:604:HOH:O	1.96	0.84
1:B:328:SER:OG	4:B:603:HOH:O	1.97	0.82
1:A:311:GLN:NE2	4:A:608:HOH:O	2.12	0.82
1:A:62:PRO:O	4:A:605:HOH:O	1.99	0.81
1:A:283:ARG:NH2	4:A:609:HOH:O	2.12	0.81
1:A:314:LYS:NZ	4:A:611:HOH:O	2.15	0.80
1:A:335:LYS:NZ	4:A:610:HOH:O	2.14	0.79
1:B:-5:VAL:O	4:B:604:HOH:O	2.02	0.76
1:A:371:LYS:N	4:A:612:HOH:O	2.19	0.75
1:B:80:THR:OG1	4:B:606:HOH:O	2.06	0.73
1:A:23:LYS:NZ	1:A:144:TYR:OH	2.20	0.73
1:A:296:THR:O	4:A:606:HOH:O	2.07	0.73
1:B:324:SER:OG	4:B:605:HOH:O	2.06	0.73
1:A:-9:SER:N	4:A:615:HOH:O	2.25	0.70
1:A:13:PRO:O	1:B:311:GLN:NE2	2.25	0.69
1:B:-9:SER:OG	4:B:607:HOH:O	2.10	0.69
1:A:317:PHE:C	4:A:602:HOH:O	2.32	0.68
1:B:50:THR:OG1	4:B:609:HOH:O	2.11	0.67
1:B:102:ASP:OD1	1:B:105:ARG:NH2	2.28	0.66
1:A:26:ASN:ND2	4:A:619:HOH:O	2.30	0.65
1:A:70:VAL:HG21	4:A:665:HOH:O	1.98	0.62
1:B:288:ASN:N	4:B:601:HOH:O	2.05	0.61
1:B:16:GLY:O	1:B:145:ARG:NH2	2.33	0.60
1:B:14:VAL:O	4:B:610:HOH:O	2.17	0.57
1:B:170:HIS:O	4:B:611:HOH:O	2.17	0.57
1:A:292:ASP:CG	4:A:604:HOH:O	2.42	0.56
1:A:13:PRO:HB2	1:B:311:GLN:HG3	1.87	0.56
1:B:257:GLU:N	4:B:617:HOH:O	2.38	0.55
1:B:410:ASN:OD1	4:B:612:HOH:O	2.18	0.55
1:B:285:TYR:C	4:B:601:HOH:O	2.50	0.50
1:A:259:SER:HB2	4:A:703:HOH:O	2.12	0.50
1:A:102:ASP:OD2	1:A:105:ARG:NH2	2.46	0.48
1:A:283:ARG:CZ	4:A:609:HOH:O	2.60	0.46
1:A:58:ASP:HA	4:A:629:HOH:O	2.15	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:138:ILE:CD1	1:B:138:ILE:N	2.78	0.46
1:A:394:ASN:ND2	4:A:607:HOH:O	2.11	0.46
1:A:264:ARG:NH2	4:A:633:HOH:O	2.49	0.45
1:A:15:ASN:HB3	1:B:311:GLN:OE1	2.17	0.45
1:A:14:VAL:CG2	1:A:20:ALA:HA	2.47	0.45
1:B:26:ASN:HA	4:B:628:HOH:O	2.17	0.44
1:B:138:ILE:N	1:B:138:ILE:HD12	2.33	0.44
1:A:242:VAL:HA	1:A:259:SER:HA	2.00	0.43
1:A:320:LYS:HB3	4:A:602:HOH:O	2.19	0.42
1:A:16:GLY:O	1:A:145:ARG:NH2	2.52	0.42
1:A:394:ASN:HB2	4:A:607:HOH:O	2.20	0.41
1:A:289:LYS:NZ	4:A:638:HOH:O	2.53	0.41
1:A:315:ASN:OD1	4:A:613:HOH:O	2.22	0.41
1:A:163:PHE:O	3:A:502:90M:N5	2.54	0.41
1:B:163:PHE:O	3:B:502:90M:N5	2.53	0.41
1:A:56:GLU:OE1	1:A:187:ARG:NH1	2.54	0.40
1:A:284:LEU:HD11	4:A:627:HOH:O	2.21	0.40
1:B:97:ARG:HA	1:B:386:ILE:HG23	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	375/444 (84%)	362 (96%)	13 (4%)	0	100	100
1	B	371/444 (84%)	358 (96%)	13 (4%)	0	100	100
All	All	746/888 (84%)	720 (96%)	26 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	349/397 (88%)	347 (99%)	2 (1%)	86	87
1	B	345/397 (87%)	343 (99%)	2 (1%)	86	87
All	All	694/794 (87%)	690 (99%)	4 (1%)	86	87

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	PHE
1	A	324	SER
1	B	51	PHE
1	B	138	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	90M	B	502	2	15,19,19	2.01	5 (33%)	14,26,26	1.01	2 (14%)
3	90M	A	502	2	15,19,19	2.00	5 (33%)	14,26,26	1.00	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	90M	B	502	2	-	0/6/10/10	0/2/2/2
3	90M	A	502	2	-	1/6/10/10	0/2/2/2

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	90M	N2-N1	4.09	1.45	1.37
3	B	502	90M	N2-N1	4.09	1.45	1.37
3	A	502	90M	C5-N5	3.88	1.47	1.39
3	B	502	90M	C5-N5	3.79	1.47	1.39
3	B	502	90M	C12-N5	2.58	1.41	1.35
3	B	502	90M	C6-C3	-2.54	1.46	1.49
3	A	502	90M	C12-N5	2.54	1.41	1.35
3	A	502	90M	C6-C3	-2.46	1.46	1.49
3	A	502	90M	C4B-C2	-2.19	1.47	1.51
3	B	502	90M	C4B-C2	-2.19	1.47	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	502	90M	C6-C3-N2	-2.07	117.23	120.78
3	A	502	90M	C13-C12-N5	2.05	117.19	114.38
3	B	502	90M	C13-C12-N5	2.05	117.19	114.38

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	502	90M	O12-C12-N5-C5


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	502	90M	1	0
3	A	502	90M	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	389/444 (87%)	0.43	28 (7%) 15 17	18, 39, 75, 98	0
1	B	385/444 (86%)	0.51	31 (8%) 12 12	16, 36, 83, 105	0
All	All	774/888 (87%)	0.47	59 (7%) 13 14	16, 38, 79, 105	0

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	140	PRO	7.3
1	B	277	LEU	7.3
1	B	70	VAL	6.1
1	B	276	SER	6.0
1	B	269	HIS	5.3
1	A	54	PRO	5.1
1	B	40	ASN	4.4
1	B	271	ALA	4.4
1	A	300	ALA	4.1
1	B	284	LEU	3.7
1	B	273	PHE	3.6
1	A	26	ASN	3.5
1	A	123	ILE	3.5
1	B	123	ILE	3.3
1	B	328	SER	3.2
1	B	62	PRO	3.2
1	A	172	VAL	3.1
1	A	29	GLN	3.0
1	A	328	SER	3.0
1	A	257	GLU	3.0
1	B	144	TYR	3.0
1	B	211	GLY	2.9
1	B	54	PRO	2.9
1	B	263	LEU	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	55	GLU	2.8
1	B	300	ALA	2.8
1	B	336	LEU	2.8
1	B	197	GLU	2.8
1	A	294	ALA	2.8
1	B	301	LYS	2.8
1	A	367	LEU	2.7
1	B	124	ASP	2.7
1	A	299	LYS	2.7
1	A	273	PHE	2.6
1	B	171	GLU	2.5
1	A	301	LYS	2.5
1	A	272	LYS	2.5
1	B	39	HIS	2.5
1	A	271	ALA	2.5
1	A	122	THR	2.4
1	A	211	GLY	2.4
1	B	257	GLU	2.4
1	A	171	GLU	2.3
1	A	62	PRO	2.3
1	A	260	PHE	2.3
1	B	143	SER	2.3
1	A	126	GLU	2.3
1	A	296	THR	2.2
1	B	280	ASN	2.2
1	B	272	LYS	2.2
1	A	310	LEU	2.2
1	A	242	VAL	2.2
1	B	243	PHE	2.1
1	B	29	GLN	2.1
1	A	309	SER	2.1
1	A	151	LEU	2.1
1	A	124	ASP	2.1
1	A	332	SER	2.0
1	B	225	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



## 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	90M	A	502	18/18	0.86	0.34	56,115,151,155	0
3	90M	B	502	18/18	0.95	0.12	31,35,42,42	0
2	ZN	B	501	1/1	0.99	0.11	39,39,39,39	0
2	ZN	A	501	1/1	0.99	0.06	46,46,46,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.