



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 8, 2020 – 02:20 AM EST

PDB ID : 6V8D
Title : Design, Synthesis, and Mechanism of Fluorine-substituted Cyclohexene Analogues of GABA-Aminobutyric Acid (GABA) as Selective Ornithine Aminotransferase Inactivators
Authors : Zhu, W.; Doubleday, P.T.; Catlin, D.S.; Weerawarna, P.; Butrin, A.; Shen, S.; Kelleher, N.L.; Liu, D.; Silverman, R.B.
Deposited on : 2019-12-10
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.15.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.15.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	404	<div> <div>0%</div> <div>91%</div> <div>8%</div> </div>
1	B	404	<div> <div>3%</div> <div>87%</div> <div>11%</div> <div>.</div> </div>
1	C	404	<div> <div>3%</div> <div>88%</div> <div>11%</div> <div>.</div> </div>

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 10072 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ornithine aminotransferase, mitochondrial.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	402	Total	C	N	O	S	0	0	0
			3150	2023	531	584	12			
1	B	402	Total	C	N	O	S	0	0	0
			3150	2023	531	584	12			
1	C	402	Total	C	N	O	S	0	0	0
			3150	2023	531	584	12			

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C₈H₁₀NO₆P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	C	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 3 is (3Z)-3-iminocyclohex-1-ene-1-carboxylic acid (three-letter code: O78) (formula: $C_7H_9NO_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			10	7	1	2		
3	B	1	Total	C	N	O	0	0
			10	7	1	2		
3	C	1	Total	C	N	O	0	0
			10	7	1	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	187	Total	O	0	0
			187	187		
4	B	196	Total	O	0	0
			196	196		
4	C	164	Total	O	0	0
			164	164		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ornithine aminotransferase, mitochondrial



- Molecule 1: Ornithine aminotransferase, mitochondrial



- Molecule 1: Ornithine aminotransferase, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	202.60Å 111.96Å 57.87Å 90.00° 104.11° 90.00°	Depositor
Resolution (Å)	31.36 – 2.25 31.36 – 2.25	Depositor EDS
% Data completeness (in resolution range)	91.7 (31.36-2.25) 91.7 (31.36-2.25)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.21 (at 2.24Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.157 , 0.196 0.157 , 0.196	Depositor DCC
R_{free} test set	2703 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	28.2	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 52.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.025 for -h-2*k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10072	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: O78, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/3223	0.48	0/4375
1	B	0.31	0/3223	0.48	1/4375 (0.0%)
1	C	0.31	0/3223	0.48	0/4375
All	All	0.30	0/9669	0.48	1/13125 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	351	LYS	CD-CE-NZ	-6.97	95.67	111.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	386	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3150	0	3153	27	0
1	B	3150	0	3153	42	0
1	C	3150	0	3154	34	1
2	A	15	0	7	0	0
2	B	15	0	7	0	0
2	C	15	0	7	0	0
3	A	10	0	0	0	0
3	B	10	0	0	0	0
3	C	10	0	0	0	0
4	A	187	0	0	6	0
4	B	196	0	0	5	0
4	C	164	0	0	7	0
All	All	10072	0	9481	90	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:351:LYS:O	1:B:351:LYS:NZ	1.85	1.09
1:C:331:ARG:NH2	4:C:601:HOH:O	1.86	1.09
1:A:176:ASN:OD1	4:A:601:HOH:O	1.90	0.88
1:B:143:VAL:HG12	1:B:179:GLY:HA3	1.56	0.86
1:B:407:THR:HG22	1:B:411:ILE:HB	1.57	0.85
1:B:63:GLU:OE1	4:B:601:HOH:O	2.03	0.76
1:C:392:LYS:NZ	4:C:603:HOH:O	2.13	0.76
1:A:154:ARG:HD3	1:B:198:GLY:HA3	1.73	0.70
1:C:64:ARG:HA	1:C:64:ARG:HE	1.54	0.70
1:C:64:ARG:HE	1:C:65:GLY:H	1.41	0.69
1:A:198:GLY:HA3	1:B:154:ARG:HD3	1.74	0.68
1:C:200:PHE:O	4:C:602:HOH:O	2.10	0.68
1:B:143:VAL:CG1	1:B:179:GLY:HA3	2.24	0.68
1:A:422:GLU:OE1	4:A:603:HOH:O	2.14	0.66
1:B:251:GLU:OE2	4:B:603:HOH:O	2.13	0.64
1:C:265:ILE:HG22	1:C:292:LYS:HD2	1.78	0.64
1:B:351:LYS:NZ	1:B:355:ILE:HG13	2.13	0.64
1:B:351:LYS:HZ1	1:B:355:ILE:HG13	1.62	0.64
1:B:265:ILE:HG22	1:B:292:LYS:HD2	1.81	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:64:ARG:NE	1:C:65:GLY:H	1.99	0.61
1:C:331:ARG:HD2	4:C:735:HOH:O	2.02	0.60
1:C:132:ASN:O	1:C:284:ARG:NH2	2.35	0.59
1:A:359:GLU:O	4:A:604:HOH:O	2.17	0.57
1:A:317:GLY:O	1:B:180:ARG:NH2	2.38	0.56
1:C:250:ARG:HD3	1:C:250:ARG:O	2.07	0.55
1:C:64:ARG:CA	1:C:64:ARG:HE	2.19	0.54
1:C:65:GLY:HA3	1:C:70:LEU:HD23	1.90	0.54
1:B:382:ILE:HG23	1:B:388:TRP:CZ2	2.43	0.53
1:C:408:HIS:CD2	1:C:411:ILE:HD12	2.44	0.53
1:C:409:GLY:N	4:C:608:HOH:O	2.41	0.53
1:A:158:TYR:OH	1:B:198:GLY:HA2	2.09	0.52
1:C:386:LYS:HG3	1:C:386:LYS:O	2.09	0.52
1:A:198:GLY:HA2	1:B:158:TYR:OH	2.09	0.52
1:A:97:ILE:HG22	1:A:298:LEU:HD22	1.91	0.52
1:A:202:PRO:HG2	1:B:199:PRO:HB2	1.91	0.52
1:B:117:ASN:ND2	4:B:608:HOH:O	2.42	0.52
1:A:199:PRO:HB2	1:B:202:PRO:HG2	1.92	0.52
1:C:64:ARG:HE	1:C:65:GLY:N	2.06	0.51
1:A:149:ALA:HB2	1:A:289:LEU:HD21	1.92	0.51
1:B:340:LEU:HD23	1:B:345:LEU:HD12	1.93	0.51
1:C:251:GLU:OE2	4:C:604:HOH:O	2.19	0.51
1:C:64:ARG:NH2	1:C:93:CYS:SG	2.84	0.51
1:B:149:ALA:HB2	1:B:289:LEU:HD21	1.94	0.49
1:C:364:PRO:HG2	1:C:367:VAL:HG12	1.95	0.49
1:C:158:TYR:OH	1:C:166:TYR:HA	2.12	0.49
1:C:278:VAL:HG22	1:C:283:VAL:HB	1.93	0.49
1:C:96:LYS:HD3	1:C:339:VAL:HG12	1.94	0.49
1:A:178:TRP:CZ2	1:A:184:ALA:HA	2.48	0.48
1:C:232:ILE:HD11	1:C:374:LYS:HD3	1.95	0.48
1:A:66:LYS:NZ	4:A:602:HOH:O	1.99	0.48
1:A:299:TYR:CZ	1:B:300:PRO:HD3	2.47	0.48
1:A:158:TYR:OH	1:A:166:TYR:HA	2.13	0.48
1:C:64:ARG:NE	1:C:64:ARG:HA	2.24	0.48
1:A:101:LEU:HD22	1:B:105:VAL:HG21	1.95	0.48
1:C:149:ALA:HB2	1:C:289:LEU:HD21	1.95	0.47
1:C:97:ILE:HG22	1:C:298:LEU:HD22	1.94	0.47
1:A:300:PRO:HD3	1:B:299:TYR:CZ	2.48	0.47
1:A:266:GLN:OE1	1:A:413:ARG:NH1	2.48	0.47
1:B:178:TRP:CZ2	1:B:184:ALA:HA	2.50	0.47
1:A:198:GLY:HA2	1:B:158:TYR:HH	1.79	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:247:MET:HG3	1:B:281:GLU:HB3	1.98	0.46
1:A:386:LYS:HD2	1:A:386:LYS:HA	1.58	0.46
1:B:82:LEU:HD13	1:B:405:LYS:HG2	1.98	0.46
1:B:158:TYR:OH	1:B:166:TYR:HA	2.16	0.45
1:A:62:LEU:HA	1:A:72:ASP:HA	1.99	0.45
1:B:387:ASP:OD1	1:B:387:ASP:N	2.50	0.45
1:C:178:TRP:CZ2	1:C:184:ALA:HA	2.52	0.45
1:C:340:LEU:HD23	1:C:345:LEU:HD12	1.98	0.45
1:A:405:LYS:HD2	1:B:114:ALA:HB1	1.98	0.44
1:A:158:TYR:HH	1:B:198:GLY:HA2	1.82	0.44
1:C:371:VAL:HG22	1:C:380:ILE:HG22	1.98	0.44
1:B:188:SER:HB2	4:B:686:HOH:O	2.18	0.44
1:A:409:GLY:HA2	4:A:630:HOH:O	2.17	0.43
1:B:390:ALA:HB1	1:B:406:PRO:HB3	2.00	0.43
1:B:351:LYS:HZ3	1:B:351:LYS:C	2.18	0.43
1:C:66:LYS:HD3	1:C:93:CYS:HB2	2.00	0.43
1:C:194:TYR:HB2	1:C:200:PHE:CE1	2.54	0.43
1:B:364:PRO:HG2	1:B:367:VAL:HG12	1.99	0.43
1:A:113:ARG:NH1	4:A:611:HOH:O	2.50	0.43
1:B:71:TRP:HA	1:B:76:ARG:O	2.19	0.43
1:B:382:ILE:HD13	1:B:382:ILE:HA	1.88	0.42
1:B:72:ASP:OD2	1:B:76:ARG:NH1	2.49	0.42
1:B:321:SER:OG	4:B:602:HOH:O	2.10	0.42
1:B:351:LYS:NZ	1:B:351:LYS:C	2.68	0.42
1:C:178:TRP:HB3	4:C:607:HOH:O	2.21	0.41
1:B:423:ASP:OD1	1:B:424:GLU:N	2.54	0.41
1:C:248:GLY:HA2	1:C:251:GLU:HG2	2.01	0.41
1:B:128:THR:HB	1:B:133:TYR:O	2.21	0.40
1:C:70:LEU:HD21	1:C:92:HIS:CE1	2.56	0.40
1:A:405:LYS:HG3	1:A:406:PRO:HD2	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:64:ARG:NH1	1:C:106:ASP:O[2_555]	2.14	0.06

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	400/404 (99%)	378 (94%)	21 (5%)	1 (0%)	41	46
1	B	400/404 (99%)	378 (94%)	20 (5%)	2 (0%)	29	29
1	C	400/404 (99%)	380 (95%)	19 (5%)	1 (0%)	41	46
All	All	1200/1212 (99%)	1136 (95%)	60 (5%)	4 (0%)	41	46

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	292	LYS
1	B	292	LYS
1	C	292	LYS
1	B	408	HIS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	336/337 (100%)	332 (99%)	4 (1%)	71	80
1	B	336/337 (100%)	331 (98%)	5 (2%)	65	75
1	C	336/337 (100%)	329 (98%)	7 (2%)	53	62
All	All	1008/1011 (100%)	992 (98%)	16 (2%)	62	73

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	183	SER
1	A	321	SER
1	A	388	TRP
1	A	405	LYS
1	B	260	PHE
1	B	366	ASP
1	B	387	ASP
1	B	388	TRP
1	B	405	LYS
1	C	66	LYS
1	C	101	LEU
1	C	183	SER
1	C	260	PHE
1	C	321	SER
1	C	388	TRP
1	C	426	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	408	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	O78	B	502	-	7,10,10	1.10	1 (14%)	7,13,13	1.74	3 (42%)
3	O78	C	502	2	7,10,10	1.13	1 (14%)	7,13,13	1.70	2 (28%)
3	O78	A	502	2	7,10,10	1.20	1 (14%)	7,13,13	1.55	2 (28%)
2	PLP	A	501	1,3	15,15,16	1.03	2 (13%)	20,22,23	0.93	0
2	PLP	B	501	-	15,15,16	0.99	1 (6%)	20,22,23	0.90	0
2	PLP	C	501	1,3	15,15,16	1.08	2 (13%)	20,22,23	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	O78	B	502	-	-	0/0/14/14	0/1/1/1
3	O78	C	502	2	-	0/0/14/14	0/1/1/1
3	O78	A	502	2	-	0/0/14/14	0/1/1/1
2	PLP	A	501	1,3	-	2/6/6/8	0/1/1/1
2	PLP	B	501	-	-	0/6/6/8	0/1/1/1
2	PLP	C	501	1,3	-	0/6/6/8	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	O78	C7-C6	2.92	1.48	1.42
3	C	502	O78	C7-C6	2.81	1.47	1.42
3	B	502	O78	C7-C6	2.72	1.47	1.42
2	A	501	PLP	C2-N1	2.46	1.38	1.33
2	C	501	PLP	C2-N1	2.44	1.38	1.33
2	B	501	PLP	C2-N1	2.34	1.38	1.33
2	A	501	PLP	C6-N1	2.10	1.38	1.34
2	C	501	PLP	C6-N1	2.08	1.38	1.34

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	502	O78	C4-C5-C6	-3.03	104.96	111.76
3	B	502	O78	C4-C5-C6	-3.00	105.04	111.76
3	C	502	O78	C5-C6-C7	2.46	120.33	117.70
3	A	502	O78	C4-C5-C6	-2.42	106.33	111.76
3	B	502	O78	C5-C6-C7	2.33	120.19	117.70
3	A	502	O78	C5-C6-C7	2.32	120.18	117.70
3	B	502	O78	C2-C7-C6	-2.08	119.01	122.62

There are no chirality outliers.


All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	PLP	C5A-O4P-P-O1P
2	A	501	PLP	C4-C5-C5A-O4P



There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	402/404 (99%)	-0.29	5 (1%) 79 81	18, 25, 40, 70	0
1	B	402/404 (99%)	-0.18	13 (3%) 47 50	18, 26, 41, 78	0
1	C	402/404 (99%)	-0.06	14 (3%) 44 46	20, 30, 49, 77	0
All	All	1206/1212 (99%)	-0.18	32 (2%) 54 57	18, 27, 44, 78	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	387	ASP	6.5
1	C	108	LEU	5.7
1	C	388	TRP	5.4
1	B	386	LYS	4.6
1	C	387	ASP	4.5
1	A	387	ASP	4.5
1	B	366	ASP	4.2
1	C	107	LYS	4.2
1	B	364	PRO	4.1
1	B	408	HIS	3.9
1	A	386	LYS	3.8
1	A	408	HIS	3.7
1	B	385	THR	3.6
1	C	75	GLY	3.5
1	C	386	LYS	3.4
1	A	198	GLY	3.2
1	B	367	VAL	3.2
1	B	351	LYS	3.1
1	B	198	GLY	2.7
1	C	384	GLU	2.7
1	C	408	HIS	2.6
1	B	384	GLU	2.5
1	C	38	PRO	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	198	GLY	2.3
1	C	39	THR	2.2
1	A	388	TRP	2.2
1	C	410	ASP	2.2
1	C	322	THR	2.2
1	B	41	ASP	2.1
1	B	382	ILE	2.1
1	B	388	TRP	2.1
1	C	320	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	O78	A	502	10/10	0.81	0.20	29,35,38,41	0
3	O78	C	502	10/10	0.84	0.22	33,38,46,48	0
3	O78	B	502	10/10	0.84	0.21	32,34,41,41	0
2	PLP	B	501	15/16	0.97	0.15	21,25,31,32	0
2	PLP	A	501	15/16	0.98	0.12	19,25,29,32	0
2	PLP	C	501	15/16	0.98	0.16	23,28,32,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around O78 A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around O78 C 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around O78 B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around PLP B 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PLP A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around PLP C 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.