



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 16, 2020 – 07:33 PM BST

PDB ID : 6V9U  
Title : Crystal structure of human TLR8 ectodomain bound to small molecule antagonist 14c  
Authors : Critton, D.A.  
Deposited on : 2019-12-16  
Resolution : 2.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	811	<div> <div>33%</div> <div>83%</div> <div>8%</div> <div>9%</div> </div>
1	B	811	<div> <div>26%</div> <div>82%</div> <div>8%</div> <div>9%</div> </div>
2	C	5	<div> <div>20%</div> <div>80%</div> </div>
2	E	5	<div> <div>60%</div> <div>40%</div> </div>
2	F	5	<div> <div>40%</div> <div>60%</div> </div>
3	D	2	<div> <div>50%</div> <div>50%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	G	3	 67% 33%
5	H	5	 60% 40%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	NAG	A	918	-	-	-	X
6	NAG	A	921	-	-	-	X
6	NAG	B	901	-	-	-	X

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 12082 atoms, of which 60 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Toll-like receptor 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	738	Total	C	N	O	S	0	2	0
			5731	3689	952	1071	19			
1	B	735	Total	C	N	O	S	0	1	0
			5646	3637	927	1063	19			

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	23	ARG	-	expression tag	UNP Q9NR97
A	24	SER	-	expression tag	UNP Q9NR97
A	25	PRO	-	expression tag	UNP Q9NR97
A	26	TRP	-	expression tag	UNP Q9NR97
A	828	GLU	-	expression tag	UNP Q9NR97
A	829	PHE	-	expression tag	UNP Q9NR97
A	830	LEU	-	expression tag	UNP Q9NR97
A	831	VAL	-	expression tag	UNP Q9NR97
A	832	PRO	-	expression tag	UNP Q9NR97
A	833	ARG	-	expression tag	UNP Q9NR97
B	23	ARG	-	expression tag	UNP Q9NR97
B	24	SER	-	expression tag	UNP Q9NR97
B	25	PRO	-	expression tag	UNP Q9NR97
B	26	TRP	-	expression tag	UNP Q9NR97
B	828	GLU	-	expression tag	UNP Q9NR97
B	829	PHE	-	expression tag	UNP Q9NR97
B	830	LEU	-	expression tag	UNP Q9NR97
B	831	VAL	-	expression tag	UNP Q9NR97
B	832	PRO	-	expression tag	UNP Q9NR97
B	833	ARG	-	expression tag	UNP Q9NR97

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	5	Total	C	N	O	0	0	0
			61	34	2	25			
2	E	5	Total	C	N	O	0	0	0
			61	34	2	25			
2	F	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	G	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	H	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 7 is 2-(3,4-dimethoxyphenyl)-5-(piperidin-4-yl)-3-(propan-2-yl)-1H-indole (three-letter code: QSM) (formula: C<sub>24</sub>H<sub>30</sub>N<sub>2</sub>O<sub>2</sub>) (labeled as "Ligand of Interest" by author).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total	C	H	N	O	30	0
			58	24	30	2	2		
7	B	1	Total	C	H	N	O	30	0
			58	24	30	2	2		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	23	Total 23	O 23	0	0
8	B	17	Total 17	O 17	0	0








- Molecule 2: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 20% 80%



- Molecule 2: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 60% 40%



- Molecule 2: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F: 40% 60%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:



- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:



- Molecule 5: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	163.59 Å   86.75 Å   151.40 Å 90.00°   119.80°   90.00°	Depositor
Resolution (Å)	131.37 – 2.65 81.69 – 2.65	Depositor EDS
% Data completeness (in resolution range)	99.3 (131.37-2.65) 99.3 (81.69-2.65)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.19 (at 2.65 Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, $R_{free}$	0.296 , 0.306 0.324 , 0.339	Depositor DCC
$R_{free}$ test set	2652 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	57.8	Xtriage
Anisotropy	0.517	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 76.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.011 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.86	EDS
Total number of atoms	12082	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	101.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, QSM, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/5861	0.58	0/7992
1	B	0.40	0/5768	0.59	1/7877 (0.0%)
All	All	0.40	0/11629	0.58	1/15869 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	601	ASP	C-N-CA	6.70	138.45	121.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5731	0	5492	27	0
1	B	5646	0	5370	27	0
2	C	61	0	52	0	0
2	E	61	0	52	0	0
2	F	61	0	52	0	0
3	D	28	0	25	0	0
4	G	39	0	34	0	0
5	H	61	0	52	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	126	0	117	0	0
6	B	112	0	104	0	0
7	A	28	30	0	0	0
7	B	28	30	0	0	0
8	A	23	0	0	0	0
8	B	17	0	0	0	0
All	All	12022	60	11350	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 52 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:670:HIS:HA	1:B:694:ASP:HB3	1.89	0.55
1:A:670:HIS:HA	1:A:694:ASP:HB3	1.89	0.54
1:A:373:HIS:HA	1:A:400:ASN:HB3	1.90	0.54
1:B:373:HIS:HA	1:B:400:ASN:HB3	1.89	0.54
1:A:538:THR:HG22	1:A:562:SER:HB2	1.89	0.53

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	728/811 (90%)	667 (92%)	58 (8%)	3 (0%)	34	48
1	B	724/811 (89%)	661 (91%)	60 (8%)	3 (0%)	34	48
All	All	1452/1622 (90%)	1328 (92%)	118 (8%)	6 (0%)	34	48

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	602	LYS
1	B	573	VAL
1	A	575	HIS
1	A	378	VAL
1	B	378	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	625/755 (83%)	610 (98%)	15 (2%)	49	67
1	B	612/755 (81%)	594 (97%)	18 (3%)	42	60
All	All	1237/1510 (82%)	1204 (97%)	33 (3%)	44	63

5 of 33 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	764	LEU
1	B	261	PHE
1	B	689	ARG
1	B	49	CYS
1	B	160	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	97	ASN
1	A	284	GLN
1	B	97	ASN
1	B	262	ASN
1	B	576	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

25 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	NAG	C	1	1,2	14,14,15	0.32	0	17,19,21	0.83	1 (5%)
2	NAG	C	2	2	14,14,15	0.31	0	17,19,21	0.85	1 (5%)
2	BMA	C	3	2	11,11,12	0.35	0	15,15,17	0.66	0
2	MAN	C	4	2	11,11,12	0.39	0	15,15,17	0.82	1 (6%)
2	MAN	C	5	2	11,11,12	0.39	0	15,15,17	0.85	1 (6%)
3	NAG	D	1	1,3	14,14,15	0.30	0	17,19,21	0.60	0
3	NAG	D	2	3	14,14,15	0.30	0	17,19,21	0.62	1 (5%)
2	NAG	E	1	1,2	14,14,15	0.27	0	17,19,21	0.60	0
2	NAG	E	2	2	14,14,15	0.28	0	17,19,21	0.52	0
2	BMA	E	3	2	11,11,12	0.32	0	15,15,17	0.48	0
2	MAN	E	4	2	11,11,12	0.37	0	15,15,17	0.80	1 (6%)
2	MAN	E	5	2	11,11,12	0.41	0	15,15,17	0.97	2 (13%)
2	NAG	F	1	1,2	14,14,15	0.29	0	17,19,21	0.83	1 (5%)
2	NAG	F	2	2	14,14,15	0.27	0	17,19,21	0.64	0
2	BMA	F	3	2	11,11,12	0.31	0	15,15,17	0.65	0
2	MAN	F	4	2	11,11,12	0.37	0	15,15,17	0.95	1 (6%)
2	MAN	F	5	2	11,11,12	0.42	0	15,15,17	1.10	1 (6%)
4	NAG	G	1	1,4	14,14,15	0.30	0	17,19,21	0.60	0
4	NAG	G	2	4	14,14,15	0.32	0	17,19,21	0.66	1 (5%)
4	BMA	G	3	4	11,11,12	0.35	0	15,15,17	0.53	0
5	NAG	H	1	1,5	14,14,15	0.26	0	17,19,21	0.60	0
5	NAG	H	2	5	14,14,15	0.28	0	17,19,21	0.54	0
5	BMA	H	3	5	11,11,12	0.30	0	15,15,17	0.47	0
5	MAN	H	4	5	11,11,12	0.41	0	15,15,17	0.94	2 (13%)



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	MAN	H	5	5	11,11,12	0.35	0	15,15,17	0.76	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	0/2/19/22	0/1/1/1
2	MAN	C	5	2	-	1/2/19/22	0/1/1/1
3	NAG	D	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
2	NAG	E	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	0/6/23/26	0/1/1/1
2	BMA	E	3	2	-	0/2/19/22	0/1/1/1
2	MAN	E	4	2	-	0/2/19/22	0/1/1/1
2	MAN	E	5	2	-	1/2/19/22	0/1/1/1
2	NAG	F	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	F	2	2	-	0/6/23/26	0/1/1/1
2	BMA	F	3	2	-	0/2/19/22	0/1/1/1
2	MAN	F	4	2	-	0/2/19/22	0/1/1/1
2	MAN	F	5	2	-	1/2/19/22	0/1/1/1
4	NAG	G	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	G	2	4	-	0/6/23/26	0/1/1/1
4	BMA	G	3	4	-	0/2/19/22	0/1/1/1
5	NAG	H	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	H	2	5	-	0/6/23/26	0/1/1/1
5	BMA	H	3	5	-	0/2/19/22	0/1/1/1
5	MAN	H	4	5	-	1/2/19/22	0/1/1/1
5	MAN	H	5	5	-	1/2/19/22	1/1/1/1

There are no bond length outliers.

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	5	MAN	C1-O5-C5	3.67	117.16	112.19
2	F	4	MAN	C1-O5-C5	3.36	116.75	112.19
2	E	5	MAN	C1-O5-C5	2.92	116.14	112.19
2	C	5	MAN	C1-O5-C5	2.58	115.69	112.19

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	4	MAN	C1-O5-C5	2.57	115.68	112.19

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	H	4	MAN	O5-C5-C6-O6
2	C	5	MAN	O5-C5-C6-O6
2	F	5	MAN	O5-C5-C6-O6
5	H	5	MAN	O5-C5-C6-O6
2	E	5	MAN	O5-C5-C6-O6


All (1) ring outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	H	5	MAN	C1-C2-C3-C4-C5-O5


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











## 5.6 Ligand geometry [i](#)

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	NAG	A	921	1	14,14,15	0.29	0	17,19,21	0.55	0
6	NAG	B	921	1	14,14,15	0.31	0	17,19,21	0.74	1 (5%)
6	NAG	B	901	1	14,14,15	0.30	0	17,19,21	0.55	0
7	QSM	A	922	-	31,31,31	0.97	1 (3%)	35,44,44	0.91	2 (5%)
6	NAG	A	901	1	14,14,15	0.44	0	17,19,21	1.60	2 (11%)
6	NAG	A	908	1	14,14,15	0.31	0	17,19,21	0.40	0
6	NAG	A	902	1	14,14,15	0.30	0	17,19,21	0.59	0
6	NAG	A	918	1	14,14,15	0.28	0	17,19,21	0.46	0
6	NAG	B	919	1	14,14,15	0.30	0	17,19,21	0.50	0
6	NAG	A	919	1	14,14,15	0.30	0	17,19,21	0.66	1 (5%)
6	NAG	A	912	1	14,14,15	0.34	0	17,19,21	0.49	0
6	NAG	A	920	1	14,14,15	0.32	0	17,19,21	0.66	1 (5%)
6	NAG	B	902	1	14,14,15	0.31	0	17,19,21	0.45	0
6	NAG	B	920	1	14,14,15	0.29	0	17,19,21	0.67	1 (5%)
7	QSM	B	922	-	31,31,31	0.96	1 (3%)	35,44,44	0.91	2 (5%)
6	NAG	B	908	1	14,14,15	0.30	0	17,19,21	0.42	0
6	NAG	B	913	1	14,14,15	0.28	0	17,19,21	0.51	0
6	NAG	B	909	1	14,14,15	0.28	0	17,19,21	0.54	0
6	NAG	A	909	1	14,14,15	0.34	0	17,19,21	1.13	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	A	921	1	-	1/6/23/26	0/1/1/1
6	NAG	B	921	1	-	0/6/23/26	0/1/1/1
6	NAG	B	901	1	-	1/6/23/26	0/1/1/1
7	QSM	A	922	-	-	2/16/24/24	0/4/4/4
6	NAG	A	901	1	-	3/6/23/26	0/1/1/1
6	NAG	A	908	1	-	2/6/23/26	0/1/1/1

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	A	902	1	-	0/6/23/26	0/1/1/1
6	NAG	A	918	1	-	1/6/23/26	0/1/1/1
6	NAG	B	919	1	-	1/6/23/26	0/1/1/1
6	NAG	A	919	1	-	0/6/23/26	0/1/1/1
6	NAG	A	912	1	-	2/6/23/26	0/1/1/1
6	NAG	A	920	1	-	0/6/23/26	0/1/1/1
6	NAG	B	902	1	-	0/6/23/26	0/1/1/1
6	NAG	B	920	1	-	0/6/23/26	0/1/1/1
7	QSM	B	922	-	-	2/16/24/24	0/4/4/4
6	NAG	B	908	1	-	2/6/23/26	0/1/1/1
6	NAG	B	913	1	-	0/6/23/26	0/1/1/1
6	NAG	B	909	1	-	0/6/23/26	0/1/1/1
6	NAG	A	909	1	-	1/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	922	QSM	C7-C4	4.27	1.45	1.40
7	B	922	QSM	C7-C4	4.22	1.45	1.40

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	901	NAG	C1-C2-N2	4.91	118.87	110.49
6	A	901	NAG	C2-N2-C7	4.23	128.93	122.90
6	A	909	NAG	C1-O5-C5	3.13	116.43	112.19
7	A	922	QSM	C-C5-C4	-3.09	119.94	122.00
7	B	922	QSM	C-C5-C4	-3.08	119.95	122.00

There are no chirality outliers.


5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	922	QSM	C13-C11-C6-N
7	B	922	QSM	C13-C11-C6-N
7	B	922	QSM	C17-C11-C6-N
6	A	908	NAG	O5-C5-C6-O6
6	B	908	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	738/811 (90%)	1.90	264 (35%) 0 0	34, 105, 188, 224	0
1	B	735/811 (90%)	1.54	214 (29%) 0 0	30, 92, 170, 203	0
All	All	1473/1622 (90%)	1.72	478 (32%) 0 0	30, 96, 181, 224	0

The worst 5 of 478 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	817	LEU	21.2
1	A	750	THR	21.1
1	B	783	PHE	14.2
1	A	805	SER	14.1
1	A	113	GLY	13.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MAN	E	4	11/12	0.66	0.28	83,88,90,92	0
5	MAN	H	5	11/12	0.66	0.19	95,98,100,101	0
2	MAN	C	5	11/12	0.69	0.23	99,102,105,107	0
5	MAN	H	4	11/12	0.72	0.18	82,86,88,92	0
4	BMA	G	3	11/12	0.77	0.20	81,85,86,87	0

*Continued on next page...*




*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MAN	F	5	11/12	0.81	0.22	92,94,95,96	0
2	MAN	E	5	11/12	0.81	0.15	81,84,89,91	0
2	MAN	F	4	11/12	0.81	0.17	89,89,90,91	0
2	MAN	C	4	11/12	0.84	0.21	97,102,107,109	0
4	NAG	G	2	14/15	0.84	0.19	65,69,73,77	0
2	BMA	C	3	11/12	0.85	0.15	90,97,102,102	0
5	BMA	H	3	11/12	0.85	0.13	63,67,73,78	0
2	BMA	E	3	11/12	0.89	0.13	71,75,79,80	0
3	NAG	D	2	14/15	0.89	0.15	51,55,57,57	0
2	BMA	F	3	11/12	0.89	0.15	79,84,89,90	0
2	NAG	F	2	14/15	0.91	0.20	63,65,68,72	0
2	NAG	E	1	14/15	0.91	0.17	44,51,58,59	0
4	NAG	G	1	14/15	0.93	0.18	50,51,55,60	0
5	NAG	H	2	14/15	0.93	0.14	44,49,53,59	0
2	NAG	F	1	14/15	0.94	0.22	47,55,60,61	0
2	NAG	C	1	14/15	0.94	0.22	60,63,65,69	0
5	NAG	H	1	14/15	0.94	0.14	30,37,42,43	0
2	NAG	C	2	14/15	0.95	0.19	72,76,81,84	0
2	NAG	E	2	14/15	0.96	0.11	54,58,62,68	0
3	NAG	D	1	14/15	0.96	0.17	39,43,45,49	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around Chain C:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain F:**




$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)









**Electron density around Chain D:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain G:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	NAG	B	901	14/15	0.30	0.77	174,174,175,176	0
6	NAG	A	901	14/15	0.45	0.37	195,200,203,204	0
6	NAG	B	908	14/15	0.48	0.40	99,108,115,117	0
6	NAG	A	902	14/15	0.51	0.39	128,138,149,149	0
6	NAG	B	919	14/15	0.55	0.39	115,126,136,140	0
6	NAG	B	909	14/15	0.61	0.26	124,127,134,134	0
6	NAG	A	918	14/15	0.63	0.42	120,134,147,151	0
6	NAG	A	921	14/15	0.67	0.48	110,112,122,122	0
6	NAG	A	920	14/15	0.68	0.36	113,123,129,133	0
6	NAG	A	908	14/15	0.69	0.27	87,89,94,97	0
6	NAG	B	921	14/15	0.70	0.30	137,143,151,151	0
6	NAG	B	902	14/15	0.71	0.30	135,139,147,152	0
6	NAG	A	909	14/15	0.74	0.34	118,124,126,128	0
6	NAG	A	912	14/15	0.76	0.29	124,128,131,134	0


*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	NAG	B	913	14/15	0.77	0.23	118,122,124,126	0
6	NAG	A	919	14/15	0.79	0.20	150,153,154,154	0
6	NAG	B	920	14/15	0.79	0.19	154,156,158,159	0
7	QSM	B	922	28/28	0.91	0.24	43,50,58,58	30
7	QSM	A	922	28/28	0.95	0.20	38,44,45,45	30

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.