



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 28, 2022 – 12:06 AM JST

PDB ID : 7VAA
Title : Crystal structure of MiCGT(W93V/V124F/ F191A/R282H) in complex with UDPs
Authors : Zhong, L.; Zhang, Z.M.
Deposited on : 2021-08-27
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.29
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.29


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	470	
1	B	470	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7036 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called UDP-glycosyltransferase 13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	453	Total	C	N	O	S	0	0	0
			3493	2238	594	645	16			
1	B	453	Total	C	N	O	S	0	0	0
			3493	2238	594	645	16			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	93	VAL	TRP	engineered mutation	UNP A0A0M4KE44
A	124	PHE	VAL	engineered mutation	UNP A0A0M4KE44
A	191	ALA	PHE	engineered mutation	UNP A0A0M4KE44
A	282	HIS	ARG	engineered mutation	UNP A0A0M4KE44
B	93	VAL	TRP	engineered mutation	UNP A0A0M4KE44
B	124	PHE	VAL	engineered mutation	UNP A0A0M4KE44
B	191	ALA	PHE	engineered mutation	UNP A0A0M4KE44
B	282	HIS	ARG	engineered mutation	UNP A0A0M4KE44

- Molecule 2 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula: $C_9H_{14}N_2O_{12}P_2$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
2	B	1	Total	C	N	O	P	0	0
			25	9	2	12	2		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: UDP-glycosyltransferase 13



• Molecule 1: UDP-glycosyltransferase 13



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	91.70Å 103.76Å 109.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.50 – 3.10 48.50 – 3.10	Depositor EDS
% Data completeness (in resolution range)	99.6 (48.50-3.10) 99.6 (48.50-3.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.92 (at 3.12Å)	Xtriage
Refinement program	PHENIX 1.9_1692+SVN	Depositor
R, R_{free}	0.245 , 0.289 0.248 , 0.293	Depositor DCC
R_{free} test set	1963 reflections (10.06%)	wwPDB-VP
Wilson B-factor (Å ²)	57.6	Xtriage
Anisotropy	0.783	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 27.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	7036	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.23	0/3578	0.42	0/4869
1	B	0.22	0/3578	0.40	0/4869
All	All	0.23	0/7156	0.41	0/9738

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3493	0	3454	48	0
1	B	3493	0	3454	44	0
2	A	25	0	11	0	0
2	B	25	0	11	1	0
All	All	7036	0	6930	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (91) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:271:GLU:HA	1:A:354:ALA:HA	1.64	0.79
1:B:271:GLU:HA	1:B:354:ALA:HA	1.71	0.73
1:A:319:GLU:O	1:A:341:LYS:NZ	2.20	0.72
1:B:463:TRP:H	1:B:464:LYS:HA	1.57	0.70
1:A:25:THR:HG22	1:A:29:ARG:HE	1.59	0.67
1:A:28:LEU:HD22	1:A:61:LEU:HD22	1.77	0.66
1:A:463:TRP:N	1:A:464:LYS:HA	2.12	0.65
1:B:16:LEU:HD23	1:B:119:ILE:HG23	1.79	0.64
1:B:84:ASN:ND2	1:B:195:ASN:O	2.31	0.63
1:A:156:PRO:HB3	1:A:203:ILE:HG23	1.79	0.63
1:B:463:TRP:N	1:B:464:LYS:HA	2.12	0.62
1:A:147:ARG:NH2	1:A:391:GLU:OE2	2.32	0.62
1:B:319:GLU:O	1:B:341:LYS:NZ	2.25	0.62
1:B:247:GLY:O	1:B:452:SER:OG	2.16	0.62
1:B:298:ILE:HD11	1:B:333:ILE:HG12	1.80	0.61
1:A:313:VAL:HG12	1:A:315:GLN:H	1.67	0.59
1:B:251:PRO:HG2	1:B:347:GLN:HG3	1.86	0.58
1:B:321:LEU:HB2	1:B:330:LEU:HD22	1.85	0.58
1:B:313:VAL:HG12	1:B:315:GLN:H	1.68	0.57
1:A:434:ARG:O	1:A:438:LEU:HB2	2.06	0.56
1:A:82:THR:HG23	1:A:98:ARG:HH22	1.70	0.56
1:A:23:HIS:HA	1:A:143:THR:HG21	1.88	0.56
1:A:29:ARG:HD3	1:A:251:PRO:HA	1.88	0.56
1:B:277:VAL:HG22	1:B:359:VAL:HB	1.88	0.55
1:A:433:VAL:HA	1:A:436:LYS:HE2	1.90	0.53
1:B:97:ARG:NH2	1:B:125:SER:OG	2.42	0.53
1:A:308:VAL:HG11	1:A:321:LEU:HD21	1.92	0.52
1:B:71:LYS:HD2	1:B:106:LEU:HD21	1.92	0.51
1:A:104:ASN:O	1:A:108:SER:OG	2.28	0.51
1:B:171:ASP:HA	1:B:184:VAL:HG22	1.93	0.51
1:B:159:VAL:HG12	1:B:165:PHE:HZ	1.76	0.50
1:B:403:LYS:HD2	1:B:403:LYS:H	1.77	0.49
1:A:340:ILE:HG12	1:A:342:ASN:H	1.77	0.49
1:A:277:VAL:HG22	1:A:359:VAL:HB	1.94	0.49
1:B:283:PHE:HE2	1:B:285:LEU:HD23	1.77	0.49
1:A:321:LEU:HB2	1:A:330:LEU:HD22	1.95	0.48
1:A:379:VAL:HB	1:A:401:TRP:HB3	1.95	0.47
1:A:159:VAL:HG12	1:A:165:PHE:HZ	1.79	0.47
1:B:401:TRP:NE1	1:B:403:LYS:HE3	2.28	0.47
1:A:171:ASP:HA	1:A:184:VAL:HG22	1.95	0.47
1:A:120:ASP:OD2	1:A:122:SER:OG	2.31	0.47
1:B:398:LEU:HD12	1:B:435:ALA:HA	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:31:ALA:HB1	1:A:41:VAL:HG11	1.97	0.47
1:A:260:SER:HB2	1:A:263:VAL:HG23	1.97	0.47
1:B:117:LEU:HD23	1:B:131:ALA:HB2	1.96	0.47
1:B:23:HIS:HA	1:B:143:THR:HG21	1.96	0.47
1:B:433:VAL:HA	1:B:436:LYS:HE2	1.96	0.47
1:A:404:ARG:NH1	1:A:417:GLU:OE1	2.48	0.46
1:A:421:ARG:HD3	1:A:421:ARG:HA	1.68	0.46
1:A:29:ARG:NH2	1:A:369:GLU:OE1	2.42	0.46
1:B:110:ILE:HG21	1:B:114:LEU:HD21	1.98	0.46
1:B:369:GLU:OE1	2:B:501:UDP:O3'	2.32	0.46
1:A:150:SER:HB3	1:A:176:ILE:HG23	1.97	0.45
1:A:153:GLU:OE1	1:A:243:TYR:OH	2.32	0.45
1:A:97:ARG:HH22	1:A:125:SER:HB3	1.82	0.45
1:B:28:LEU:HD22	1:B:61:LEU:HD22	1.99	0.44
1:A:21:MET:O	1:A:25:THR:OG1	2.32	0.44
1:B:329:VAL:O	1:B:333:ILE:HG13	2.18	0.44
1:B:429:ASN:HA	1:B:432:ARG:HG3	2.00	0.44
1:A:25:THR:HG21	1:A:29:ARG:HH21	1.83	0.43
1:B:234:LYS:HE3	1:B:234:LYS:HB2	1.80	0.43
1:A:104:ASN:HB3	1:A:105:PRO:HD3	1.99	0.43
1:A:147:ARG:HH21	1:A:182:VAL:HG22	1.82	0.43
1:A:49:THR:OG1	1:A:51:SER:O	2.31	0.43
1:A:361:HIS:CE1	1:A:363:GLY:H	2.37	0.43
1:B:379:VAL:HB	1:B:401:TRP:HB3	2.01	0.43
1:B:16:LEU:HB3	1:B:119:ILE:HA	2.01	0.43
1:B:155:PHE:O	1:B:159:VAL:HG13	2.19	0.42
1:B:251:PRO:CG	1:B:347:GLN:HG3	2.46	0.42
1:B:194:LEU:HA	1:B:199:THR:HG21	2.00	0.42
1:B:421:ARG:HD3	1:B:421:ARG:HA	1.77	0.42
1:A:298:ILE:HD11	1:A:333:ILE:HG12	2.02	0.42
1:B:187:VAL:HA	1:B:188:PRO:HD2	1.84	0.42
1:A:235:ARG:NH2	1:A:241:PRO:O	2.47	0.42
1:A:66:PRO:HD2	1:B:36:GLN:HG2	2.00	0.42
1:A:279:PHE:HD1	1:A:361:HIS:HB3	1.85	0.42
1:A:457:LYS:HB3	1:A:457:LYS:HE2	1.89	0.42
1:B:393:ILE:O	1:B:396:SER:OG	2.24	0.41
1:A:332:LYS:HE2	1:A:332:LYS:HB3	1.88	0.41
1:B:309:LYS:HE3	1:B:309:LYS:HB3	1.96	0.41
1:B:360:THR:HG21	1:B:377:MET:HE3	2.01	0.41
1:B:142:PHE:CE2	1:B:144:SER:HB3	2.55	0.41
1:A:311:LYS:HB2	1:A:317:ASP:HB2	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:124:PHE:CZ	1:A:212:ALA:HB2	2.56	0.41
1:A:213:ASN:OD1	1:A:213:ASN:N	2.53	0.41
1:A:426:MET:O	1:A:432:ARG:NH1	2.52	0.41
1:B:75:LEU:HD22	1:B:96:ILE:HA	2.03	0.41
1:A:371:VAL:HG22	1:A:398:LEU:HD22	2.03	0.40
1:B:378:LEU:HB2	1:B:425:LEU:HD22	2.03	0.40
1:A:156:PRO:HA	1:A:159:VAL:HG22	2.03	0.40
1:B:230:GLY:O	1:B:234:LYS:HB2	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	449/470 (96%)	425 (95%)	21 (5%)	3 (1%)	22	57
1	B	449/470 (96%)	433 (96%)	16 (4%)	0	100	100
All	All	898/940 (96%)	858 (96%)	37 (4%)	3 (0%)	41	73

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	188	PRO
1	A	172	ASP
1	A	137	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	384/410 (94%)	361 (94%)	23 (6%)	19	49
1	B	384/410 (94%)	374 (97%)	10 (3%)	46	74
All	All	768/820 (94%)	735 (96%)	33 (4%)	29	62

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	LEU
1	A	35	VAL
1	A	82	THR
1	A	161	SER
1	A	164	ASN
1	A	171	ASP
1	A	174	ILE
1	A	213	ASN
1	A	228	LEU
1	A	267	ASP
1	A	358	PHE
1	A	366	SER
1	A	383	PHE
1	A	402	VAL
1	A	412	LEU
1	A	417	GLU
1	A	421	ARG
1	A	432	ARG
1	A	438	LEU
1	A	439	ARG
1	A	440	GLU
1	A	451	SER
1	A	459	LEU
1	B	164	ASN
1	B	171	ASP
1	B	174	ILE
1	B	318	GLU
1	B	361	HIS
1	B	398	LEU
1	B	417	GLU
1	B	421	ARG
1	B	432	ARG
1	B	439	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	365	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	UDP	B	501	-	24,26,26	0.94	0	37,40,40	1.59	7 (18%)
2	UDP	A	501	-	24,26,26	0.95	0	37,40,40	1.59	6 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UDP	B	501	-	-	4/16/32/32	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UDP	A	501	-	-	4/16/32/32	0/2/2/2

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	UDP	C4-N3-C2	-5.00	119.98	126.58
2	A	501	UDP	C4-N3-C2	-4.92	120.10	126.58
2	A	501	UDP	N3-C2-N1	3.96	120.15	114.89
2	B	501	UDP	N3-C2-N1	3.95	120.13	114.89
2	B	501	UDP	C5-C4-N3	3.24	119.69	114.84
2	A	501	UDP	C5-C4-N3	3.22	119.66	114.84
2	A	501	UDP	PA-O3A-PB	-2.96	122.68	132.83
2	B	501	UDP	O4-C4-C5	-2.83	120.18	125.16
2	A	501	UDP	O4-C4-C5	-2.78	120.26	125.16
2	B	501	UDP	PA-O3A-PB	-2.64	123.75	132.83
2	A	501	UDP	C3'-C2'-C1'	2.17	105.56	101.43
2	B	501	UDP	C3'-C2'-C1'	2.07	105.36	101.43
2	B	501	UDP	O2-C2-N1	-2.06	120.05	122.79

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	UDP	C3'-C4'-C5'-O5'
2	A	501	UDP	PA-O3A-PB-O2B
2	B	501	UDP	C3'-C4'-C5'-O5'
2	B	501	UDP	PA-O3A-PB-O2B
2	A	501	UDP	O4'-C4'-C5'-O5'
2	B	501	UDP	O4'-C4'-C5'-O5'
2	A	501	UDP	PA-O3A-PB-O3B
2	B	501	UDP	PA-O3A-PB-O3B


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	UDP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	453/470 (96%)	0.32	9 (1%) 65 44	20, 35, 56, 86	0
1	B	453/470 (96%)	0.87	73 (16%) 1 1	68, 85, 116, 146	0
All	All	906/940 (96%)	0.59	82 (9%) 9 3	20, 69, 103, 146	0

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	330	LEU	11.4
1	B	314	ASP	10.6
1	B	312	LYS	7.2
1	B	315	GLN	7.1
1	B	282	HIS	5.7
1	B	334	LYS	5.6
1	B	320	SER	5.5
1	B	399	GLY	5.3
1	B	327	ARG	4.8
1	B	316	GLU	4.6
1	B	313	VAL	4.6
1	B	398	LEU	4.6
1	B	331	GLU	4.3
1	B	293	LEU	4.2
1	B	227	ILE	4.1
1	B	137	PRO	3.9
1	B	400	MET	3.7
1	A	224	GLU	3.7
1	B	340	ILE	3.7
1	B	306	TRP	3.7
1	B	321	LEU	3.6
1	B	169	GLN	3.6
1	B	173	VAL	3.6
1	B	308	VAL	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	243	TYR	3.5
1	B	310	CYS	3.5
1	B	336	TYR	3.5
1	B	217	ILE	3.4
1	B	80	PRO	3.4
1	A	218	ASN	3.3
1	B	84	ASN	3.2
1	B	297	LEU	3.1
1	B	338	PHE	3.1
1	B	333	ILE	3.1
1	B	224	GLU	3.1
1	B	431	LEU	3.0
1	B	328	ASP	3.0
1	B	165	PHE	3.0
1	B	179	PHE	2.9
1	B	146	THR	2.9
1	B	418	ILE	2.9
1	A	14	ALA	2.8
1	B	339	VAL	2.8
1	B	317	ASP	2.7
1	B	341	LYS	2.7
1	B	228	LEU	2.7
1	A	356	GLY	2.7
1	B	244	CYS	2.7
1	B	198	PHE	2.6
1	B	155	PHE	2.6
1	B	378	LEU	2.6
1	B	401	TRP	2.5
1	B	147	ARG	2.5
1	B	38	HIS	2.5
1	B	192	LEU	2.5
1	B	380	TRP	2.5
1	B	171	ASP	2.5
1	B	387	LYS	2.4
1	B	288	GLU	2.4
1	A	338	PHE	2.4
1	A	116	ALA	2.4
1	B	397	GLY	2.4
1	B	318	GLU	2.4
1	B	346	GLN	2.3
1	B	86	THR	2.3
1	A	108	SER	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	27	CYS	2.3
1	B	170	LEU	2.2
1	B	414	LYS	2.2
1	B	307	VAL	2.2
1	B	226	GLY	2.2
1	B	151	LEU	2.1
1	A	246	VAL	2.1
1	B	231	ILE	2.1
1	B	396	SER	2.1
1	B	175	GLU	2.1
1	B	361	HIS	2.1
1	A	106	LEU	2.1
1	B	385	ASP	2.0
1	B	216	LEU	2.0
1	B	309	LYS	2.0
1	B	391	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	UDP	B	501	25/25	0.76	0.31	78,87,104,113	0
2	UDP	A	501	25/25	0.89	0.26	23,41,48,55	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around UDP B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.