



wwPDB X-ray Structure Validation Summary Report ⓘ

May 18, 2020 – 09:41 pm BST

PDB ID : 2VBI
Title : Holostructure of pyruvate decarboxylase from *Acetobacter pasteurianus*
Authors : Gocke, D.; Berthold, C.L.; Schneider, G.; Pohl, M.
Deposited on : 2007-09-14
Resolution : 2.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	566	84% 12% ..
1	B	566	83% 13% ..
1	C	566	83% 13% ..
1	D	566	82% 14% ..
1	E	566	82% 14% ..
1	F	566	81% 15% ..
1	G	566	83% 13% ..
1	H	566	83% 13% ..

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 33860 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PYRUVATE DECARBOXYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	554	Total	C	N	O	S	0	1	0
			4194	2637	731	801	25			
1	B	554	Total	C	N	O	S	0	1	0
			4194	2637	731	801	25			
1	C	554	Total	C	N	O	S	0	1	0
			4194	2637	731	801	25			
1	D	554	Total	C	N	O	S	0	1	0
			4194	2637	731	801	25			
1	E	554	Total	C	N	O	S	0	1	0
			4194	2637	731	801	25			
1	F	554	Total	C	N	O	S	0	1	0
			4194	2637	731	801	25			
1	G	554	Total	C	N	O	S	0	1	0
			4194	2637	731	801	25			
1	H	554	Total	C	N	O	S	0	1	0
			4194	2637	731	801	25			

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	25	ALA	GLY	conflict	UNP Q8L388
A	207	GLU	LYS	conflict	UNP Q8L388
A	208	LYS	ASN	conflict	UNP Q8L388
A	209	SER	ARG	conflict	UNP Q8L388
A	210	ALA	PRO	conflict	UNP Q8L388
A	211	SER	ALA	conflict	UNP Q8L388
A	297	ALA	GLY	conflict	UNP Q8L388
A	298	TRP	MET	conflict	UNP Q8L388
A	397	PRO	ALA	conflict	UNP Q8L388
B	25	ALA	GLY	conflict	UNP Q8L388
B	207	GLU	LYS	conflict	UNP Q8L388
B	208	LYS	ASN	conflict	UNP Q8L388
B	209	SER	ARG	conflict	UNP Q8L388

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	210	ALA	PRO	conflict	UNP Q8L388
B	211	SER	ALA	conflict	UNP Q8L388
B	297	ALA	GLY	conflict	UNP Q8L388
B	298	TRP	MET	conflict	UNP Q8L388
B	397	PRO	ALA	conflict	UNP Q8L388
C	25	ALA	GLY	conflict	UNP Q8L388
C	207	GLU	LYS	conflict	UNP Q8L388
C	208	LYS	ASN	conflict	UNP Q8L388
C	209	SER	ARG	conflict	UNP Q8L388
C	210	ALA	PRO	conflict	UNP Q8L388
C	211	SER	ALA	conflict	UNP Q8L388
C	297	ALA	GLY	conflict	UNP Q8L388
C	298	TRP	MET	conflict	UNP Q8L388
C	397	PRO	ALA	conflict	UNP Q8L388
D	25	ALA	GLY	conflict	UNP Q8L388
D	207	GLU	LYS	conflict	UNP Q8L388
D	208	LYS	ASN	conflict	UNP Q8L388
D	209	SER	ARG	conflict	UNP Q8L388
D	210	ALA	PRO	conflict	UNP Q8L388
D	211	SER	ALA	conflict	UNP Q8L388
D	297	ALA	GLY	conflict	UNP Q8L388
D	298	TRP	MET	conflict	UNP Q8L388
D	397	PRO	ALA	conflict	UNP Q8L388
E	25	ALA	GLY	conflict	UNP Q8L388
E	207	GLU	LYS	conflict	UNP Q8L388
E	208	LYS	ASN	conflict	UNP Q8L388
E	209	SER	ARG	conflict	UNP Q8L388
E	210	ALA	PRO	conflict	UNP Q8L388
E	211	SER	ALA	conflict	UNP Q8L388
E	297	ALA	GLY	conflict	UNP Q8L388
E	298	TRP	MET	conflict	UNP Q8L388
E	397	PRO	ALA	conflict	UNP Q8L388
F	25	ALA	GLY	conflict	UNP Q8L388
F	207	GLU	LYS	conflict	UNP Q8L388
F	208	LYS	ASN	conflict	UNP Q8L388
F	209	SER	ARG	conflict	UNP Q8L388
F	210	ALA	PRO	conflict	UNP Q8L388
F	211	SER	ALA	conflict	UNP Q8L388
F	297	ALA	GLY	conflict	UNP Q8L388
F	298	TRP	MET	conflict	UNP Q8L388
F	397	PRO	ALA	conflict	UNP Q8L388
G	25	ALA	GLY	conflict	UNP Q8L388

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	207	GLU	LYS	conflict	UNP Q8L388
G	208	LYS	ASN	conflict	UNP Q8L388
G	209	SER	ARG	conflict	UNP Q8L388
G	210	ALA	PRO	conflict	UNP Q8L388
G	211	SER	ALA	conflict	UNP Q8L388
G	297	ALA	GLY	conflict	UNP Q8L388
G	298	TRP	MET	conflict	UNP Q8L388
G	397	PRO	ALA	conflict	UNP Q8L388
H	25	ALA	GLY	conflict	UNP Q8L388
H	207	GLU	LYS	conflict	UNP Q8L388
H	208	LYS	ASN	conflict	UNP Q8L388
H	209	SER	ARG	conflict	UNP Q8L388
H	210	ALA	PRO	conflict	UNP Q8L388
H	211	SER	ALA	conflict	UNP Q8L388
H	297	ALA	GLY	conflict	UNP Q8L388
H	298	TRP	MET	conflict	UNP Q8L388
H	397	PRO	ALA	conflict	UNP Q8L388

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	G	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0
2	E	1	Total Mg 1 1	0	0
2	H	1	Total Mg 1 1	0	0
2	B	1	Total Mg 1 1	0	0
2	C	1	Total Mg 1 1	0	0
2	A	1	Total Mg 1 1	0	0
2	F	1	Total Mg 1 1	0	0

- Molecule 3 is THIAMINE DIPHOSPHATE (three-letter code: TPP) (formula: C₁₂H₁₉N₄O₇P₂S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total 26	C 12	N 4	O 7	P 2	S 1	0	0
3	B	1	Total 26	C 12	N 4	O 7	P 2	S 1	0	0
3	C	1	Total 26	C 12	N 4	O 7	P 2	S 1	0	0
3	D	1	Total 26	C 12	N 4	O 7	P 2	S 1	0	0
3	E	1	Total 26	C 12	N 4	O 7	P 2	S 1	0	0
3	F	1	Total 26	C 12	N 4	O 7	P 2	S 1	0	0
3	G	1	Total 26	C 12	N 4	O 7	P 2	S 1	0	0
3	H	1	Total 26	C 12	N 4	O 7	P 2	S 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	6	Total O 6 6	0	0
4	B	5	Total O 5 5	0	0
4	C	14	Total O 14 14	0	0
4	D	15	Total O 15 15	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	20	Total 20	O 20	0	0
4	F	8	Total 8	O 8	0	0
4	G	11	Total 11	O 11	0	0
4	H	13	Total 13	O 13	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: PYRUVATE DECARBOXYLASE




• Molecule 1: PYRUVATE DECARBOXYLASE




• Molecule 1: PYRUVATE DECARBOXYLASE






• Molecule 1: PYRUVATE DECARBOXYLASE

Chain D: 82% 14% ..




• Molecule 1: PYRUVATE DECARBOXYLASE

Chain E: 82% 14% ..




• Molecule 1: PYRUVATE DECARBOXYLASE

Chain F: 81% 15% ..





• Molecule 1: PYRUVATE DECARBOXYLASE

Chain G: 83% 13% ..



● Molecule 1: PYRUVATE DECARBOXYLASE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	179.62Å 162.43Å 169.33Å 90.00° 102.71° 90.00°	Depositor
Resolution (Å)	30.00 – 2.75	Depositor
% Data completeness (in resolution range)	97.1 (30.00-2.75)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.224 , 0.243	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	33860	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, TPP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/4282	0.55	0/5829
1	B	0.42	0/4282	0.55	0/5829
1	C	0.45	0/4282	0.56	0/5829
1	D	0.46	0/4282	0.57	0/5829
1	E	0.46	0/4282	0.58	0/5829
1	F	0.45	0/4282	0.57	0/5829
1	G	0.42	0/4282	0.56	0/5829
1	H	0.43	0/4282	0.55	0/5829
All	All	0.44	0/34256	0.56	0/46632

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4194	0	4148	39	0
1	B	4194	0	4148	38	0
1	C	4194	0	4148	42	0
1	D	4194	0	4148	44	2
1	E	4194	0	4148	46	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	4194	0	4148	51	1
1	G	4194	0	4148	43	0
1	H	4194	0	4148	43	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
3	A	26	0	16	1	0
3	B	26	0	16	3	0
3	C	26	0	16	0	0
3	D	26	0	16	1	0
3	E	26	0	16	1	0
3	F	26	0	16	1	0
3	G	26	0	16	2	0
3	H	26	0	16	3	0
4	A	6	0	0	1	0
4	B	5	0	0	1	0
4	C	14	0	0	3	0
4	D	15	0	0	1	0
4	E	20	0	0	2	0
4	F	8	0	0	2	0
4	G	11	0	0	1	0
4	H	13	0	0	2	0
All	All	33860	0	33312	300	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 300 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:137:SER:HA	1:D:165:ASP:HB2	1.47	0.96
1:E:137:SER:HA	1:E:165:ASP:HB2	1.49	0.95
1:G:137:SER:HA	1:G:165:ASP:HB2	1.48	0.95
1:B:137:SER:HA	1:B:165:ASP:HB2	1.50	0.94
1:A:137:SER:HA	1:A:165:ASP:HB2	1.49	0.92

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:360:GLY:N	1:F:533:ARG:NH1[3_555]	1.95	0.25
1:D:509:GLU:OE2	1:E:496:GLU:CG[3_555]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	553/566 (98%)	538 (97%)	14 (2%)	1 (0%)	47	69
1	B	553/566 (98%)	539 (98%)	13 (2%)	1 (0%)	47	69
1	C	553/566 (98%)	538 (97%)	14 (2%)	1 (0%)	47	69
1	D	553/566 (98%)	538 (97%)	14 (2%)	1 (0%)	47	69
1	E	553/566 (98%)	538 (97%)	14 (2%)	1 (0%)	47	69
1	F	553/566 (98%)	539 (98%)	13 (2%)	1 (0%)	47	69
1	G	553/566 (98%)	538 (97%)	14 (2%)	1 (0%)	47	69
1	H	553/566 (98%)	538 (97%)	14 (2%)	1 (0%)	47	69
All	All	4424/4528 (98%)	4306 (97%)	110 (2%)	8 (0%)	47	69

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	74	SER
1	B	74	SER
1	C	74	SER
1	D	74	SER
1	E	74	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	440/450 (98%)	404 (92%)	36 (8%)	11	20
1	B	440/450 (98%)	404 (92%)	36 (8%)	11	20
1	C	440/450 (98%)	404 (92%)	36 (8%)	11	20
1	D	440/450 (98%)	404 (92%)	36 (8%)	11	20
1	E	440/450 (98%)	405 (92%)	35 (8%)	12	21
1	F	440/450 (98%)	403 (92%)	37 (8%)	11	19
1	G	440/450 (98%)	404 (92%)	36 (8%)	11	20
1	H	440/450 (98%)	404 (92%)	36 (8%)	11	20
All	All	3520/3600 (98%)	3232 (92%)	288 (8%)	11	20

5 of 288 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	320	ASP
1	E	276	THR
1	H	272	GLU
1	D	371	ASN
1	E	106	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 59 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	329	GLN
1	E	329	GLN
1	H	64	ASN
1	D	430	HIS
1	E	44	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	TPP	H	2000	2	22,27,27	1.44	5 (22%)	29,40,40	1.83	8 (27%)
3	TPP	C	2000	2	22,27,27	1.69	5 (22%)	29,40,40	1.73	8 (27%)
3	TPP	A	2000	2	22,27,27	1.45	4 (18%)	29,40,40	1.86	9 (31%)
3	TPP	F	2000	2	22,27,27	1.52	4 (18%)	29,40,40	1.89	7 (24%)
3	TPP	G	2000	2	22,27,27	1.38	4 (18%)	29,40,40	1.80	8 (27%)
3	TPP	D	2000	2	22,27,27	1.47	5 (22%)	29,40,40	1.63	6 (20%)
3	TPP	E	2000	2	22,27,27	1.46	6 (27%)	29,40,40	1.72	8 (27%)
3	TPP	B	2000	2	22,27,27	1.56	5 (22%)	29,40,40	1.76	7 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TPP	H	2000	2	-	0/16/17/17	0/2/2/2
3	TPP	C	2000	2	-	2/16/17/17	0/2/2/2
3	TPP	A	2000	2	-	0/16/17/17	0/2/2/2
3	TPP	F	2000	2	-	1/16/17/17	0/2/2/2
3	TPP	G	2000	2	-	3/16/17/17	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TPP	D	2000	2	-	5/16/17/17	0/2/2/2
3	TPP	E	2000	2	-	0/16/17/17	0/2/2/2
3	TPP	B	2000	2	-	1/16/17/17	0/2/2/2

The worst 5 of 38 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	2000	TPP	C4-N3	-4.33	1.35	1.39
3	C	2000	TPP	C4-N3	-4.05	1.36	1.39
3	C	2000	TPP	C2'-N3'	3.52	1.40	1.34
3	C	2000	TPP	C4'-N3'	3.16	1.39	1.35
3	E	2000	TPP	C4-N3	-3.15	1.37	1.39

The worst 5 of 61 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	2000	TPP	C6-C5-C4	6.57	132.71	127.43
3	A	2000	TPP	C6-C5-C4	5.07	131.50	127.43
3	B	2000	TPP	C6-C5-C4	4.81	131.29	127.43
3	H	2000	TPP	C6-C5-C4	4.50	131.04	127.43
3	C	2000	TPP	C6-C5-C4	4.49	131.04	127.43

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	2000	TPP	PA-O3A-PB-O3B
3	G	2000	TPP	C5-C6-C7-O7
3	G	2000	TPP	C7-O7-PA-O1A
3	D	2000	TPP	C7-O7-PA-O1A
3	D	2000	TPP	C7-O7-PA-O2A

There are no ring outliers.

7 monomers are involved in 12 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	H	2000	TPP	3	0
3	A	2000	TPP	1	0
3	F	2000	TPP	1	0
3	G	2000	TPP	2	0
3	D	2000	TPP	1	0

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	2000	TPP	1	0
3	B	2000	TPP	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





Ligand TPP C 2000




Ligand TPP A 2000







Ligand TPP D 2000



Ligand TPP E 2000





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.