



wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 08:35 am BST

PDB ID : 5VEI
Title : Crystal structure of the SH3 domain of human sorbin and SH3 domain-containing protein 2
Authors : Liu, Y.; Tempel, W.; Huang, H.; Gu, J.; Liu, K.; Sidhu, S.S.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.M.; Min, J.; Structural Genomics Consortium (SGC)
Deposited on : 2017-04-04
Resolution : 1.33 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1385 (1.36-1.32)
Clashscore	141614	1417 (1.36-1.32)
Ramachandran outliers	138981	1397 (1.36-1.32)
Sidechain outliers	138945	1397 (1.36-1.32)
RSRZ outliers	127900	1369 (1.36-1.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	110	<div> <div>4%</div> <div>69%</div> <div>.</div> <div>27%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 700 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sorbin and SH3 domain-containing protein 2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	A	80	651	423	106	122	0	15	0

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	832	MET	-	initiating methionine	UNP H7C1R7
A	833	LYS	-	expression tag	UNP H7C1R7
A	834	ILE	-	expression tag	UNP H7C1R7
A	835	GLU	-	expression tag	UNP H7C1R7
A	836	GLU	-	expression tag	UNP H7C1R7
A	837	HIS	-	expression tag	UNP H7C1R7
A	838	HIS	-	expression tag	UNP H7C1R7
A	839	HIS	-	expression tag	UNP H7C1R7
A	840	HIS	-	expression tag	UNP H7C1R7
A	841	HIS	-	expression tag	UNP H7C1R7
A	842	HIS	-	expression tag	UNP H7C1R7
A	843	SER	-	expression tag	UNP H7C1R7
A	844	SER	-	expression tag	UNP H7C1R7
A	845	GLY	-	expression tag	UNP H7C1R7
A	846	ARG	-	expression tag	UNP H7C1R7
A	847	GLU	-	expression tag	UNP H7C1R7
A	848	ASN	-	expression tag	UNP H7C1R7
A	849	LEU	-	expression tag	UNP H7C1R7
A	850	TYR	-	expression tag	UNP H7C1R7
A	851	PHE	-	expression tag	UNP H7C1R7
A	852	GLN	-	expression tag	UNP H7C1R7
A	853	GLY	-	expression tag	UNP H7C1R7
A	854	GLY	-	expression tag	UNP H7C1R7
A	855	ALA	-	expression tag	UNP H7C1R7
A	856	ALA	-	expression tag	UNP H7C1R7
A	857	GLN	-	expression tag	UNP H7C1R7
A	858	PRO	-	expression tag	UNP H7C1R7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	859	ALA	-	expression tag	UNP H7C1R7
A	860	MET	-	expression tag	UNP H7C1R7
A	861	ALA	-	expression tag	UNP H7C1R7
A	862	GLN	-	expression tag	UNP H7C1R7
A	863	GLY	-	expression tag	UNP H7C1R7
A	864	ALA	-	expression tag	UNP H7C1R7
A	865	LEU	-	expression tag	UNP H7C1R7
A	922	GLY	-	expression tag	UNP H7C1R7
A	923	SER	-	expression tag	UNP H7C1R7
A	924	ALA	-	expression tag	UNP H7C1R7
A	925	ALA	-	expression tag	UNP H7C1R7
A	926	ALA	-	expression tag	UNP H7C1R7
A	927	LEU	-	expression tag	UNP H7C1R7
A	928	ARG	-	expression tag	UNP H7C1R7
A	929	THR	-	expression tag	UNP H7C1R7
A	930	GLY	-	expression tag	UNP H7C1R7
A	931	GLU	-	expression tag	UNP H7C1R7
A	932	ALA	-	expression tag	UNP H7C1R7
A	933	TYR	-	expression tag	UNP H7C1R7
A	934	LEU	-	expression tag	UNP H7C1R7
A	935	ARG	-	expression tag	UNP H7C1R7
A	936	TYR	-	expression tag	UNP H7C1R7
A	937	VAL	-	expression tag	UNP H7C1R7
A	938	ASP	-	expression tag	UNP H7C1R7
A	939	ALA	-	expression tag	UNP H7C1R7
A	940	ALA	-	expression tag	UNP H7C1R7
A	941	ALA	-	expression tag	UNP H7C1R7

- Molecule 2 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	27	Total X 27 27	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	21	Total O 22 22	0	2

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Sorbin and SH3 domain-containing protein 2

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	46.20Å 27.49Å 58.83Å 90.00° 101.01° 90.00°	Depositor
Resolution (Å)	12.62 – 1.33 23.51 – 1.33	Depositor EDS
% Data completeness (in resolution range)	99.7 (12.62-1.33) 99.6 (23.51-1.33)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 1.33Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.185 , 0.206 0.200 , 0.219	Depositor DCC
R_{free} test set	819 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	14.6	Xtriage
Anisotropy	0.559	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 66.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	700	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UNX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/678	0.68	0/917

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	651	0	612	2	0
2	A	27	0	0	0	0
3	A	22	0	0	0	0
All	All	700	0	612	2	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (2) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:865:LEU:HB3	1:A:891:TYR:HB3	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:867:PRO:HB2	1:A:920[A]:LEU:HD12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	86/110 (78%)	84 (98%)	2 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	60/85 (71%)	60 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 27 ligands modelled in this entry, 27 are unknown - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	80/110 (72%)	0.09	4 (5%)	28 33	12, 20, 29, 34	7 (8%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	864	ALA	3.7
1	A	863[A]	GLY	2.8
1	A	862[A]	GLN	2.2
1	A	941[A]	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	UNX	A	1008	1/1	0.62	0.36	37,37,37,37	0
2	UNX	A	1015	1/1	0.75	0.10	44,44,44,44	0
2	UNX	A	1017	1/1	0.75	0.26	35,35,35,35	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	UNX	A	1023	1/1	0.83	0.08	33,33,33,33	0
2	UNX	A	1013	1/1	0.85	0.27	32,32,32,32	1
2	UNX	A	1004	1/1	0.86	0.10	29,29,29,29	0
2	UNX	A	1012	1/1	0.88	0.07	36,36,36,36	0
2	UNX	A	1027	1/1	0.90	0.15	30,30,30,30	0
2	UNX	A	1002	1/1	0.90	0.11	31,31,31,31	0
2	UNX	A	1019	1/1	0.90	0.08	26,26,26,26	0
2	UNX	A	1018	1/1	0.91	0.08	34,34,34,34	0
2	UNX	A	1021	1/1	0.91	0.13	23,23,23,23	0
2	UNX	A	1022	1/1	0.92	0.15	29,29,29,29	0
2	UNX	A	1003	1/1	0.92	0.21	30,30,30,30	0
2	UNX	A	1014	1/1	0.94	0.06	28,28,28,28	0
2	UNX	A	1001	1/1	0.95	0.09	27,27,27,27	0
2	UNX	A	1007	1/1	0.95	0.13	22,22,22,22	0
2	UNX	A	1005	1/1	0.95	0.08	30,30,30,30	0
2	UNX	A	1011	1/1	0.96	0.08	24,24,24,24	0
2	UNX	A	1016	1/1	0.96	0.06	26,26,26,26	0
2	UNX	A	1024	1/1	0.96	0.23	22,22,22,22	0
2	UNX	A	1020	1/1	0.97	0.15	21,21,21,21	0
2	UNX	A	1006	1/1	0.97	0.13	19,19,19,19	0
2	UNX	A	1010	1/1	0.97	0.06	20,20,20,20	0
2	UNX	A	1009	1/1	0.98	0.07	31,31,31,31	0
2	UNX	A	1026	1/1	0.98	0.09	19,19,19,19	0
2	UNX	A	1025	1/1	1.00	0.18	12,12,12,12	0

6.5 Other polymers ⓘ

There are no such residues in this entry.