



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 14, 2020 – 05:52 am BST

PDB ID : 3VGU  
Title : E134A mutant nucleoside diphosphate kinase derived from Halomonas sp. 593  
Authors : Okazaki, N.; Yonezawa, Y.; Arai, S.; Matsumoto, F.; Tamada, T.; Tokunaga, H.; Ishibashi, M.; Tokunaga, M.; Kuroki, R.  
Deposited on : 2011-08-21  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	141	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, green 1%, green 84%, yellow 84%, yellow 96%, grey 96%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> <span>84%</span> <span>16%</span> </div> </div>
1	B	141	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 2%, green 2%, green 78%, yellow 78%, yellow 98%, grey 98%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> <span>78%</span> <span>20%</span> </div> </div>
1	C	141	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 9%, green 9%, green 79%, yellow 79%, yellow 97%, grey 97%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> <span>79%</span> <span>18%</span> </div> </div>
1	D	141	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 4%, green 4%, green 77%, yellow 77%, yellow 98%, grey 98%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> <span>77%</span> <span>21%</span> </div> </div>
1	E	141	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 6%, green 6%, green 79%, yellow 79%, yellow 97%, grey 97%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> <span>79%</span> <span>18%</span> </div> </div>
1	F	141	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 4%, green 4%, green 80%, yellow 80%, yellow 97%, grey 97%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> <span>80%</span> <span>17%</span> </div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	141	<div><div>6%</div><div><div></div><div>75%</div><div>21%</div><div></div></div><div>• •</div></div>
1	H	141	<div><div>9%</div><div><div></div><div>71%</div><div>26%</div><div></div></div><div>• •</div></div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8894 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleoside diphosphate kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	140	Total	C	N	O	S	0	0	0
			1060	661	182	213	4			
1	B	140	Total	C	N	O	S	0	0	0
			1060	661	182	213	4			
1	C	140	Total	C	N	O	S	0	0	0
			1060	661	182	213	4			
1	D	140	Total	C	N	O	S	0	0	0
			1060	661	182	213	4			
1	E	140	Total	C	N	O	S	0	0	0
			1060	661	182	213	4			
1	F	140	Total	C	N	O	S	0	0	0
			1060	661	182	213	4			
1	G	140	Total	C	N	O	S	0	0	0
			1060	661	182	213	4			
1	H	138	Total	C	N	O	S	0	0	0
			1042	650	177	211	4			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	134	ALA	GLU	ENGINEERED MUTATION	UNP Q83WH5
B	134	ALA	GLU	ENGINEERED MUTATION	UNP Q83WH5
C	134	ALA	GLU	ENGINEERED MUTATION	UNP Q83WH5
D	134	ALA	GLU	ENGINEERED MUTATION	UNP Q83WH5
E	134	ALA	GLU	ENGINEERED MUTATION	UNP Q83WH5
F	134	ALA	GLU	ENGINEERED MUTATION	UNP Q83WH5
G	134	ALA	GLU	ENGINEERED MUTATION	UNP Q83WH5
H	134	ALA	GLU	ENGINEERED MUTATION	UNP Q83WH5


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	57	Total	O	0	0
			57	57		
2	B	69	Total	O	0	0
			69	69		
2	C	37	Total	O	0	0
			37	37		
2	D	58	Total	O	0	0
			58	58		
2	E	46	Total	O	0	0
			46	46		
2	F	63	Total	O	0	0
			63	63		
2	G	59	Total	O	0	0
			59	59		
2	H	43	Total	O	0	0
			43	43		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Nucleoside diphosphate kinase





- Molecule 1: Nucleoside diphosphate kinase




- Molecule 1: Nucleoside diphosphate kinase




- Molecule 1: Nucleoside diphosphate kinase




- Molecule 1: Nucleoside diphosphate kinase




- Molecule 1: Nucleoside diphosphate kinase



- Molecule 1: Nucleoside diphosphate kinase



- Molecule 1: Nucleoside diphosphate kinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	186.49Å 93.15Å 68.41Å 90.00° 103.60° 90.00°	Depositor
Resolution (Å)	22.98 – 2.30 22.98 – 2.30	Depositor EDS
% Data completeness (in resolution range)	93.2 (22.98-2.30) 93.3 (22.98-2.30)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.26 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.5.0070	Depositor
R, $R_{free}$	0.189 , 0.235 0.185 , 0.227	Depositor DCC
$R_{free}$ test set	2402 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.3	Xtriage
Anisotropy	0.227	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 47.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8894	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.40% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.00	1/1076 (0.1%)	0.81	0/1450
1	B	1.04	1/1076 (0.1%)	0.78	0/1450
1	C	1.00	0/1076	0.81	1/1450 (0.1%)
1	D	0.97	1/1076 (0.1%)	0.79	0/1450
1	E	0.98	0/1076	0.80	0/1450
1	F	1.00	0/1076	0.88	1/1450 (0.1%)
1	G	1.05	2/1076 (0.2%)	0.90	2/1450 (0.1%)
1	H	1.00	1/1056 (0.1%)	0.82	2/1421 (0.1%)
All	All	1.01	6/8588 (0.1%)	0.82	6/11571 (0.1%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	45	GLU	CG-CD	7.17	1.62	1.51
1	G	45	GLU	CB-CG	6.71	1.64	1.52
1	B	15	VAL	CB-CG1	-5.28	1.41	1.52
1	D	15	VAL	CB-CG1	-5.26	1.41	1.52
1	A	78	GLU	CD-OE1	-5.25	1.19	1.25

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	26	ARG	NE-CZ-NH1	6.43	123.52	120.30
1	F	40	LEU	CA-CB-CG	6.08	129.28	115.30
1	C	40	LEU	CA-CB-CG	5.72	128.46	115.30
1	G	26	ARG	NE-CZ-NH2	-5.64	117.48	120.30
1	H	106	ASP	CB-CG-OD2	5.52	123.27	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1060	0	1035	13	0
1	B	1060	0	1035	26	0
1	C	1060	0	1035	15	0
1	D	1060	0	1035	22	0
1	E	1060	0	1035	20	0
1	F	1060	0	1035	20	0
1	G	1060	0	1035	23	0
1	H	1042	0	1014	25	0
2	A	57	0	0	0	0
2	B	69	0	0	1	0
2	C	37	0	0	3	0
2	D	58	0	0	3	0
2	E	46	0	0	1	0
2	F	63	0	0	3	0
2	G	59	0	0	1	0
2	H	43	0	0	1	0
All	All	8894	0	8259	153	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 153 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:15:VAL:HG21	1:A:71:PRO:O	1.73	0.88
1:B:40:LEU:HD23	1:B:40:LEU:N	1.92	0.85
1:B:57:ARG:HH21	1:B:58:PRO:HD2	1.44	0.83
1:B:18:ASN:ND2	1:F:141:ARG:HH12	1.80	0.79
1:C:26:ARG:HD2	2:C:149:HOH:O	1.85	0.75

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	138/141 (98%)	135 (98%)	3 (2%)	0	100	100
1	B	138/141 (98%)	132 (96%)	6 (4%)	0	100	100
1	C	138/141 (98%)	131 (95%)	7 (5%)	0	100	100
1	D	138/141 (98%)	131 (95%)	7 (5%)	0	100	100
1	E	138/141 (98%)	131 (95%)	6 (4%)	1 (1%)	22	26
1	F	138/141 (98%)	134 (97%)	4 (3%)	0	100	100
1	G	138/141 (98%)	132 (96%)	5 (4%)	1 (1%)	22	26
1	H	134/141 (95%)	132 (98%)	2 (2%)	0	100	100
All	All	1100/1128 (98%)	1058 (96%)	40 (4%)	2 (0%)	47	58

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	81	ASN
1	G	109	GLN

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	108/109 (99%)	105 (97%)	3 (3%)	43	60
1	B	108/109 (99%)	103 (95%)	5 (5%)	27	38
1	C	108/109 (99%)	101 (94%)	7 (6%)	17	23

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	108/109 (99%)	105 (97%)	3 (3%)	43	60
1	E	108/109 (99%)	105 (97%)	3 (3%)	43	60
1	F	108/109 (99%)	104 (96%)	4 (4%)	34	48
1	G	108/109 (99%)	102 (94%)	6 (6%)	21	29
1	H	106/109 (97%)	100 (94%)	6 (6%)	20	28
All	All	862/872 (99%)	825 (96%)	37 (4%)	29	40

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	112	ASP
1	E	114	ASN
1	H	62	ASP
1	D	123	GLU
1	E	3	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	86	ASN
1	F	81	ASN
1	G	109	GLN
1	E	81	ASN
1	H	81	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	140/141 (99%)	-0.03	2 (1%) 75 80	15, 24, 43, 50	0
1	B	140/141 (99%)	0.13	3 (2%) 63 70	13, 25, 43, 50	0
1	C	140/141 (99%)	0.39	13 (9%) 8 11	15, 29, 52, 65	0
1	D	140/141 (99%)	0.08	6 (4%) 35 42	14, 26, 43, 50	0
1	E	140/141 (99%)	0.39	9 (6%) 19 25	17, 31, 55, 62	0
1	F	140/141 (99%)	0.06	6 (4%) 35 42	11, 22, 45, 59	0
1	G	140/141 (99%)	0.22	8 (5%) 23 30	12, 24, 46, 50	0
1	H	138/141 (97%)	0.27	13 (9%) 8 11	13, 24, 65, 81	0
All	All	1118/1128 (99%)	0.19	60 (5%) 25 32	11, 26, 48, 81	0

The worst 5 of 60 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2	ALA	5.1
1	C	58	PRO	5.0
1	E	2	ALA	4.6
1	C	109	GLN	4.2
1	H	59	PHE	4.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.