



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 05:55 am BST

PDB ID : 1VHO
Title : Crystal structure of a putative peptidase/endoglucanase
Authors : Structural GenomiX
Deposited on : 2003-12-01
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	346	<div> <div>3%</div> <div>79%</div> <div>12%</div> <div>8%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	A	353	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2717 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called endoglucanase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	319	2454	1569	410	465	3	7	0	4	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MSE	-	cloning artifact	UNP Q9X0D8
A	0	SER	-	cloning artifact	UNP Q9X0D8
A	1	LEU	-	cloning artifact	UNP Q9X0D8
A	3	MSE	MET	modified residue	UNP Q9X0D8
A	10	MSE	MET	modified residue	UNP Q9X0D8
A	114	MSE	MET	modified residue	UNP Q9X0D8
A	129	PRO	LEU	variant	UNP Q9X0D8
A	144	GLY	ASP	variant	UNP Q9X0D8
A	224	MSE	MET	modified residue	UNP Q9X0D8
A	306	MSE	MET	modified residue	UNP Q9X0D8
A	312	MSE	MET	modified residue	UNP Q9X0D8
A	335	GLU	-	cloning artifact	UNP Q9X0D8
A	336	GLY	-	cloning artifact	UNP Q9X0D8
A	337	GLY	-	cloning artifact	UNP Q9X0D8
A	338	SER	-	cloning artifact	UNP Q9X0D8
A	339	HIS	-	cloning artifact	UNP Q9X0D8
A	340	HIS	-	cloning artifact	UNP Q9X0D8
A	341	HIS	-	cloning artifact	UNP Q9X0D8
A	342	HIS	-	cloning artifact	UNP Q9X0D8
A	343	HIS	-	cloning artifact	UNP Q9X0D8
A	344	HIS	-	cloning artifact	UNP Q9X0D8

- Molecule 2 is CACODYLATE ION (three-letter code: CAC) (formula: $C_2H_6AsO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	As	C	O	0	0
			4	1	2	1		
2	A	1	Total	As	C	O	0	0
			4	1	2	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	245	Total 245	O 245	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: endoglucanase



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	120.45Å 120.45Å 57.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.59 – 1.86 42.59 – 1.86	Depositor EDS
% Data completeness (in resolution range)	(Not available) (42.59-1.86) 99.2 (42.59-1.86)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.34 (at 1.86Å)	Xtriage
Refinement program	REFMAC 4.0	Depositor
R, R_{free}	0.212 , 0.245 0.193 , 0.227	Depositor DCC
R_{free} test set	1796 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	25.2	Xtriage
Anisotropy	0.094	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 53.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2717	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CAC, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	0/2503	1.14	2/3379 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	172	ARG	NE-CZ-NH2	-7.36	116.62	120.30
1	A	279	ARG	CD-NE-CZ	5.80	131.73	123.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2454	0	2476	27	0
2	A	8	0	0	2	0
3	A	10	0	0	2	0
4	A	245	0	0	4	0
All	All	2717	0	2476	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:36:PHE:HA	1:A:186:LYS:HE3	1.59	0.84
1:A:199:SER:HB2	2:A:351:CAC:C2	2.17	0.75
1:A:206:CYS:SG	1:A:285:ASP:HB3	2.35	0.67
1:A:146:ARG:HD3	4:A:513:HOH:O	1.97	0.64
1:A:228:PHE:HB2	1:A:303:LEU:O	1.99	0.62
1:A:36:PHE:HD1	1:A:186:LYS:HG3	1.67	0.58
1:A:96:ALA:CB	1:A:153:ILE:HD11	2.38	0.54
1:A:315:PRO:O	1:A:319:GLU:HG3	2.09	0.52
1:A:284:THR:HG22	1:A:287:VAL:HG23	1.93	0.51
1:A:96:ALA:HB2	1:A:153:ILE:HD11	1.94	0.48
1:A:251:VAL:HG22	4:A:460:HOH:O	2.13	0.48
1:A:105:ASN:ND2	3:A:353:SO4:O1	2.48	0.46
1:A:153:ILE:HD12	4:A:446:HOH:O	2.16	0.45
1:A:36:PHE:CD1	1:A:186:LYS:HG3	2.50	0.45
1:A:178:LEU:HB3	1:A:196:VAL:CG1	2.48	0.44
1:A:141:CYS:HA	2:A:350:CAC:C1	2.48	0.43
1:A:287:VAL:O	1:A:290:VAL:HG22	2.18	0.43
1:A:190:HIS:HA	1:A:191:PRO:HD3	1.77	0.43
1:A:143:ARG:HB3	3:A:353:SO4:O1	2.19	0.43
1:A:225:ASP:OD2	1:A:278:GLY:HA2	2.20	0.42
1:A:34:GLU:HB2	1:A:35:PRO:HD3	2.01	0.42
1:A:114:MSE:CE	1:A:140:LEU:HD13	2.50	0.41
1:A:3[B]:MSE:HE2	4:A:595:HOH:O	2.19	0.41
1:A:284:THR:CG2	1:A:287:VAL:HG23	2.50	0.41
1:A:63:PHE:CD2	1:A:206:CYS:HB3	2.56	0.41
1:A:249:GLY:O	1:A:252:VAL:HG22	2.21	0.41
1:A:239:GLU:OE2	1:A:242:LYS:NZ	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	315/346 (91%)	303 (96%)	10 (3%)	2 (1%)	25 12

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	131	TYR
1	A	228	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	270/287 (94%)	261 (97%)	9 (3%)	38 21

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	THR
1	A	95	TYR
1	A	104	LYS
1	A	138	LEU
1	A	143	ARG
1	A	207	LEU
1	A	273	GLU
1	A	333[A]	GLU
1	A	333[B]	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	105	ASN
1	A	161	ASN
1	A	255	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CAC	A	351	1	0,3,4	0.00	-	0,3,6	0.00	-
3	SO4	A	353	-	4,4,4	0.58	0	6,6,6	0.13	0
3	SO4	A	352	-	4,4,4	0.70	0	6,6,6	0.38	0
2	CAC	A	350	1	0,3,4	0.00	-	0,3,6	0.00	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	351	CAC	1	0
3	A	353	SO4	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	350	CAC	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	313/346 (90%)	0.28	9 (2%)	51 50	12, 24, 45, 60	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	131	TYR	6.7
1	A	132	ASP	5.6
1	A	95	TYR	5.1
1	A	130	THR	4.7
1	A	96	ALA	3.9
1	A	44	ARG	2.9
1	A	228	PHE	2.9
1	A	290	VAL	2.8
1	A	134	ILE	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	A	353	5/5	0.80	0.31	79,79,79,80	0
3	SO4	A	352	5/5	0.84	0.20	57,58,59,60	0
2	CAC	A	350	4/5	0.96	0.19	51,52,53,53	0
2	CAC	A	351	4/5	0.97	0.14	31,33,34,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.