



Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 03:54 pm BST

PDB ID : 1VIF
Title : STRUCTURE OF DIHYDROFOLATE REDUCTASE
Authors : Narayana, N.; Matthews, D.A.; Howell, E.E.; Xuong, N.-H.
Deposited on : 1996-10-03
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	62	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FOL	A	1[A]	-	X	-	-
2	FOL	A	1[B]	-	X	-	-

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 527 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DIHYDROFOLATE REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	60	Total	C	N	O	S	0	0	0
			457	287	81	87	2			

- Molecule 2 is FOLIC ACID (three-letter code: FOL) (formula: C₁₉H₁₉N₇O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	1
			26	14	10	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	44	Total	O	0	0
			44	44		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: DIHYDROFOLATE REDUCTASE

Chain A: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	68.75 Å 68.75 Å 52.59 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.80	Depositor
% Data completeness (in resolution range)	100.0 (10.00-1.80)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.176 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	527	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: FOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.12	5/469 (1.1%)	1.44	7/636 (1.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	20	SER	C-N	10.36	1.57	1.34
1	A	73	ALA	C-N	-6.24	1.19	1.34
1	A	58	GLU	CD-OE2	5.91	1.32	1.25
1	A	53	GLU	CD-OE1	-5.60	1.19	1.25
1	A	60	GLU	CD-OE2	5.24	1.31	1.25

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	76	ARG	NE-CZ-NH1	8.33	124.47	120.30
1	A	28	ASP	CB-CG-OD2	7.59	125.13	118.30
1	A	31	ARG	NE-CZ-NH1	7.19	123.89	120.30
1	A	28	ASP	CB-CG-OD1	-6.76	112.22	118.30
1	A	31	ARG	NE-CZ-NH2	-5.53	117.53	120.30
1	A	78	ASN	N-CA-CB	5.43	120.37	110.60
1	A	45	TRP	N-CA-CB	-5.33	101.01	110.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	74	LEU	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	457	0	436	13	2
2	A	26	0	8	6	0
3	A	44	0	0	1	1
All	All	527	0	444	14	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67:GLN:HG3	2:A:1[B]:FOL:C4	2.06	0.85
2:A:1[A]:FOL:N5	3:A:121:HOH:O	2.16	0.79
1:A:67:GLN:HG3	2:A:1[A]:FOL:C4	2.16	0.76
1:A:19:PRO:C	1:A:21:ASN:H	2.10	0.54
1:A:33:LYS:HD2	1:A:74:LEU:C	2.31	0.50
1:A:45:TRP:HZ3	1:A:53:GLU:HG3	1.77	0.48
1:A:19:PRO:HG2	1:A:22:ALA:HB2	1.97	0.47
1:A:19:PRO:C	1:A:21:ASN:N	2.67	0.46
1:A:19:PRO:O	1:A:21:ASN:N	2.48	0.46
1:A:67:GLN:HG3	2:A:1[B]:FOL:C4A	2.46	0.45
1:A:67:GLN:HG3	2:A:1[A]:FOL:N3	2.28	0.43
1:A:19:PRO:HG2	1:A:22:ALA:CB	2.49	0.42
1:A:31:ARG:NE	1:A:75:GLU:OE2	2.53	0.41
1:A:67:GLN:HG3	2:A:1[B]:FOL:O4	2.19	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:21:ASN:ND2	1:A:21:ASN:ND2[8_667]	1.47	0.73
1:A:48:THR:OG1	3:A:111:HOH:O[13_646]	1.65	0.55

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	58/62 (94%)	55 (95%)	1 (2%)	2 (3%)	3 0

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	20	SER
1	A	21	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	46/48 (96%)	38 (83%)	8 (17%)	2 0

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	SER
1	A	21	ASN
1	A	33	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	41	GLN
1	A	45	TRP
1	A	50	LEU
1	A	67	GLN
1	A	74	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	GLN
1	A	41	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FOL	A	1[A]	-	14,14,34	4.39	7 (50%)	17,20,47	3.22	12 (70%)
2	FOL	A	1[B]	-	14,14,34	2.53	6 (42%)	17,20,47	3.36	10 (58%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FOL	A	1[A]	-	-	-	0/2/2/3
2	FOL	A	1[B]	-	-	-	0/2/2/3

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1[A]	FOL	C4-C4A	11.94	1.61	1.41
2	A	1[A]	FOL	C2-N3	6.50	1.47	1.35
2	A	1[A]	FOL	C4A-N5	5.01	1.40	1.33
2	A	1[B]	FOL	C2-NA2	-4.79	1.24	1.33
2	A	1[B]	FOL	C4-N3	-4.24	1.25	1.33
2	A	1[A]	FOL	C4A-C8A	-4.19	1.33	1.40
2	A	1[A]	FOL	C8A-N1	3.88	1.44	1.36
2	A	1[B]	FOL	C6-N5	3.37	1.40	1.33
2	A	1[A]	FOL	O4-C4	3.27	1.32	1.24
2	A	1[A]	FOL	C6-N5	-3.24	1.27	1.33
2	A	1[B]	FOL	C4A-N5	3.11	1.37	1.33
2	A	1[B]	FOL	C8A-N8	-2.70	1.33	1.37
2	A	1[B]	FOL	C8A-N1	2.22	1.41	1.36

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1[B]	FOL	C4-N3-C2	6.99	127.03	115.93
2	A	1[B]	FOL	C7-N8-C8A	-6.51	110.15	116.69
2	A	1[A]	FOL	C4A-C4-N3	-5.89	115.37	123.43
2	A	1[A]	FOL	N1-C2-N3	-4.91	120.67	127.22
2	A	1[B]	FOL	C4A-C4-N3	-4.78	116.90	123.43
2	A	1[A]	FOL	C9-C6-C7	4.48	130.18	121.42
2	A	1[A]	FOL	C8A-C4A-N5	-4.47	117.26	122.33
2	A	1[B]	FOL	N1-C2-N3	-4.13	121.71	127.22
2	A	1[A]	FOL	C4-N3-C2	4.11	122.45	115.93
2	A	1[A]	FOL	C2-N1-C8A	3.62	119.49	115.36
2	A	1[B]	FOL	NA2-C2-N1	3.62	123.68	117.79
2	A	1[A]	FOL	C7-C6-N5	-3.36	113.35	123.02
2	A	1[B]	FOL	C6-C7-N8	3.28	129.95	121.41
2	A	1[B]	FOL	C6-N5-C4A	-3.25	110.10	118.40
2	A	1[A]	FOL	C4-C4A-N5	3.01	122.04	118.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1[A]	FOL	NA2-C2-N1	2.68	122.17	117.79
2	A	1[B]	FOL	C4-C4A-C8A	-2.66	118.19	119.95
2	A	1[A]	FOL	C6-N5-C4A	2.65	125.17	118.40
2	A	1[A]	FOL	C7-N8-C8A	-2.51	114.17	116.69
2	A	1[B]	FOL	C8A-C4A-N5	2.45	125.12	122.33
2	A	1[A]	FOL	C6-C7-N8	2.23	127.22	121.41
2	A	1[B]	FOL	N8-C8A-N1	-2.05	113.48	115.82

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

2 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1[A]	FOL	3	0
2	A	1[B]	FOL	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand FOL A 1 (A)



Ligand FOL A 1 (B)



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	73:ALA	C	74:LEU	N	1.19

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.